Technical repor Volume 1 – Findings of the review

> Prepared for National Blood Authority

Project Update of Patient Blood Management Guideline for Adults with Critical Bleeding

The Commonwealth of Australia as represented by the National Blood Authority

> prepared by HTANALYSTS Pty Ltd

> > August 2023

Contents

Report information

This evidence evaluation report has been developed by **HT**ANALYSTS in conjunction with the National Blood Authority (NBA) and a Clinical/Consumer Reference Group (CRG). It describes the main body of evidence related to a systematic review of the evidence for the management of people with critical bleeding with regards to blood components and blood conservation strategies. Supplementary data (Appendices A to E) are provided in the Technical Reports (volume 2 and volume 3).

History

The 2011 Patient Blood Management (PBM) Guidelines Module 1: Critical Bleeding/ Massive Transfusion were developed by the NBA (in collaboration with a CRG) to improve patient outcomes; by ensuring that the focus of a patients' medical and surgical management was on improving and conserving the patient's own blood (1). Approval of the 2011 PBM Guidelines: Module 1 was granted by The National Health and Medical Research Council (NHMRC) in 2011 after undergoing public consultation, peer review and an independent AGREE II assessment.

To ensure the 2011 PBM Guidelines: Module 1 reflect the best available evidence, and remain current and relevant for the Australian context, **HT**ANALYSTS were engaged in April 2018 to update the evidence-based recommendations and practice points made in the 2011 PBM Guidelines: Module 1 as informed by a systematic review of the evidence.

NHMRC approval has not been sought for this update, however, all associated materials have been developed in a robust and transparent manner (including public consultation and an independent AGREE II assessment) in accordance with relevant best practice standards (2-5).

Dates

The 2023 Patient blood management guideline for adults with critical bleeding was released on 10 August 2023.

The draft guideline was made available for public consultation from 28 September 2022 to 9 November 2022. All feedback was considered by the CRG and responses to comments recorded at a meeting held 23-24 November 2022.

The evidence review informing the 2023 *Patient blood management guideline for adults with critical bleeding* includes studies published up until 29 September 2021. This technical report (and associated appendices) outlining the best available evidence were presented to the NBA and CRG over several meetings held between December 2021 and August 2022.

The research protocol outlining the methodology to be used to systematically review the evidence to support an update of the *2011 PBM Guidelines: Module 1* received approval from the CRG and NBA on 03 July 2018.

Contents

Funding

The review was funded by the Commonwealth of Australia, as represented by the National Blood Authority.

Authors

¹ HTANALYSTS, Level 8, 46 Kippax Street, Surry Hills New South Wales 2010 Australia

The Research Protocol and the Evidence Evaluation and Technical Reports were written and developed by **HT**ANALYSTS in conjunction with the NBA. Expert advice was provided by the CRG, particularly in relation to development of the research questions, eligibility criteria of identified studies, interpretation of the evidence in relation to clinical important effects and the development of recommendations.

The following named authors (and specific contributions) are outlined in the table below:

Name	Contribution
Dr Margaret	Project Lead and methodological oversight.
Jorgensen	Draft protocol, development of search strategy with contributions from other authors. Oversight of study selection/eligibility, data extraction and data synthesis Meeting
	attendance to facilitate GRADE summary of findings, evidence to decisions and
	development of recommendations.
	Continentation of public consultation reedback.
Alison Miles	Senior Project Manager 2021-2022
	Selection of studies (screening), data extraction and critical appraisal.
	Data synthesis and meeting attendance to facilitate GRADE summary of findings.
Stephanie	Senior Project Manager 2018-2019
Allerdice	Draft methods and development of literature search strategy.
	Selection of studies (screening), data extraction and critical appraisal.
	Data synthesis and meeting attendance to facilitate GRADE summary of findings.
Jessica Shi	2021-2022
	Selection of studies (screening), data extraction and critical appraisal.
	Data synthesis and meeting attendance to facilitate evidence to decisions and
	development of recommendations.
Jack Hide	2021-2022
	Preliminary data extraction and critical appraisal of selected studies.
	Record of meetings, documentation of public consultation feedback.

Acknowledgments

Dr Santwona Baidya, Wendy van Zuijlen, Adrian Peacock, and Kevin Phan are not listed as authors but contributed to the 2018-2019 evidence review (including conduct of the literature search, screening for eligible studies and preliminary data extraction and critical appraisal).

Declarations of interest

All named authors declare they have no financial, personal or professional interests that could be construed to have influenced the conduct or results of the systematic review. Funding, Secretariat and Project Management was provided by the National Blood Authority. The development of the final recommendations was not influenced by the views or interests of the funding body.

Contents

In line with NBA processes, CRG members completed a conflict of interests register before commencement of guideline development as outlined in the NBA conflicts of interest policy. Any additional conflicts of interest were declared at the start of each meetings and appropriately recorded. If a member declared a conflict in relation to a specific intervention (with the exception of conducting research), the member did not participate in the discussion or decision-making for the intervention.

Contents

Re	eport	infor	mation	2
С	onten	ts		5
Li	st of]	Tables	S	8
Li	st of I	Figure	es	
AI	bbrev	iatior	15	
1	Bac	karou	Ind	10
•	11	Docor	intion of condition and sotting	20
	1.1	Desci	intion of intervention and how it might work	
_	1.2	Descr	iption of intervention and now it might work	
2	Rati	onale	and objectives	22
3	Met	hods		23
	3.1	Criter	ia for selecting studies for this review	
		3.1.1	Types of participants	
		3.1.2	Types of interventions	
		3.1.3	Types of outcome measures	
		3.1.4	Types of studies	
	3.2	Searcl	h methods for identification of studies	
		3.2.1	Search terms	
		3.2.2	Databases	
		3.2.3	Other sources	
	3.3	Scree	ning of studies	
		3.3.1	Title/abstract screening	
		3.3.2	Full text screening	
		3.3.3	Screening process	
	3.4	Critica	al appraisal, data collection and evidence synthesis	
		3.4.1	Critical appraisal process	
		3.4.2	Critical appraisal tools	
		3.4.3	Data collection	
		3.4.4	Data synthesis	
	3.5	Sumn	nary of findings and draft recommendations	
		3.5.1	GRADE evidence profiles	
		3.5.2	Evidence to decisions	

	3.6	Conse	nsus process	47
		3.6.1	Consensus guiding principles and values	47
		3.6.2	Consensus ground rules	49
4	Finc	lings	of the systematic review	50
	4.1	Litera	ture search results	50
		4.1.1	Flow of studies	
		4.1.2	Studies awaiting classification or not included	
		4.1.3	Included studies	50
	4.2	Progn	ostic factors (Question 1)	55
		4.2.1	Methods	55
		4.2.2	Summary of evidence	56
		4.2.3	Results	63
	4.3	Define	ed major haemorrhage protocol (Question 2)	76
		4.3.1	Methods	76
		4.3.2	Summary of evidence	77
		4.3.3	Results	85
	4.4	Dose,	timing and ratio (algorithm) of RBC to blood component therapy	
	(Qu	iestion	3)	104
		4.4.1	Methods	104
		4.4.2	Summary of evidence	105
		4.4.3	Results	116
	4.5	Volum	ne of RBC transfused (Question 4)	129
		4.5.1	Methods	129
		4.5.2	Summary of evidence	130
		4.5.3	Results	135
	4.6	Recor	nbinant factor VIIa (Question 5)	140
		4.6.1	Methods	140
		4.6.2	Summary of evidence	
		4.6.3	Results	
	4.7	Blood	components (Question 6)	175
		4.7.1	Methods	175
		4.7.2	Summary of evidence	176
		4.7.3	Results	
	4.8	Antifik	prinolytics (Question /)	242
		4.8.1	Methods	242

4.8.2	Summary of evidence	
4.8.3	Results	
4.9 Viscoe	elastic haemostatic assays (Question 8)	
4.9.1	Methods	
4.9.2	Summary of evidence	279
4.9.3	Results	
4.10 Cell sa	alvage (Question 9)	
4.10.1	Methods	
4.10.2	2 Summary of evidence	320
4.10.3	Results	
5 Reference	es	
Appendix A	- Literature search results	
Appendix B	- Literature screening results	
Appendix C -	- List of excluded studies	
Appendix D	- Critical appraisal	
Appendix E -	- Data extraction forms	

List of Tables

Table 3.1	Types of participants eligible for inclusion	24
Table 3.2	List of eligible interventions	. 25
Table 3.3	Characteristics of the ideal evidence base for each question	. 27
Table 3.4	AMSTAR-2: Domain classification	36
Table 3.5	Evidence to decision framework	44
Table 4.1	Overview of studies identified for each question	51
Table 4.2	Characteristics and quality of included systematic reviews	56
Table 4.3	Characteristics and quality of additional primary studies included in the review	61
Table 4.4	Results for physiologic, biochemical and metabolic (including temperature parameters indicative of critical physiologic derangement: Patients with critical bleeding – Mortality	e) 65
Table 4.5	Results for physiologic, biochemical and metabolic (including temperature parameters indicative of critical physiologic derangement: Patients with critical bleeding – Transfusion volume	e) 72
Table 4.6	Characteristics and quality of systematic reviews by clinical setting: defined MHPs versus no defined MHPs	d 77
Table 4.7	Overlap table primary studies included in the identified systematic reviews defined MHPs versus no defined MHPs	s: 79
Table 4.8	Characteristics and quality of observational and cohort studies by clinical setting: defined MHPs versus no defined MHPs	.80
Table 4.9	Results for defined MHP versus no defined MHP: Patients <i>with</i> critical bleeding – Mortality	88
Table 4.10	Results for defined MHP versus no defined MHP: Patients <i>with</i> critical bleeding – Transfusion volume, red blood cells	93
Table 4.11	Results for defined MHP versus no defined MHP: Patients <i>with</i> critical bleeding – Transfusion volume, fresh frozen plasma	96
Table 4.12	Results for defined MHP versus no defined MHP: Patients <i>with</i> critical bleeding – Transfusion volume, platelets	99
Table 4.13	Results for defined MHP versus no defined MHP: Patients with critical bleeding – Wastage of blood components	102
Table 4.14	Results for defined MHP versus no defined MHP: Patients with critical bleeding – Time to delivery of blood components	103
Table 4.15	Characteristics and quality of systematic review evidence: ratio of RBC to blood component therapy	106

Table 4.16	Overlap table of RCTs identified by included systematic reviews: RBC:FFP and RBC:PLT ratios
Table 4.17	Characteristics and quality of RCT evidence: ratio of RBC to blood component therapy
Table 4.18	Overlap table of observational and cohort primary studies identified by included systematic reviews that meet the 1:1:1 ratio inclusion criterion: RBC:FFP and RBC:PLT ratios
Table 4.19	Overlap table of observational and cohort primary studies identified by included systematic reviews that do not meet the 1:1:1 ratio inclusion criterion: RBC:FFP and RBC:PLT ratios
Table 4.20	Characteristics and quality of observational and cohort evidence: RBC:FFP, RBC:PLT or RBC:CRYO ratio112
Table 4.21	Results for high ratio of blood components versus low ratio of blood components: Patients <i>with</i> critical bleeding – Mortality
Table 4.22	Results for high ratio of blood components versus low ratio of blood components: Patients <i>with</i> critical bleeding – Morbidity, thromboembolic events
Table 4.23	Results for high ratio of blood components versus low ratio of blood components: Patients <i>with</i> critical bleeding – Morbidity, critical complications
Table 4.24	Results for high ratio of blood components versus low ratio of blood components: Patients <i>with</i> critical bleeding – Transfusion volumes, red blood cells
Table 4.25	Results for high ratio of blood components versus low ratio of blood components: Patients <i>with</i> critical bleeding – Transfusion volumes, other blood components*
Table 4.26	Characteristics and quality of systematic review evidence: increased RBC transfusion
Table 4.27	Overlap table showing included systematic reviews and identified cohort studies: increased RBC transfusion131
Table 4.28	Characteristics and quality of prospective cohort evidence: increased RBC transfusion
Table 4.29	Characteristics and quality of retrospective cohort evidence: increased RBC transfusion
Table 4.30	Results for increased volume of RBC transfused versus decreased volume of RBC transfused: Patients <i>at risk of</i> critical bleeding – Mortality
Table 4.31	Results for increased volume of RBC transfused versus decreased volume of RBC transfused: Patients <i>at risk of</i> critical bleeding – Morbidity
Table 4.32	Characteristics and quality of systematic review evidence: rFVIIa142
Table 4.33	Overlap table of RCTs identified by included systematic reviews: rFVIIa

Table 4.34	Overlap table of nonrandomised cohort studies identified by included systematic reviews: rFVIIa
Table 4.35	Characteristics and quality of RCT evidence: rFVIIa
Table 4.36	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – Mortality 151
Table 4.37	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – Morbidity: thromboembolic events
Table 4.38	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – Morbidity: other adverse events
Table 4.39	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – Morbidity: other second-line interventions (obstetrics and maternity only)
Table 4.40	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – RBC transfusion volume
Table 4.41	Results for rFVIIa versus no rFVIIa: Patients <i>with</i> critical bleeding – transfusion volume, other blood components
Table 4.42	Characteristics and quality of systematic reviews by clinical setting: blood components
Table 4.43	Overlap table of RCTs identified by included systematic reviews: blood components
Table 4.44	Overlap table of cohort studies identified by included systematic reviews: blood components
Table 4.45	Characteristics and quality of RCTs by clinical setting: blood components 181
Table 4.46	Characteristics and quality of observational and cohort studies by clinical setting: the effect of blood component therapy on patient outcomes
Table 4.47	Results for FFP versus no FFP: Patients with critical bleeding – Mortality 191
Table 4.48	Results for FFP versus no FFP: Patients with critical bleeding – Morbidity: any adverse outcome
Table 4.49	Results for FFP versus no FFP: Patients with critical bleeding – Transfusion volumes
Table 4.50	Results for FFP versus no FFP: Patients with critical bleeding – Length of stay
Table 4.51	Results for CRYO versus no CRYO: Patients with critical bleeding – Mortality
Table 4.52	Results for CRYO versus no CRYO: Patients with critical bleeding – Morbidity
Table 4.53	Results for CRYO versus no CRYO: Patients with critical bleeding – RBC transfusion volume
Table 4.54	Results for CRYO versus no CRYO: Patients with critical bleeding – Transfusion volume, other blood components

Table 4.55	Results for CRYO versus no CRYO: Patients with critical bleeding – Length of stay
Table 4.56	Results for FC versus no FC: Patients with critical bleeding – Mortality211
Table 4.57	Results for FC versus no FC: Patients with critical bleeding – Morbidity: critical complications
Table 4.58	Results for FC versus no FC: Patients with critical bleeding – Morbidity: ARDS 220
Table 4.59	Results for FC versus no FC: Patients with critical bleeding – RBC transfusion volume
Table 4.60	Results for FC versus no FC: Patients with critical bleeding – Transfusion volume, other blood components
Table 4.61	Results for FC versus no FC: Patients with critical bleeding – Hospital LOS232
Table 4.62	Results of FC versus no FC: Patients with critical bleeding – ICU LOS
Table 4.63	Results for PCC versus no PCC: Patients with critical bleeding – Mortality, latest timepoint
Table 4.64	Results for PCC versus no PCC: Patients with critical bleeding – Morbidity: critical complications
Table 4.65	Results for PCC versus no PCC: Patients with critical bleeding – RBC transfusion volume
Table 4.66	Characteristics and quality of systematic review evidence: Antifibrinolytics versus no antifibrinolytics
Table 4.67	Overlap table of studies identified by included systematic reviews: Antifibrinolytics
Table 4.68	Characteristics and quality of RCT evidence: Antifibrinolytics versus no antifibrinolytics
Table 4.69	Characteristics and quality of cohort evidence: Antifibrinolytics versus no antifibrinolytics
Table 4.70	Results for TXA versus no TXA: Patients with critical bleeding – Mortality254
Table 4.71	Results for TXA versus no TXA: Patients with critical bleeding – Morbidity264
Table 4.72	Results for TXA versus no TXA: Patients with critical bleeding – Blood loss273
Table 4.73	Results for TXA versus no TXA: Patients with critical bleeding – Transfusion volume
Table 4.74	Characteristics and quality of systematic review evidence: TEG or ROTEM versus usual care
Table 4.75	Overlap table of RCTs identified by included systematic reviews: TEG or ROTEM versus usual care
Table 4.76	Overlap table of nonrandomised cohort studies identified by included systematic reviews: TEG or ROTEM versus usual care

Table 4.77	Characteristics and quality of RCT evidence: TEG or ROTEM versus usual care 285
Table 4.78	Characteristics and quality of Observational and cohort studies evidence: TEG or ROTEM versus usual care
Table 4.79	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – Mortality293
Table 4.80	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – Morbidity (thromboembolic events) 299
Table 4.81	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – Major morbidities
Table 4.82	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – RBC transfusion volume
Table 4.83	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – FFP transfusion volume
Table 4.84	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients <i>with</i> critical bleeding – PLT, CRYO, PCC transfusion volume 316
Table 4.85	Characteristics and quality of SR and MA evidence: cell salvage versus no cell salvage
Table 4.86	Overlap table showing systematic reviews and included primary studies: cell salvage versus no cell salvage
Table 4.87	Characteristics and quality of RCT evidence: cell salvage versus no cell salvage
Table 4.88	Characteristics and quality of observation and cohort studies: cell salvage versus no cell salvage
Table 4.89	Results for cell salvage versus no cell salvage: Patients with critical bleeding – Mortality
Table 4.90	Results for cell salvage versus no cell salvage: Patients with critical bleeding – Morbidity: post-operative complications
Table 4.91	Results for cell salvage versus no cell salvage: Patients with critical bleeding – Transfusion volume
Table 4.92	Results for cell salvage versus no cell salvage: Patients with critical bleeding – Cost

Contents

List of Figures

Figure 3.1	Schematic representation of literature review hierarchy	33
Figure 3.2	Consensus Process Flow chart	18
Figure 4.1	Summary of the process used to identify and select studies for the assessment of Question 1 (prognostic factors)	52
Figure 4.2	Summary of the process used to identify and select studies for the assessment of Question 2 (major haemorrhage protocols), Question 3 (ratios of blood components), Question 4 (RBC transfusion volume), and Question 4 (individual blood components)	; 5 53
Figure 4.3	Summary of the process used to identify and select studies for the assessment of Question 5 (rFVIIa), Question 7 (TXA), Question 8 (viscoelastic haemostatic assays) and Question 9 (cell salvage)	54
Figure 4.4	PPO criteria: Question 1 – physiologic, biochemical and metabolic paramete	rs 55
Figure 4.5	PICO criteria: Question 2 – defined MHPs	76
Figure 4.6	Forest plot of comparison: MHPs vs no MHPs, outcome: Mortality, 24 hours.8	36
Figure 4.7	Forest plot of comparison: MHPs vs no MHPs, outcome: Mortality, latest timepoint	37
Figure 4.8	Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, red blood cells	, 72
Figure 4.9	Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, FFP	- 95
Figure 4.10	Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, platelets	, 98
Figure 4.11	PICO criteria: Question 3 – dose, timing and ratio of different ratios of red blood cells)4
Figure 4.12	Forest plot of comparison: high ratio vs low ratio blood components, outcome: Mortality, latest timepoint1	17
Figure 4.13	Forest plot of comparison: high ratio vs low ratio blood components, outcome: Morbidity, thromboembolic events12	20
Figure 4.14	Forest plot of comparison: high ratio vs low ratio blood components, outcome: Morbidity, multiple organ failure1	21
Figure 4.15	Forest plot of comparison: high ratio vs low ratio blood components, outcome: Transfusion volume, red blood cells12	25
Figure 4.16	Forest plot of comparison: high ratio vs low ratio blood components, outcome: Transfusion volume, FFP12	26
Figure 4.17	PPO criteria: Question 4 – effect of transfusion of increased volumes of RBC	29

Figure 4.18	PICO criteria: Question 5 – recombinant factor VIIa140
Figure 4.19	Forest plot of comparison: rFVIIa vs placebo, outcome: Mortality, latest timepoint
Figure 4.20	Forest plot of comparison: rFVIIa vs placebo, outcome: total thromboembolic events
Figure 4.21	Forest plot of comparison: rFVIIa vs placebo, outcome: Other adverse events (trauma setting)
Figure 4.22	Forest plot of comparison: rFVIIa vs placebo, outcome: Morbidity – need for second-line intervention (obstetrics and maternity)164
Figure 4.23	Forest plot of comparison: rFVIIa vs placebo, outcome: Morbidity - other second-line interventions (obstetrics and maternity)
Figure 4.24	Forest plot of comparison: rFVIIa vs placebo, outcome: RBC transfusion volume, Units
Figure 4.25	Forest plot of comparison: rFVIIa vs placebo, outcome: transfusion volume (other blood components), Units172
Figure 4.26	PICO criteria: Question 6 – effect of blood component therapy on patient outcomes
Figure 4.27	Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Mortality, all-cause (at 24 hours)190
Figure 4.28	Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Mortality, all-cause (latest reported timepoint)190
Figure 4.29	Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Morbidity
Figure 4.30	Forest plot of comparison: CRYO vs no CRYO (or varying administration of), outcome: Mortality, latest timepoint
Figure 4.31	Forest plot of comparison: CRYO vs no CRYO (or varying administration of), outcome: Morbidity, thromboembolic events
Figure 4.32	Forest plot of comparison: CRYO vs no CRYO (or varying administration of), outcome: Morbidity, other
Figure 4.33	Forest plot of comparison: FC vs no FC (or varying concentration of), outcome: Mortality, all-cause (latest timepoint)
Figure 4.34	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Morbidity, thromboembolic events
Figure 4.35	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Morbidity, multiple organ failure
Figure 4.36	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: RBC transfusion volume, units
Figure 4.37	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Transfusion volume, other blood components, FFP (trauma)226

Figure 4.38	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Transfusion volume, other blood components, FFP (surgical)226
Figure 4.39	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Length of stay, hospital (days)231
Figure 4.40	Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Length of stay, ICU (days)231
Figure 4.41	Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: Mortality (trauma setting)
Figure 4.42	Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: Morbidity, thromboembolic events (trauma setting)237
Figure 4.43	Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: RBC transfusion volume, Units (trauma setting)239
Figure 4.44	PICO criteria: Question 7 – antifibrinolytics
Figure 4.45	Forest plot of comparison: TXA vs no TXA, outcome: Mortality, latest timepoint
Figure 4.46	Forest plot of comparison: TXA vs no TXA, outcome: Mortality, latest timepoint (trauma only)
Figure 4.47	Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, vascular events (any)
Figure 4.48	Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, venous and arterial events (GI bleeding)
Figure 4.49	Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, other (obstetrics)
Figure 4.50	Forest plot of comparison: TXA vs no TXA, outcome: RBC transfusion volume (trauma)
Figure 4.51	Forest plot of comparison: TXA vs no TXA, outcome: RBC transfusion volume (trauma)
Figure 4.52	PICO criteria: Question 8 – viscoelastic haemostatic assays
Figure 4.53	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: Mortality, latest timepoint
Figure 4.54	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: Mortality, by setting
Figure 4.55	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: thromboembolic events
Figure 4.56	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: morbidity (multiorgan failure, need for hysterectomy)
Figure 4.57	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: RBC transfusion volume (units), by study design

Figure 4.58	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: RBC transfusion volume (units), by setting	05
Figure 4.59	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: FFP transfusion volume (units), by study design	09
Figure 4.60	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: FFP transfusion volume (units), by setting	510
Figure 4.61	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: PLT transfusion volume (units), by study design	314
Figure 4.62	Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: PLT transfusion volume (units), by setting	315
Figure 4.63	PICO criteria: Question 9 – cell salvage	319
Figure 4.64	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Mortality any timepoint up to 30 days	/, 26
Figure 4.65	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Morbidit - post-operative complications	у 29
Figure 4.66	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Morbidit - post-operative complications (urgent AAA repair)	у 30
Figure 4.67	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (RBC)	34
Figure 4.68	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (FFP)	35
Figure 4.69	Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (PLT)	36

Contents

Abbreviations

AAA	Abdominal aortic aneurysm
AGREE	Appraisal of Guidelines for Research and Evaluation
ALI	Acute lung injury
AMSTAR	A MeaSurement Tool to Assess systematic Reviews
AOD	Aortoiliac occlusive disease
ARDS	Acute respiratory distress syndrome
APTT	activated partial thromboplastin time
AUC	Area under the curve
C.A.T.S	Continuous AutoTransfusion System
CRG	Clinical/Consumer Reference Group
CRYO	Cryoprecipitate
BCT	Blood component therapy
DVT	Deep vein thrombosis
ED	Emergency department
EtD	Evidence to Decision
FC	Fibrinogen concentrate
FFP	Fresh frozen plasma
GRADE	Grading of Recommendations Assessment, Development and Evaluation
HSCT	Haematopoietic stem cell transplant
HTA	Health technology assessment
ICU	Intensive care unit
INR	International normalised ratio
IQR	Interquartile range
LOS	Length of stay
MD	Mean difference
MHP	Major haemorrhage protocol
MODS	Multiorgan dysfunction syndrome
MOF	Multiple organ failure
MTP	Massive transfusion protocol
NBA	National Blood Authority
NHMRC	National Health and Medical Research Council
NRSI	Nonrandomised study of intervention
OR	Odds ratio
PBM	Patient blood management
PCC	Prothrombin complex concentrate
pCoh	Prospective cohort study
PE	Pulmonary embolus

PICO	Population, Intervention, Comparator, Outcome
PPH	Primary postpartum haemorrhage
PPO	Population, Prognostic factor, Comparator, Outcomes
PT	Prothrombin time
PLT	Platelets
PT	Prothrombin time
RBC	Red blood cells
rCoh	Retrospective cohort study
RCT	Randomised controlled trial
ROC	Receiver operating characteristic curve
rFVIIa	Recombinant activated factor VII
RR	Risk ratio/Relative risk
ROTEM	Rotational thromboelastometry
SBP	Systolic blood pressure
SC	Single centre
SEM	Standard error of the mean
SoC	Standard of care
SMD	Standardised mean difference
SR	Systematic review
RoB	Risk of bias
TBI	Traumatic brain injury
TEG	Thromboelastography
ТХА	Tranexamic acid
UGIB	Upper gastrointestinal bleeding
VHAs	viscoelastic haemostatic assays

1 Background

The National Blood Authority (NBA), in collaboration with a Clinical/Consumer Reference Group (CRG), has updated the 2011 Patient Blood Management (PBM) Guidelines: Module 1 to ensure it reflects the best available evidence, is current and relevant for the Australian context. Module 1 is part of a series of PBM Guidelines that aims to improve patient outcomes by ensuring that the focus of the patient's medical and surgical management is on improving and conserving the patient's own blood (1).

Based on the best available evidence and knowledge at the time, the 2011 PBM Guidelines: Module 1 made 2 recommendations:

- 1. Institutions should develop a standardised massive transfusion protocol (MTP) that includes the dose, timing, and ratio of blood component therapy for use in trauma patients with, or at risk of, critical bleeding requiring massive transfusion.
- 2. The routine use of rFVIIa in trauma patients with critical bleeding requiring massive transfusion is not recommended, because of its lack of effect of mortality and variable effect on morbidity.

However, no recommendations could be made on:

- the dose, timing, ratio of blood components or use of individual blood components
- the effect of variation of physiologic, biochemical and metabolic parameters on morbidity, mortality and transfusion rate
- the effect of rFVIIa on morbidity, mortality and transfusion rate in patients with critical bleeding.

A review of Module 1 commenced in late 2015 with the establishment of a multidisciplinary CRG. **HT**ANALYSTS were contracted in 2018 to conduct the systematic review of the scientific literature to update or inform new sections of the *2011 PBM Guidelines: Module 1*.

A Research Protocol was then developed to describe the methodology intended to be used to: (i) source the clinical evidence by performing a systematic literature search of the literature, (ii) selecting the best available evidence; (iii) critically appraising and presenting the evidence, and (iv) determining the quality of the evidence base for each question, using a structured assessment of the body of evidence in accordance GRADE¹ methodology.

¹ Grading of Recommendations Assessment, Development and Evaluation. Available at <u>http://gdt.guidelinedevelopment.org/app/handbook/handbook.html</u>

Background

1.1 Description of condition and setting

Critical bleeding is a term used to describe a range of clinical scenarios where bleeding may result in significant patient morbidity or mortality (1). Critical bleeding results in decreased circulating volume, loss of oxygen-carrying capacity, and coagulopathy (impaired clot formation).

Broadly, critical bleeding falls into one of 2 categories (which may overlap):

- major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion (greater than or equal to 5 units of red blood cells in 4 hours (6, 7), or
- 2. haemorrhage of a smaller volume in a critical area or organ (e.g. intracranial, intraspinal or intraocular), resulting in patient morbidity or mortality.

For the purpose of this document, critical bleeding refers only to the first category.

1.2 Description of intervention and how it might work

Different transfusion interventions have been used to restore the circulating blood volume and achieve haemostasis² – 2 key elements of critical bleeding management (1). These interventions include massive transfusion with different combinations or ratios of blood components – such as red blood cells (RBC), fresh frozen plasma (FFP), platelets, cryoprecipitate, fibrinogen concentrate, prothrombin complex concentrate (PCC), whole blood, lyophilised platelets, lyophilised plasma or liquid plasma. Blood component therapy involves the separation of whole blood into specific cellular and plasma components (RBC, FFP, etc.), which are then then transfused separately according to patients' perceived need.

Major haemorrhage is defined based on the volume of blood loss or on the volume transfused. Various definitions exist in the literature and include the loss or transfusion of one blood volume (about 7% of body weight in adults) over 24 hours, or approximately 10 units of RBC (8-11). Alternatively, 'real-time' definitions include replacement of half a blood volume within 4 hours, or blood loss of more than 150 mL per minute (1).

Traditionally, it has been assumed that massive transfusion benefits all patients with critical bleeding. While in certain circumstances such therapy can save lives, a benefit of massive transfusion has not been demonstrable in many clinical scenarios (1). In addition, massive transfusion of blood components is not without risk. Blood transfusion in trauma, surgery and critical care is an independent predictor of multiple organ failure, systemic inflammatory response syndrome, increased infection, and increased mortality (12). It is increasingly clear that the decision to transfuse must be made with great care, and the evidence for the risks and benefits need to be regularly reviewed.

Controversy over the benefits of interventions that aim to improve haemostasis (e.g. FFP, cryoprecipitate, fibrinogen concentrate and PCC) in both surgical and nonsurgical

² a function of balance between procoagulant systems (platelets, coagulation cascade) and anticoagulant systems (APC/protein S, fibrinolysis, serpins).

Background

settings also exist. The use of these interventions may be associated with infection, allergic reactions, haemolysis, transfusion-related circulatory overload and transfusion-related acute lung injury. Monitoring dynamic changes in haemostasis in patients with critical bleeding may help guide transfusion of blood component therapy. Whole blood coagulation analysers are viscoelastic haemostatic assays that may help clinicians assess the cause of bleeding and improve the care of patients with critical bleeding.

Other interventions that may play a role in the management of critical bleeding are tranexamic acid (TXA), recombinant activated factor VII (rFVIIa), and cell salvage.

- TXA acts as an antifibrinolytic by competitively inhibiting the activation of plasminogen to plasmin, a molecule responsible for the degradation of fibrin.
- rFVIIa is a synthetic form of blood factor VIIa, which activates the formation of prothrombinase complex. It is indicated for the treatment or prevention of bleeding in patients with inhibitors to coagulation factor VIII or factor IX, congenital factor VII deficiency and Glanzmann's thrombasthenia.
- Cell salvage is the process that allows blood lost from surgical procedures to be collected, filtered, and washed for re-transfusion to the patient to minimise or prevent allogeneic red cell transfusion.

Objectives

2 Rationale and objectives

The rationale for conducting this review was to update and enhance the evidence and guidance used to inform the 2011 PBM Guidelines: Module 1 (i.e. to identify whether any new high-quality studies had been published and to address the evidence gaps noted). This was to ensure recommendations relating to the appropriate use of blood components and strategies that aim to minimise blood loss in patients who are critically bleeding remain relevant and up-to-date.

In brief, the objectives of the review were to systematically examine the evidence relating to optimisation of blood volume and red blood cell mass, minimisation of blood loss, use of viscoelastic haemostatic assays to assess coagulation function, antifibrinolytics or rFVIIa in patients who are critically bleeding.

The specific questions to be investigated were reviewed and/or developed by the CRG at meetings held in September 2016 and June 2018 and are outline in Box 1 below.

Box 1 Research questions for the update of the 2011 PBM Guidelines: Critical Bleeding

Q1 – In patients with critical bleeding, which physiologic, biochemical and metabolic (including temperature) parameters should be measured early and frequently, and what values of these parameters are indicative of critical physiologic derangement?

Q2 – In patients with critical bleeding, what is the effectiveness of major haemorrhage protocols.

Q3 – In patients with critical bleeding, what is the optimal dose, timing and ratio (algorithm) to RBC, of blood component therapy to reduce morbidity, mortality and transfusion?

Q4 – In patients at risk of critical bleeding, is the transfusion of increased volumes of RBC associated with an increased risk of mortality or adverse effects?

Q5 – In patients with critical bleeding, what is the effect of rFVIIa treatment on morbidity, mortality and transfusion rate?

Q6 – In patients with critical bleeding, what is the effect of FFP, cryoprecipitate, fibrinogen concentrate, PCC and/or platelet transfusion on RBC transfusion and patient outcomes?

Q7 – In patients with critical bleeding, what is the effect of antifibrinolytics on blood loss, RBC transfusion and patient outcomes?

Q8 – In patient with critical bleeding, does the use of viscoelastic haemostatic assays change patient outcomes?

Q9 – In patients with critical bleeding, what is the effect of cell salvage on patient outcomes?

Methods

3 Methods

Methods reported here were based on those described in the <u>Cochrane Handbook for</u> <u>Systematic Reviews of Interventions</u> (13, 14) and relevant sections in the <u>JBI Manual for</u> <u>Evidence Synthesis</u> (15, 16). <u>Covidence</u>, a web-based platform for producing systematic reviews, was used for screening citations and recording decisions made. RevMan 5.4 was used for the main analyses and <u>GRADEPro</u> was used to derive an overall GRADE relating to the certainty of evidence (high, moderate, low, or very low) for each outcome (guided by the <u>GRADE handbook</u>). <u>MAGICApp</u> was then used to record evidence to decisions regarding the development of recommendations.

To identify the evidence for the 9 clinical questions detailed in Box 1, a systematic search of published medical literature was conducted. All potentially relevant studies were identified after applying prespecified inclusion and exclusion criteria as outlined in the research protocol. For eligible studies, the risk of bias was assessed, appropriate data was extracted into data extraction tables, and the results summarised into appropriate categories according to each question. Details on the methods and approach used to conduct the evidence evaluation are provided below.

3.1 Criteria for selecting studies for this review

3.1.1 Types of participants

The types of participants specific to each question are outlined in Table 3.1.

In all questions, the specified population was *people who are critically bleeding*, defined as: *people who have decreased circulating volume*, loss of oxygen-carrying capacity, and coagulopathy due to major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion³.

In question 3, the specific population of interest was people *who received* a massive transfusion.

In question 4, the population included people who were *at risk of* critical bleeding. The broader definition was included to account for patients with penetration injuries who *may* not develop critical bleeding, but if over-transfused before haemorrhage control may go on to do so.

In question 5, the focus was *people who fail to reach adequate haemostasis* and did not include patients with hereditary bleeding disorders such as haemophilia⁴ or those after cardiopulmonary bypass.

In question 9, the focus was on *people in the emergency setting*, and did not include patients in the elective setting.

³ Because the definition of massive transfusion varies across centres, this was to be defined by the literature. ⁴ An X-linked congenital bleeding disorder caused by a deficiency of coagulation factor VIII (FVIII) (in

haemophilia A) or factor IX (FIX) (in haemophilia B). The deficiency is the result of mutations of the respective clotting factor genes. See <u>https://www.blood.gov.au/haemophilia-guidelines</u>

Methods

Question	Types of participants
Q1, Q2 and Q6	People with critical bleeding
Q3	People with critical bleeding Focus: patients who received a massive transfusion
Q4	People <i>at risk</i> of critical bleeding
Q5	People with critical bleeding, Focus: patients who fail to achieve adequate haemostasis despite surgical management and appropriate blood component therapy ^a
Q7	People with critical bleeding Subgroup: patients who have received a massive transfusion
Q8	People with critical bleeding. Subgroups: trauma, obstetrics, perioperative [surgical bleeding, cardiothoracic, liver transplantation], other settings
Q9	People with critical bleeding <i>Focus: patients</i> in the emergency setting (not elective)

Table 3.1 Types of participants eligible for inclusion

Adequate haemostasis is a function of balance between procoagulant systems (platelets, coagulation cascade) and anticoagulant systems (APC/protein S, fibrinolysis, serpins). If haemostasis is out of balance due to a defect in one of these systems, then either thrombosis (too much clotting) or bleeding (not enough clotting) may be the result (17).

Settings: Studies were stratified according to the following patient settings:

- Trauma (meaning physical trauma involving a serious injury to the body). The 2 main types of physical trauma were: (i) blunt force trauma—when an object or force strikes the body, often causing concussions, deep cuts, or broken bones, and (ii) penetrating trauma—when an object pierces the skin or body, usually creating an open wound.⁵
- Surgical (or perioperative meaning around the time of surgery).
 This usually lasts from the time the patient goes into the hospital or doctor's office for surgery until the time the patient goes home⁶. Here, studies were usually in (but were not limited) cardiothoracic or major abdominal surgery, with patients often considered life-threatening, emergency⁷, or urgent⁸.
- *Obstetrics & maternity*: referring to medical care of women during pregnancy and childbirth and in the diagnosis and treatment of diseases of the female reproductive organs⁹.
- *Paediatrics*: referring to medical care of infants (1–23 months of age), children (2–12 years of age), and adolescent (13–18 years of age).

Restrictions: There were no limits to age or ethnicity.

Geographical restrictions: There were no geographical restrictions.

⁵ https://www.nigms.nih.gov/education/pages/Factsheet_Trauma.aspx

⁶ https://www.cancer.gov/publications/dictionaries/cancer-terms/def/perioperative

⁷ Patient requires immediate attention

⁸ Patient requires urgent attention but is not a life-threatening situation

⁹ https://www.cancer.gov/publications/dictionaries/cancer-terms/def/obstetrics-and-gynecology

3.1.2 Types of interventions

Question 1 and Q4 were prognostic questions:

For question 1, studies examining the following parameters as predictors of mortality were eligible for inclusion: temperature, acid-base status, ionised calcium, haemoglobin, platelet count, PT/INR, APTT or fibrinogen level.

For question 4, studies examining the volume of RBC transfused as a predictor for mortality or adverse effects were eligible for inclusion.

Questions 2, 3, 5, 6, 7, 8 and 9 were interventional questions, with eligible studies being those that evaluated the effects of the interventions outlined in Table 3.2. Restrictions on the product type, mode of administration, number of doses or dosage are noted in the table.

Table 3.2	List of	eligible	interventions
-----------	---------	----------	---------------

Question	Intervention
Q2	A defined major haemorrhage protocol
Q3	MHP, RBC:FFP ratio, RBC:PLT ratio, RBC:CRYO ratio
Q5	rFVIIa (as treatment)
Q6	FFP, CRYO, PLT, fibrinogen concentrate, PCC ^a
Q7	Antifibrinolytics (TXA or aprotinin) ^b
Q8	Use of viscoelastic haemostatic assays to guide transfusion of blood component therapy $^{\rm c}$
Q9	Cell salvage

CRYO, cryoprecipitate; FFP, fresh frozen plasma; MHP, major haemorrhage protocol; PCC, prothrombin complex concentrate; PLT, platelets; RBC, red blood cell; rFVIIa, recombinant activated factor seven; ROTEM, thromboelastometry; TEG, thromboelastography; TXA, tranexamic acid

a. Three-factor and four-factor preparations were eligible for inclusion; noting the three-factor preparation is used in Australia/ New Zealand.

b. Limited to IV only (not oral).

c. Limited to TEG or ROTEM (not Sonoclot).

3.1.3 Types of outcome measures

The critical outcome measure to inform decisions on benefits was all-cause mortality reported at 30-days or at the latest measured timepoint. Other measures related to mortality (e.g. death due to bleeding) were also recorded.

The critical outcome measures to inform decisions on harms were related to morbidity. Specifically, any prespecified adverse event relevant to the included population and typically associated with the intervention, such as:

- thromboembolic events¹⁰ (TEs),
- acute respiratory distress syndrome (ARDS),
- time on mechanical ventilator,
- transfusion-related acute lung injury (TRALI),

¹⁰ Inclusive of myocardial infarction, pulmonary embolism and stroke

Methods

- transfusion-associated circulatory overload, and
- multiorgan system failure.

Important or critical outcome measures related to resource use included:

- volume of blood component or blood product transfused (RBC, FFP, PLT, CRYO, fibrinogen concentrate, PCC),
- wastage of blood components,
- time to delivery of blood components, and
- length of hospital or ICU stay.

3.1.4 Types of studies

Eligible studies were those designed to measure a prognostic (Q1 and Q4) or intervention effect (Q2, Q3, Q5, Q6, Q7, Q8, Q9) (18). A summary of the types of studies eligible for each question is provided in Table 3.3.

For prognostic questions, studies with the following design labels were eligible for inclusion¹¹:

- A systematic review of prospective cohort studies (Level I)
- A prospective cohort study (Level II)
- 'All or none' (Level III-1)
- Analysis of prognostic factors among persons in a single arm of an RCT (Level III-2)
- A retrospective cohort study (Level III-3)

For interventional question, studies with the following design labels were eligible for inclusion¹¹:

- A systematic review of RCTs (Level I)
- An RCT (Level II)
- A pseudo (or quasi) RCT (Level III-1)
- A comparative study with concurrent controls including non-randomised, experimental trials, cohort studies, case-control studies and interrupted time series with a control group (Level III-2)
- A comparative study without concurrent controls including historical control studies, 2 or more single arm studies, interrupted time series without a parallel control group (Level III-3).

Level IV evidence¹² was not eligible for any research question, irrespective of whether insufficient higher-level evidence was found to address all critical and important outcomes for that question. This is because results from these studies were likely to lead to misinformed judgements about the effect estimate.

¹¹ <u>https://www.nhmrc.gov.au/ files nhmrc/file/guidelines/developers/nhmrc levels grades evidence 120423.pdf</u>

¹² For prognostic questions Level IV = i.e. case series; for interventional questions Level IV = single arm studies with either post-test, or pre-test and post-test outcomes

Methods

Table 3.3 Characteristics of the ideal evidence base for each question

Question	What type of question is this?	What type of evidence is appropriate?	What size of study is acceptable?	How should the impact of time be considered?	What publication time frame is appropriate?
1	Prognostic	Narrative or systematic review A prospective cohort study Analysis of prognostic factors among persons in a single arm of an RCT A retrospective cohort study Cohort studies of persons with different degrees of bleeding	Any. No restrictions to be applied. Preference to be given to observational studies with at least 500 subjects	Reported as per included studies No restrictions to be applied	January 2009 to current (previous search conducted in June 2009)
2	Interventional	Comparative study with concurrent or historical controls: RCTs Quasi- or pseudo-RCT Cohort study Case-control study Interrupted time series (with or without parallel control group) Include all studies that compare outcomes from the one institution before and after the implementation of a major haemorrhage protocol	Any. Small studies (N≥20 subjects) are acceptable. If observational studies are included, preference is to be given to studies with at least 500 subjects	Reported as per included studies Sensible timeframe for outcomes related to blood component use is 24 h and 48 h; and out to the latest reported measure of mortality	From January 2013 to current (search span for the SR by Mitra 2013 was January 1990 – June 2013)
3	Interventional	SR of RCTs Individual RCTs <i>If little is found, then SR of</i> <i>observational studies can be</i> <i>assessed.</i> Individual observational studies will be excluded ¹³	Any. No restrictions to be applied	Reported as per included studies No restrictions to be applied	From June 2015 to current (search span for SR by Monash was May 2009 ¹⁴ to December 2015)

¹³ CRG noted that evaluating all the available observational studies would not add significantly to the RCT data available however, evidence to support the use of ratios is likely to come from observational studies and that this may be best included by way of the most recent fulsome SR of observational studies (Johansson 2012).

¹⁴ last date of search for current PBM guideline Module 1.

Methods

Question	What type of question is this?	What type of evidence is appropriate?	What size of study is acceptable?	How should the impact of time be considered?	What publication time frame is appropriate?
4	Prognostic	A meta-analysis of interventional and/or observational studies A prospective cohort study Analysis of RBC transfusion volume among persons in a single arm of an RCT A retrospective cohort study Case series, or cohort study of persons with different RBC transfusion volumes Preference to be given to SRs and meta-analyses published by others (i.e. Patel, 2014). Only individual studies published since the Patel 2014 search date to be included.	Any. No restrictions to be applied. If individual observational studies are to be included, preference will be given to studies with at least 500 subjects and/or studies that undertook a multivariate analysis and present adjusted finding.	Reported as per included studies No restrictions to be applied	From January 2012 to current (search span for SR by Patel, 2014 was 1947 – 2012)
5	Interventional	SR of RCTs (with or without meta- analysis) Individual RCT <i>Preference to be given to published</i> <i>SRs and meta-analyses (in broad</i> <i>populations and in specific</i> <i>subpopulations). Inclusion of</i> <i>individual studies only as required.</i> Non-RCTs and observational studies (with concurrent or noncurrent controls) will be excluded.	Any. No restrictions to be applied. For the interpretation of harms, preference will be given to larger studies.	Reported as per included studies No restrictions to be applied	From January 2009 to current (Previous NBA search span was 1990 – June 2009; also provides overlap with relevant Cochrane review (Simpson 2012), and other SRs published in 2011) Note: rFVIIa has been available since the early 2000s
6	Interventional	Individual RCTs Prospective observational studies	Any size RCT. Observational studies with at least 500 participants in total (this is the same restriction applied in the previous SR for this question)	Reported as per included studies Latest reported mortality	For PCC: from 1990 to current For all other interventions: from 2009 to current (Previous NBA search span was 1990 – June 2009, but it did not include PCC)

Methods

Question	What type of question is this?	What type of evidence is appropriate?	What size of study is acceptable?	How should the impact of time be considered?	What publication time frame is appropriate?
		Retrospective observational studies and interrupted time series will be excluded.			
7	Interventional	SR of RCTs RCTs Observational studies Regarding observational studies, preference to be given to prospective studies.	Any size RCT. Observational studies with at least 500 participants	Reported as per included studies Latest reported mortality	From 2000 to current (Cochrane Targeted Update conducted for the NBA in August 2015 relied on a search conducted in January 2015)
8	Interventional	SR of RCTs RCTs Comparative observational studies	Any. No restrictions to be applied. <i>Limited results will be available on this topic.</i> <i>Consider exclusions after the search.</i>	Reported as per included studies Latest reported mortality	From 2000 to current Relevant Cochrane Review (Wikkelsø, 2016) search date January 2016
9	Interventional	SR of RCTs RCTs Cohort studies (prospective or retrospective) Interrupted time series	Any. No restrictions to be applied. For observational studies preference will be given to larger studies	Reported as per included studies Latest reported mortality	From 1990 to current (New question for Module 1)

PCC, prothrombin complex concentrate; RCT, randomised controlled trial; SR, systematic review

Methods

3.2 Search methods for identification of studies

3.2.1 Search terms

The search strategy was developed in Ovid (for Embase and MEDLINE) based on key elements of the research questions (i.e. PICO/PPO criteria). The search strategy was then adapted to suit the Cochrane Library (database of systematic reviews, other reviews, clinical trials, technology assessments, economic evaluations) and PubMed (limited to inprocess citations and citations not indexed in MEDLINE).

Search terms and results for each question are provided in **Appendix A** (see technical report, volume 1).

In developing the search strategy, we appraised and adapted the search strategies provided in the technical report, other health technology assessment reports (including Module 5 and Module 6 of the PBM Guidelines) and those suggested in the scoping report; with terms or concepts proven not suitable removed and other terms added. The overall approach was based on the search methods described in the Technical Report of the *2011 PBM Guidelines: Module 1* (19).

The searches were not limited by outcome, but rather by population, intervention (or prognostic factor), and then study type (applied using the stepped approach outlined above). Methodological filters for identifying different levels of evidence (Level I, Level II, and Level III) developed previously for the *PBM Guidelines* were applied (these filters are based on those developed by NHMRC¹⁵ and SIGN¹⁶) with exclusions for publication types added. The search syntax from embase.com was converted to the Ovid platform.

To facilitate the search and screening of studies, and to minimise duplication of effort, the literature searches were grouped and run under 3 categories:

- Question 1 (prognostic) [screened independently]
- Questions 2, 3, 4, & 6 (blood components or blood products) [screened simultaneously]
- Questions 5, 7, 8, & 9 (blood conservation strategies) [screened simultaneously]

No date, language or geographic limitations were applied when conducting the search:

- Literature search start dates defined by the CRG for each question are provided in Table 3.3. These date limits were applied once citations were imported into the bibliographic management database (EndNote).
- Non-English databases were not searched, however, publications in languages other than English were considered if an English language abstract was available. English language abstracts that appeared to meet the inclusion criteria were supplied to the CRG to confirm if translation in English of the full article was required.
- All studies were considered by the CRG regardless of enrolment country, with a judgement on the applicability of the evidence to the Australian health care

¹⁵ National Health and Medical Research Council

¹⁶ Scottish Intercollegiate Guidelines Network

Methods

context (participant and health system resources) made when assessing the indirectness of the evidence (see Section 3.5).

3.2.2 Databases

In addition to the primary databases listed above (Embase, MedLine, the Cochrane Library and PubMed), searches of additional secondary databases were conducted. This included:

- OpenGrey
- Clinical trial registries (ClinicalTrials.gov and WHO ICTRP¹⁷)
- Health technology assessment/government websites (NICE¹⁸, CADTH¹⁹, and AHRQ²⁰)
- Guideline databases (Guidelines International Network, National Guidelines Clearing House)

3.2.3 Other sources

The review considered both peer reviewed literature, as well as unpublished and grey literature. Studies recommended by CRG members, and potentially relevant studies/systematic reviews identified in the scoping report were also included if they satisfied eligibility criteria and were published within the specified search period of the systematic review.

To maintain the rigour of the systematic review process, studies published after the literature search date of the systematic review were not eligible for inclusion in the technical report. However, pivotal new evidence could be discussed in the guidelines and could be used to inform consensus-based recommendations.

3.3 Screening of studies

3.3.1 Title/abstract screening

Citations (title/abstracts) retrieved by the literature searches for each category were imported into EndNote and duplicates across the databases removed. Citations were then imported into Covidence and screened for inclusion against the eligibility criteria for each question.

At title/abstract stage, one systematic reviewer independently screened each citation who discarded ineligible studies (marked as irrelevant and tagged with a reason for exclusion) and retained potentially eligible ones (marked as relevant or maybe). Where there was uncertainty regarding relevance, a decision was made through discussion with the lead reviewer (MJ), who decided to either mark the citation as irrelevant or take it through for full text review. A second reviewer then checked the screening process to ensure any citation marked as irrelevant did not meet the eligibility criteria. Any differences were resolved by discussion.

¹⁷ World Health Organization International Clinical Trials Registry Platform

¹⁸ National Institute for Health and Care and Excellence

¹⁹ Canadian Agency for Drugs and Technologies in Health

²⁰ Agency for Healthcare Research and Quality

Methods

3.3.2 Full text screening

Full text articles identified for possible inclusion in the evidence synthesis were retrieved then assessed for inclusion independently by 2 reviewers. Where there was uncertainty regarding inclusion, a decision was made through discussion with the lead reviewer (MJ), or advice was sought from the CRG to confirm eligibility based on PICO/PPO criteria.

A prespecified, hierarchical approach was used to annotate reasons for exclusion, with the results of the study selection process illustrated in a PRISMA flow diagram.

Studies were excluded based on hierarchical, prespecified exclusion criteria as follows:

- · Study published prior to search date specified in the protocol
- Duplicate citation
- Non-human study
- PICO out of scope
- Publication type out of scope (e.g. nonsystematic review, editorial, commentary, conference abstract)
- Study type out of scope (e.g. not a comparative clinical study), or
- Other (e.g. study superseded, withdrawn)

Additional prespecified criteria for excluding studies included the following:

- No usable data (systematic review does not provide data relating to the primary studies)
- Primary study (RCT and/or nonrandomised study) already assessed and included in a systematic review
- Sample size (as specified in the protocol for each question)
- · Insufficient or no adjustment for confounders (observational studies only)

Trial registration numbers, author names and study titles, locations and dates were used to identify multiple citations arising from the same study. Ongoing trials and studies published as abstracts were also identified and are listed under "Studies awaiting classification".

3.3.3 Screening process

To minimises the potential for bias, a hierarchical approach to the screening for each question was conducted as illustrated in Figure 3.1.

Using a stepped process, the highest 'level' of evidence was assessed before studies with other design labels were considered. This meant that a systematic review of Level II studies was considered the highest level of evidence (Level I) for all question types (see section 3.1.4 for study design labels).

If high-quality (see Section 3.4.2) systematic review evidence was available to address the specified outcomes of interest, assessment of studies with other design labels (Level II or Level III) was not conducted²¹.

²¹ Noting that eligible RCTs (or Level II studies) published since the search date of the key systematic review were identified and incorporated into the review.

Methods

If there were no relevant systematic reviews available for a specific research question, the citations retrieved from the Level II²² search were screened, and if no Level II studies were identified the process was repeated for Level III studies (to the level specified in the PICO/PPO criteria).

For critical and important outcomes not addressed in higher-level evidence, the screening of lower-level evidence was targeted to that outcome only.

Where there was insufficient or no evidence available to answer a research question, a 'consensus recommendation' or 'good practice statement' was made.



Figure 3.1 Schematic representation of literature review hierarchy

In 2018, for each group of questions, all citations identified in the search for systematic reviews (meta-analyses, guidelines etc.) were screened according to the date limits indicated in Table 3.3. Date limits specific to each question were applied within EndNote.

Systematic reviews and meta-analyses identified for potential inclusion were scrutinised and assessed for eligible primary studies (RCTs and/or nonrandomised studies) for each question. Any systematic review that provided duplicate information (duplicate data) or did not provide enough information about the primary studies (no usable data) was excluded, with the most comprehensive and most recent systematic review retained. Reviews that did not include any eligible primary studies were excluded.

Based on the literature search dates of the most recent systematic review, a date limit was then applied to Level II and PubMed searches, which was then screened for additional studies not already identified.

At this point), a list of potentially relevant studies, and the existing clinical questions to which they applied, was supplied to the NBA and CRG with an understanding of the

²² For prognostic questions Level II = prospective cohort studies; for interventional questions = RCTs

Methods

scope of new evidence to undergo full critical appraisal and data extraction reached before proceeding to the next stage of this review (either continue screening for RCTs and observational searches or move on the critical appraisal, data collection and evidence synthesis).

Here, the CRG was consulted to advise whether the identified evidence would likely answer and address each research question, and whether the inclusion of additional lower-level evidence would likely substantially change the overall results.

Specifically, each question custodian was asked to respond to the following 3 questions:

- What is missing from the list of identified primary studies?
- Are there any 'landmark' studies that are not included in the proposed included studies?
- If there are no 'landmark' studies that would capture the missing data or they are insufficient to answer the question or the missing data is unknown, should any of the following options be considered?
 - Search the Level II and/or Level III evidence bases for studies published after publication of the most recent systematic review literature.
 - Conduct a targeted review focusing only on components not included in the systematic review evidence. What are these components?
 - o Other?

An agreement to stop screening was reached in December 2018 after all relevant systematic reviews had been identified (and the included primary studies) and any pivotal new studies had been included. While some CRG members noted the paucity of systematic review evidence, they acknowledged the considerable work required to investigate primary studies may not be justified.

The searches were re-run in 2019 and again in 2021, with the stepped process again used when screening the body of evidence. This occurred after the application of date limits that incorporated a minimum 6 months prior to the previous search date (see Appendix A, technical report, volume 2).

In 2019, citations retrieved in the search for systematic reviews and published between January 2018 to August 2019 were screened; however due to a pause in the project in 2020, screening of citations retrieved in the lower-level searches did not proceed.

In 2021, citations retrieved in the Level I search (systematic reviews) and published between January 2019 to September 2021 were screened, followed by the application of date limits according to the most recent and comprehensive systematic review identified for each question. A new date limit was then applied to remaining searches (Level II, Level III, PubMed), which were screened for additional studies published after the systematic review search date. Methods

3.4 Critical appraisal, data collection and evidence synthesis

3.4.1 Critical appraisal process

The methodological quality of included systematic reviews and the of risk of bias of primary studies was assessed using a variety of assessment tools according to the type of study. Here, the clarity and completeness or reporting, strengths and weaknesses of methods and processes used, as well as the underlying assumptions and limitations of a study was assessed. For each systematic review or primary study, supporting information and a rationale for each judgement is provided in **Appendix D** (see technical report, volume 2).

Critical appraisal of each included systematic review or primary study was assessed by one reviewer. A second reviewer then checked and confirmed each assessment made. Disagreements were resolved through discussion, with advice sought from a third reviewer if needed.

3.4.2 Critical appraisal tools

3.4.2.1 Systematic reviews and meta-analyses

Systematic reviews and meta-analysis of RCTs and/or observational studies were assessed using the AMSTAR-2 quality assessment checklist (20). The AMSTAR-2 consists of 16 domain questions (classified as critical flaws or weaknesses as outlined in Table 3.4) that are answered as 'yes', 'no', 'partial yes'. A 'yes' answer denotes a positive result.

The overall quality of the systematic review was summarised based on the criteria outlined in Box 2.

Prior to 2019, systematic reviews and meta-analyses of RCTs were assessed using the AMSTAR²³ quality assessment checklist (21), which consists of 11 signalling questions that are answered as 'yes', 'no', 'can't answer', or 'not applicable'. A 'yes' answer denotes a positive result.

The overall quality of systematic reviews assessed with the original AMSTAR checklist was summarised based on the following criteria: (i) high quality, scoring 'yes' on 9 or more questions, (ii) moderate quality, scoring 'yes' on between 6 and 8 questions, and (iii) low quality, scoring 'yes' on 5 or less questions.

It is noted that AMSTAR and AMSTAR-2 lead to a judgement of methodological quality (or limitations) of a systematic review, not a judgement about risk of bias of the body of evidence included within the systematic review. The risk of bias of primary studies included within the systematic review (if reported) were noted during data collection (see **Appendix E** [technical report, volume 3]).

²³ A MeaSurement Tool to Assess systematic Reviews

Methods

Table 3.4 AMSTAR-2: Domain classification

Critical weakness	Critical flaw
Domain 1: Inclusion of PICO in research	Domain 4: Adequacy of the literature search
questions and inclusion criteria	Domain 8: Detailed description of included
Domain 2: Registration of protocol before	studies
commencement of the review	Domain 9: Risk of bias from individual studies
Domain 3: Discussion of selection of study	being included in the review
designs for inclusion	Domain 11: Appropriateness of meta-analytical
Domain 5: Duplicate study selection	methods
Domain 6: Duplicate data extraction	Domain 13: Consideration of risk of bias when
Domain 7: Justification for excluding individual	interpreting the results of the review
studies	
Domain 10: Review of sources of funding for	
included studies	
Domain 12: Discussion of impact of risk of bias	
of included studies on meta-analysis results	
Domain 14: Discussion of heterogeneity	
Domain 15: Assessment of presence and likely	
impact of publication bias	
Domain 16: Reporting of potential sources of	
conflict of interest including any funding	
received	

Source: (20)

Box 2 Overall quality of included systematic reviews

Overall quality of included systematic reviews

High quality (no or one noncritical weakness) – the systematic review provides an accurate and comprehensive summary of the results of the available studies that address the question of interest.

Moderate quality (more than one noncritical weakness) – the systematic review has more than one weakness but no critical flaws. It may provide an accurate summary of the results of the available studies that were included in the review.

Low quality (one critical flaw with or without noncritical weaknesses) – the review has a critical flaw and may not provide an accurate and comprehensive summary of the available studies that address the question of interest.

Critically low quality (more than one critical flaw with or without noncritical weaknesses) – the review has more than one critical flaw and should not be relied on to provide an accurate and comprehensive summary of the available studies.
Methods

3.4.2.2 Randomised controlled trials

The risk of bias of included RCTs was assessed using the Cochrane Collaboration's Risk of Bias tool (22). This tool is made up of 6 bias domains assessing 7 sources of bias including selection bias (random sequence generation and allocation concealment), performance bias (blinding of researchers and patients), detection bias (blinding of outcome assessment), attrition bias (incomplete outcome data), reporting bias (selective reporting) and other bias.

For each domain, concerns of bias were raised (recorded as 'high', 'low', or 'unclear') when it was considered plausible (i.e. likely, probable, possible or conceivable) that bias was present. Supporting information and a rationale for each judgement is provided in **Appendix D** (see technical report, volume 2).

The overall risk of bias for an RCT was determined based on the criteria outlined in Box 3.

Box 3 Overall risk of bias within identified RCTs

Overall risk of bias within identified RCTs

Overall low risk – low risk of bias for ALL key domains

Overall unclear risk - low or unclear risk of bias for ALL key domains

Overall high risk – high risk of bias for one or more key domains

3.4.2.3 Observational (nonrandomised) cohort studies

The risk of bias of observational cohort studies was guided by GRADE²⁴, with the focus being on bias relating to the following 4 domains: selection of participants, measurement of exposure/outcomes, confounding and follow-up.

It is noted that formal risk of bias assessment of cohort studies using the Cochrane Collaboration's ROBINS-I (or <u>another appropriate tool</u>) was not conducted.

For each domain, concerns of bias were raised (recorded as 'low', 'moderate', 'serious', 'critical', or 'no information provided') when it was considered plausible (i.e. likely, probable, possible or conceivable) that bias was present. Supporting information and a rationale for each judgement is provided in **Appendix D** (see technical report, volume 2).

The overall risk of bias for observational studies was determined based on the guide outlined in Box 4.

²⁴ Table 5.5 in GRADE handbook http://gdt.guidelinedevelopment.org/app/handbook/handbook.html#h.m9385o5z3li7

Methods

Box 4 Overall risk of bias within identified observational (cohort) studies

Overall risk of bias within identified observational (cohort) studies

Overall low risk of bias – the study is comparable to a well-performed RCT and is judged to be a low risk of bias for ALL domains

Overall moderate risk of bias – the study appears to provide sound evidence for a nonrandomised study but cannot be considered comparable to a well-performed randomised trial. The study is judged to be a low or moderate risk of bias for ALL domains

Overall serious risk of bias – the study has some important problems and is judged to be at serious risk of bias *in* at least ONE domain, but not a critical risk of bias in any domain

Overall critical risk of bias – the study is too problematic with regards to this domain to provide any useful evidence. The study is judged to be at critical risk of bias in at least ONE domain

No information – there is no information on which to base a judgement about overall risk of bias. There is no clear indication that the study is at serious or critical risk of bias AND there is a lack of information in one or more key domains of bias.

3.4.3 Data collection

The characteristics and results of each included systematic review or primary study were extracted by a single evidence reviewer using standardised data collection forms (see **Appendix E** [technical report, volume 3]). Data extraction forms were then checked by a second reviewer, with any disagreements resolved through discussion.

The following characteristics of included studies was extracted:

- study design
- year conducted
- funding sources and funder involvement in study
- setting and location (such as prehospital, trauma setting, military zone)
- participant characteristics (including enrolment number and any notable demographics or comorbidities)
- intervention and comparator characteristics (including product, timing, dose and administration technique)
- outcomes measure and results (including measurement method, timing or severity)

Only data from systematic reviews judged to be of high or moderate quality were used to inform the evidence base. Here, data was extracted from the systematic review or metaanalyses and a return to the source documents (primary studies) to verify data was not done. A return to source documents occurred if their where concerns about the completeness of data reported in the systematic review. Methods

3.4.4 Data synthesis

After data collection, the available effect estimates (including 95% confidence intervals, *p*-values) for critical and important outcomes and those relating to resource use were presented in evidence summary tables, alongside the population and intervention characteristics. The evidence summary tables were structured by question, comparisons, study design and outcome measure (see results tables in Section 4). All available information was reported, including if the results were incompletely reported (e.g. no effect estimate, but the direction of effect with a *p*-value was reported). Implications of the missing outcome data were considered when interpreting the evidence (see Section 3.5).

Where possible, data synthesis of results within each comparison was performed²⁵ according to methods described in Chapter 6 of the Cochrane Handbook (23). Using RevMan 5.4, effect estimates were combined across studies for each outcome using a random effects model, with data from RCTs and observational studies presented separately. Forest plots were used to visually depict the results. If the reported information allowed for direct calculation of effect estimates or imputation of missing statistics (e.g. standard deviations), calculations were performed within the computer program²⁶ (23).

Heterogeneity was assessed by visually by inspecting the overlap of confidence intervals on the forest plots, formally test for heterogeneity using the Chi² test (using a significance level of α = 0.1), and quantify heterogeneity using the l² statistic (24).

Indirect treatment comparisons were not conducted.

3.5 Summary of findings and draft recommendations

For each comparison, setting and outcome, the available evidence was assessed using the GRADE approach (25), which provides a framework for rating the certainty of the evidence for each outcome (see Box 5) and grading recommendations in health care (see Box 6).

3.5.1 GRADE evidence profiles

For each question, evidence profiles were initially developed by the lead reviewer (MJ), using the GRADEpro GDT software (<u>www.gradepro.org</u>), with the CRG then considering each profile and relevance to the Australian context. In the absence of data, a narrative summary was provided. All critical and important outcomes were reported, regardless of whether the findings demonstrate a clinically meaningful change.

²⁵ i.e. the PICO criteria and study design features were considered sufficiently homogenous or suitable to be combined.

²⁶ Usually transformed from published confidence intervals or standard errors of the mean

Methods

Box 5 GRADE certainty of evidence (per outcome)

GRADE certainty of evidence

High $(\oplus \oplus \oplus \oplus)$: further research is very unlikely to change the confidence in the estimate of effect

Moderate $(\oplus \oplus \oplus \ominus)$: further research is likely to have an important impact in the confidence in the estimate of effect

Low $(\oplus \oplus \ominus \ominus)$: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate

Very low $(\oplus \ominus \ominus \ominus)$: any estimate of effect is very uncertain

Box 6 GRADE Recommendations



Methods

Beginning with the study design (RCTs or observational studies), 5 factors were considered that can reduce the certainty of evidence and 3 factors were considered that can increase the certainty of evidence (see Box 7). Here, scoring of the certainty of the evidence begins as 'high' for RCTs (score=4), and 'low' for observational studies (score=2).

For each outcome, a judgement was recorded against each factor that could reduce the certainty of evidence (no concerns, serious or very serious). Each factor was downgraded by –1 for serious concerns or –2 for very serious concerns. Footnotes were used to record judgements made about downgrading or upgrading the evidence. Factors that can increase the certainty of evidence were considered only where relevant.

Box 7 GRADE factors that can reduce or increase the certainty of the evidence

Factors that can reduce the certainty of the evidence

Risk of bias. Based on the summary risk of bias assessment across studies for each outcome reported for a comparison (26).

Inconsistency. Based on heterogeneity in the observed effects across studies that suggests important differences in the effect of the intervention and whether this can be explained (27).

Imprecision. Based on interpretation of the upper and lower confidence limits of the pooled result and whether the intervention has a clinically important effect (28).

Indirectness. Based on important differences between the review questions and the characteristics of included studies that may lead to important differences in the intervention effects (29).

Publication bias. Based on the extent to which the evidence is available and the likely non-reporting of results (30).

Factors that can increase the certainty of the evidence

Large magnitude of effect

All plausible confounding would reduce the demonstrated effect or increase the effect if no effect was observed

Dose-response gradient

Risk of bias

For GRADE assessments it was necessary to first draw conclusions about the overall risk of bias for each outcome within a study, and then summarise risk of bias assessments across studies for each outcome. These summary assessments of risk of bias were used in determining the overall certainty of evidence, and the basis for each was reported as footnotes to the GRADE summary of findings tables.

Methods

Serious concerns were raised if the outcome result was influenced by the inclusion of studies judged to be at high risk of bias (i.e. removing these studies changed the size of the effect). Serious concerns were also raised if it was considered plausible (i.e. likely, probable or conceivable) that missing outcome data made a difference to the estimated effect (considering the weight of studies that had substantial missing data).

Inconsistency

For GRADE assessments we considered measures of statistical heterogeneity (e.g. I² statistic) as well as any non-overlap of confidence intervals that could not be explained, suggesting important difference in the observed effect.

Inconsistency was not downgraded when there was only one study.

Imprecision

For GRADE assessments we considered the upper and lower confidence limits of the pooled result in relation to a minimal clinically important threshold (i.e. the confidence interval includes both appreciable benefit and harm); and whether the optimal information size has been reached (i.e. the total number of patients meets the required sample size for a sufficiently powered individual study).

In determining the clinical importance, a rough threshold guide was used: for dichotomous outcomes a 25% relative risk reduction (or increase); for continuous outcomes we used Cohen's guidance (31) for interpreting the magnitude of the SMD: 0.2 represents a small difference, 0.5 is moderate, and 0.8 is a large difference.

Indirectness

Studies were downgraded for indirectness if a large proportion of participants included in the study did not meet the definition for critical bleeding or when clinical decisions relating to transfusion of blood components or blood conservation strategies differed to that typically used in Australian practice.

Publication bias.

Judgements regarding missing results across the identified studies were made based an assessment of 'known-unknowns' (i.e. selective non-reporting or non-inclusion of results from identified studies). This included checking for missing outcome results in published studies, checking the ongoing studies and studies awaiting classification and making a judgement on whether the studies were not complete, failed to report an outcome, were not published (or translated) due to the nature of their results (e.g. results were in favour of the comparator, or no observed effect) and if the missing result for the outcome would materially influence the meta-analysis results. Given most of the outcome results came from small studies, any missing results due to non-reporting was considered likely to impact the results.

A judgement about 'unknown-unknowns' was made based on the likelihood that missing data from studies not identified was likely to have included that outcome. Publication bias was suspected when the evidence for an outcome was limited to a small number of small trials. all reporting a positive effect. No additional statistical analysis for testing for small-study effects (e.g. contour enhanced funnel plots) was conducted.

Methods

3.5.2 Evidence to decisions

GRADE evidence profiles were transitioned into MAGICApp and summary of findings tables reporting estimates of treatment effects for each outcome as absolute and relative risks were presented and discussed with the CRG. Here, an evidence statement pertaining to each outcome was included. The evidence statement was guided by the format prescribed in MAGICApp (32).

The evidence to decisions framework provided within MAGICApp was used to inform translation of the evidence into recommendations for use in the clinical guidance chapter.

Recommendations were made after considering the following key concepts (see Table 3.5):

- Benefits and harms
- · Certainty of evidence
- Values and preferences
- · Resources
- Equity
- · Acceptability
- · Feasibility

As noted by GRADE (33):

"In the context of a systematic review, the ratings of the quality of evidence reflect the extent of our confidence that the estimates of the effect are correct. In the context of making recommendations, the quality ratings reflect the extent of our confidence that the estimates of an effect are adequate to support a particular decision or recommendation."

As such, the certainty of the evidence was used to inform the strength of any evidencebased recommendations that are made, with higher certainty evidence resulting in a strong recommendation for or against a particular action, and lower certainty resulting in a weak or conditional recommendation for or against a particular action.

A consensus process (see Section 3.6) was used to ensure the clinical guidance is consistent with the evidence presented. Any dissenting opinions regarding the wording or grading of recommendations was documented.

Methods

Table 3.5 Evidence to decision framework

Decision domain	Questions to consider	Ratings (from MAGICapp)	Summary of judgement (from MAGICapp)
Benefits and harms <i>A narrative summary of the most</i> <i>important benefits and harms</i>	How substantial are the benefits? How substantial are the harms? Are you confident that the benefits outweigh the harms or burden for most patients? Is the baseline risk similar across different patients? Should there be separate recommendations for different patients?	Not set Trivial/no benefits Small benefits Moderate benefits Large benefits Varies Don't know n/a	Not set Small net benefit, or little difference between alternatives Substantial net benefit of the recommended alternative Important harms
Certainty of evidence <i>Overall certainty in effect estimates across</i> <i>outcomes</i>	What is the certainty of the evidence?	Not set Very low Low Moderate High n/a	Not set Very low Low Moderate High
Values and preferences Typical patient preferences and values? Common issues or expected variability?	Considering values and preference of patients and their carer's, are you confident the benefits outweigh the harms and burdens for most patients?		Not set Substantial variability is expected or uncertain No substantial variability expected We expect few to want the intervention
Resources Issues with costs or resource use? For whom? Cost benefit analysis? Implementation or other issues?	Do the resources used (including cost, personnel time, etc.) favour the intervention or the comparator?	Not set Large savings Moderate savings Negligible cost or savings Moderate cost	Not set Factor not considered Important issues or potential issues not investigated

Methods

Decision domain	Questions to consider	Ratings (from MAGICapp)	Summary of judgement (from MAGICapp)
		Large cost Varies Don't know n/a	No important issues with the recommended alternative Important negative issues
Equity How do the different alternatives affect equity?	Are there reasons for anticipating differences in the effectiveness of the intervention for any disadvantaged patients or settings (e.g. place of residence, ethnicity, education, socioeconomic status)? Is the recommendation likely to reduce existing inequity? Are there different baseline conditions across patients or settings, that would change the absolute effectiveness of the intervention, or is the issue more/less important, for any disadvantaged patients or settings? Are there important considerations that should be made when implementing the option to ensure inequities are not increased?	Not set Increased equity Probably increased equity Probably no impact Probably reduced equity Reduced equity Varies Don't know n/a	Not set Factor not considered Important issues or potential issues not investigated No important issues with the recommended alternative Intervention likely increases inequity
Acceptability Is the option acceptable to key stakeholders?	Are there key stakeholders (e.g. private and public hospitals, consumers and carer's, health professions, Hospitals and Health Services, state health departments) that would not accept the distribution of the benefits, harms and costs? Are there key stakeholders that would not accept the costs or undesirable effects in the short term for desirable effects (benefits) in the future? Are there stakeholders that would put more value (relative importance) on the undesirable consequences than the desirable consequences or costs?	Not set Acceptable Probably acceptable Probably not acceptable Not acceptable Varies Don't know n/a	Not set Factor not considered Important issues or potential issues not investigated No important issues with the recommended alternative Intervention is likely poorly accepted

Methods

Decision domain	Questions to consider	Ratings (from MAGICapp)	Summary of judgement (from MAGICapp)
	Are there key stakeholders that would disapprove of the intervention morally, i.e. in relationship to ethical principles such as autonomy, nonmaleficence, beneficence or justice?		
Feasibility How feasible will it be to implement the different alternatives? Any issues?	Is there lots of variability in the resource requirements across settings? Is the intervention feasible to implement for either patients and their carer's or for health professionals?	Not set Feasible Probably feasible Probably not feasible Not feasible Varies Don't know n/a	Not set Factor not considered Important issues or potential issues not investigated No important issues with the recommended alternative Intervention is likely

Source: (34, 35)

Methods

3.6 Consensus process

The consensus process for developing evidence-based recommendation and expert opinion points is illustrated in Figure 3.2:

- *Stage 1 Introduction.* The Chair describes the consensus process, participants' roles and responsibilities, ground rules and the guiding principles.
- Stage 2 Open discussion. The Chair opens the floor to a general discussion and suggestions for recommendation / expert opinion wording, noting that recommendations will be based on the GRADE framework. The Chair provides an opportunity for concerns or issues to be raised.
- Stage 3 Resolve concerns. The Chair has the first option to resolve concerns by clarifying or changing the wording, or seeing whether those with concerns will stand aside. Where concerns are not resolved and the time is short, the discussion will be carried over to a later meeting.
- Stage 4 First call for consensus. The Chair calls for consensus.
- Stage 5 Second call for consensus. If consensus is not reached, the CRG will consider the consensus process guiding principles and values, and:
 - o the member stands aside, and the differing schools of thought are documented
 - the member is not willing to withdraw the concern or stand aside, and the CRG declares itself blocked – the recommendation or expert opinion is not accepted
 - o the member withdraws their concern and consensus is reached

3.6.1 Consensus guiding principles and values

- Consensus is reached where all members agree with the recommendation / expert opinion point. Consensus is not achieved on the basis of a 'majority'.
- The opinions of all members of the group are equally valid/important, notwithstanding that some members may have discipline-specific expert opinion.
- Where consensus is not reached, the dissenting members may present their case. This may be done immediately in the current meeting or be carried over to the subsequent meeting to allow the members to succinctly formulate their concerns or provide other documentation/ research.
- Issues of semantics, language or content, while recognised as important, should preferably not absorb discussion time within the meetings.
- Members are respectfully asked to reflect upon their own values and conflicts of interests and be mindful of the extent to which these may influence their opinions.

Methods



Figure 3.2 Consensus Process Flow chart

Methods

3.6.2 Consensus ground rules

- Members agree to take turns speaking and not interrupt each other.
- Members agree to stay away from establishing hard positions or express themselves in terms of personal needs and interests and the outcomes that they wish to realise.
- Members recognise that, even if they do not agree with it, each of them is entitled to their own perspective.
- Members will not dwell on things that did not work in the past, but instead will focus on the future they would like to create.
- Members agree to make a conscious, sincere effort to refrain from unproductive arguing, venting, or narration, and agree to use their time to work towards what they perceive to be their fairest and most constructive agreement possible.
- Members will speak up if something is not working for them during the consensus process.
- Members will request a break when they need to.
- Members will point out if they feel the Chair is not being impartial.

4 Findings of the systematic review

4.1 Literature search results

4.1.1 Flow of studies

The medical literature was searched on 11 August 2018 to identify relevant systematic reviews and primary studies published from database inception to the literature search date. The searches were repeated on 09 August 2019 and again on 29 September 2021²⁷ to ensure the most recent and relevant evidence had been identified to inform clinical guidance.

Searches were conducted using the databases and sources described in Section 3.2, with citations returned by the literature searches screened based on information in the publication title and abstract using a stepped process as described in Section 3.3.

Search terms and search results are described in **Appendix A** (see technical report, volume 2). Details on the application of the study selection criteria are provided in **Appendix B** (see technical report, volume 2).

A PRISMA flow summarising the number of studies at each stage of the search and screening process for Question 1 is shown in Figure 4.1.

A PRISMA flow summarising the number of studies at each stage of the search and screening process for Questions 2, 3, 4, and 6 is shown in Figure 4.2.

A PRISMA flow summarising the number of studies at each stage of the search and screening process for Questions 5, 7, 8, and 9 is shown in Figure 4.3.

4.1.2 Studies awaiting classification or not included

No language limits were applied to the search strategy, however eligible studies published in a language other than English were not included. These studies, and other studies that could not be retrieved or those that met the inclusion criteria but contained insufficient or inadequate data for inclusion are listed in **Appendix C** (see technical report, volume 2).

4.1.3 Included studies

Overall, the systematic review and handsearching process identified 73 systematic reviews (that had assessed 156 eligible primary studies) and 23 additional primary studies covering the 9 research questions.

An overview of the number studies that informed the evidence is provided in Table 4.1. Details are provided within the summary of evidence section relating to each intervention.

²⁷ Due to unforeseen challenges and delays (including COVID).

Intervention	SRs (k)	Additional RCTs	Additional NRSIs	Total evidence base	Section
Physiological parameters	12 (k=50) ª	0	3 pCoh 4 rCoh ^b 2 RCT analysis	59 primary studies	4.2.2
Major haemorrhage protocol	8 (k=5 pCoh, 24 rCoh)	0	0	5 pCoh 24 rCoh	4.3.2
RBC ratios, timing and dose	16 (k=12 RCTs, 2 pCoh, 20 rCoh)	0	0	12 RCTs 2 pCoh 20 rCoh	4.4.2
RBC transfusion volume	2 (k=9 pCoh, 12 rCoh)	0	1 pCoh 1 rCoh	10 pCoh 13 rCoh	4.5.2
Recombinant activated factor VII	8 (k= 9 RCTs)	2	Not eligible	11 RCTs	4.6.2
Blood components and/or products	11 (k=15 RCTs, 3 pCoh, 13 rCoh)	0	1 rCoh	15 RCTs 3 pCoh 14 rCoh	4.7.2
Antifibrinolytics	13 (k=5 RCTs, 4 pCoh, 12 rCoh)	1	0	5 RCTs 4 pCoh 12 rCoh	4.8.2
Viscoelastic haemostatic assays	12 (k=6 RCTs, 1 pCoh, 13 rCoh)	1	1rCoh	7 RCTs 1 pCoh 14 rCoh	4.9.2
Cell salvage	3 (k=1 RCT, 2 pCoh, 3 rCoh)	0	1 pCoh	1 RCT 3 pCoh 3 rCoh	4.10.2

Table 4.1 Overview of studies identified for each question

Coh, observational or cohort study; pCoh, prospective cohort; rCoh, retrospective cohort; NRSI, nonrandomised study of an intervention; RCT, randomised controlled trial; SRs, systematic reviews

k = number of eligible primary studies included within the SRs

a. Study design features of primary studies included within the SRs were not always specified. Includes prospective cohort studies and retrospective analyses with before and after design.

b. Retrospective cohort studies are inclusive of before and after studies that include an historical control.





a. Search for SRs, RCTs, and cohort studies conducted via Ovid (Embase, MEDLINE, EBM Reviews) and PubMed (in-process and citations not indexed in MEDLINE).

b. Protocol date limits as follows: Q1 SRs – studies published prior to 2009; Q1 RCTs, cohort, PubMed – studies published prior to 2016 (initial search); Q1 (updated search) – studies published prior to 2019.

Figure 4.2 Summary of the process used to identify and select studies for the assessment of Question 2 (major haemorrhage protocols), Question 3 (ratios of blood components), Question 4 (RBC transfusion volume), and Question 6 (individual blood components)



- Search for Level I studies conducted via Ovid (Embase, Medline, EBM Reviews) and PubMed (in-process and citations not indexed in Medline), where possible date limits from 2019 (not applied to PubMed), restricted to Q2, Q3, Q4 and Q6.
- b. Search for Level II studies conducted via Ovid (Embase, Medline, EBM Reviews) and PubMed (in-process and citations not indexed in Medline), where possible date limits from 2019 (not applied to PubMed), restricted to Q2, Q3, Q4 and Q6.
- c. Studies outside data limits and/or were not tagged as either level I or level II study design.

Figure 4.3 Summary of the process used to identify and select studies for the assessment of Question 5 (rFVIIa), Question 7 (TXA), Question 8 (viscoelastic haemostatic assays) and Question 9 (cell salvage)



a. Search for SRs, RCTs, cohort studies conducted via Ovid (Embase, MEDLINE, EBM Reviews) and PubMed (in-process and citations not indexed in MEDLINE).

b. Protocol date limits as follows: Q5 – studies published prior to 2009; Q7 – studies published prior to 2000; Q8 – studies published prior to 2000; Q9 – studies published prior to 1990. In PubMed, studies published prior to 2015 were not screened (initial search); Updated search, studies published prior to 2019.

4.2 Prognostic factors (Question 1)

Question 1 - (Prognostic)

In patients with critical bleeding, which physiologic, biochemical and metabolic (including temperature) parameters should be measured early and frequently, and what values of these parameters are indicative of critical physiologic derangement?

4.2.1 Methods

This review sought to identify 8 potential prognostic factors associated with increased mortality and transfusion volume requirements in patients who are critically bleeding (i.e. major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion), regardless of age or clinical setting, as outlined in Figure 4.4 below.

Figure 4.4 PPO criteria: Question 1 – physiologic, biochemical and metabolic parameters



APTT, activated partial thromboplastin time; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; INR, international normalised ratio; PLT, platelets; PT, prothrombin time; RBC, red blood cells

Notes:

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. c. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2009, noting primary studies published prior to 2009 that had been included in a systematic review were also eligible for inclusion. There were no restrictions applied to study sample size. Assuming all relevant primary studies had been identified in the included systematic review; screening for lower-level evidence was not conducted.

The literature search was updated in August 2019²⁸ and again in September 2021 to identify any new systematic review studies meeting the eligibility criteria. In 2021, a systematic screen for prospective cohort studies was carried out and limited to studies published from 2019. This is because the most recent studies identified in the updated review did not cover all prognostic factors. No further limits were applied.

4.2.2 Summary of evidence

4.2.2.1 Systematic reviews

Twelve systematic reviews (36-47) were found that searched for, and assessed, at least one of the 8 prognostic factors of interest in this review. The characteristics of each review and the relevant prognostic factor/s and outcomes assessed are shown in Table 4.2.

One systematic review (Kamyszek 2019) searched for evidence relating to transfusion triggers in paediatric patients but found no data about the prognostic factors and their association with the outcomes of interest for this review. One systematic review (Razzaghi 2012) also did not identify any data in patients with thrombocytopenia in the setting of nonvariceal upper gastrointestinal bleeding. In the absence of data, these reviews are not further discussed in this report.

Review ID	Study design	Population	Prognostic factor/s	Outcomes
Review quality	(No. of studies)			
Vasudeva 2021 (36) <i>Critically low</i>	SR of RCTs and observational studies (3 studies)	Adult trauma patients with an admission ionised calcium measurement	lonised calcium	Mortality Transfusion volume
Kamyszek 2019 (37) <i>Critically low</i>	SR of observational studies (29 studies)	Paediatric population requiring massive blood transfusion	PT/INR, temperature	No relevant outcomes identified
Shih 2019 (38) <i>Critically low</i>	SR of observational studies (45 studies)	Trauma patients	Any scores or predictors of massive transfusion	Transfusion volume
Lilitsis 2018 (39) <i>Critically low</i>	Narrative SR (NR studies)	Severely injured patients	Any predictive factors that describe patient's status including coagulation	Mortality
Tran 2018 (40) Critically low	SR / MA (84 studies)	Adult patients with traumatic torso injuries	Any clinical, laboratory or imaging predictors available during the initial hour of resuscitation	Transfusion volumes
Levy 2017 (41) <i>Critically low</i>	Narrative review (8 studies)	Bleeding patients in the perioperative setting	Platelet transfusion in relation to triggers, dose and assessment of haemostatic efficacy	Transfusion volumes

Table 4.2	Characteristics and quality of included systematic reviews
Table 4.2	Characteristics and quality of included systematic review

²⁸ One additional systematic review was found (Kamyszek 2019).

Review ID <i>Review quality</i>	Study design (No. of studies)	Population	Prognostic factor/s	Outcomes
Baxter 2016 (43) Critically low	SR / MA of RCTs and observational studies (28 studies)	Adult trauma patients presenting to the ED	Blood lactate	Mortality Transfusion volumes
Poole 2016 (42) Critically low	SR / MA of RCTs and observational studies (5 studies)	Adult trauma patients in non-military setting (excluding TBI)	Measures of coagulopathy (fibrinogen, APTT, platelet count, INR)	Mortality
Haas 2015 (44) Critically low	SR of RCTs and observational studies (64 studies)	Patients in the perioperative setting or with massive bleeding	PT/INR or APTT	Mortality Transfusion volumes
Abdul-Kadir 2014 (45) <i>Critically low</i>	Narrative SR (NR studies)	Women with PPH	Fibrinogen levels	Transfusion volume
Pacagnella 2013 (46) <i>Critically low</i>	SR of observational studies (11 studies)	Obstetrics patients with haemorrhage	Shock index, heart rate, systolic blood pressure	Mortality Blood loss
Razzaghi 2012 (47) <i>Critically low</i>	SR of RCTs and cohort studies (18 studies)	Patients with thrombocytopenia in the setting of nonvariceal upper Gl bleeding.	Platelets	No studies identified specifically assessing patients with active GI haemorrhage

APTT, activated partial thromboplastin time; ED, emergency department; GI, gastrointestinal; INR, international normalised ratio; MA, meta-analysis; NR, not reported; PPH, postpartum haemorrhage; PT, prothrombin time; RCT, randomised controlled trial; SR, systematic review; TBI, traumatic brain injury

Overall, there were 50 primary studies reported in the included systematic reviews, which were often judged by the review authors to have high or moderate concerns of bias related to study design features and likely reporting bias.

(Martin 2005, Balvers 2016, Aslar 2004, Callaway 2009, Duane 2008, Lavery 2000, Mizushima 2011, Neville 2011, Odom 2012, Regnier 2012, Vandromme 2010, Gale 2016, Odom 2013, Heinonen 2014, Cherry 2006, Vasudeva 2020, Hagemo 2014, Mitra 2010, MacLeod 2003, Hess 2009, Mitra 2007, Rourke 2012, Sambasivan 2011, Ciavarella 1987, Callcut 2011, Vandromme 2011, Baron 2004, Ipekci 2013, Paulus 2014, Callcut 2013, Leemann 2010, Schöchl 2011, Schreiber 2007, Arnold 2006, Fayed 2013, McGrath 2008, Premaratne 2001, Tanaka 2014, Wu 2014, van Hout 2017, Mannucci 1982, Murray 1998, Charbit 2007, Cortet 2012, Peyvandi 2012, Rouse 2006, Nakamura 2017, Magnotti 2011, Sperry 2018 [PAMPer], Moore 2018 [COMBAT]).

Temperature

Two reviews (Lilitsis 2018, Shih 2019) reported evidence from 3 observational studies relating to temperature in critically bleeding patients. Two studies (Balvers 2016, Martin 2005) examined the association between temperature with mortality and one study (Callcut 2011) assessed the association between temperature and transfusion requirements. The studies were carried out in trauma centres in the US (2 studies) or The Netherlands (one study).

Two studies (Balvers 2016, Martin 2005,) were judged by Lilitsis 2018 to have concerns of bias related to related to study design, and one study (Callcut 2011) was judged by Shih 2019 to be of good methodological quality with no serious concerns of bias.

Acid-base status

Three reviews (Lilitsis 2018, Tran 2018, Baxter 2016) reported evidence from 15 observational studies relating to acid-base status in critically bleeding patients. Twelve studies (Gale 2016, Heinonen 2014, Odom 2013, Odom 2012, Regnier 2012, Mizushima 2011, Neville 2011, Vandromme 2010, Callaway 2009, Duane 2008, Aslar 2004, Lavery 2000) assessed the association between lactate levels and mortality and 5 studies (Ipekci 2013, Regnier 2012, Vandromme 2011, Vandromme 2010, Baron 2004) assessed the association between lactate levels and transfusion volume.

The studies were carried out in various trauma centres in the US, France, Switzerland and South Africa. The overall risk of bias was judged to be moderate or high due to attrition, confounding and reporting biases.

lonised calcium

Two reviews (Vasudeva 2021, Shih 2019) reported evidence from 3 studies relating to ionised calcium in critically bleeding patients. Three observational studies (Vasudeva 2020, Magnotti 2011, Cherry 2006) assessed the association between ionised calcium and mortality and 2 studies (Magnotti 2011, Vasudeva 2020) reported on transfusion volume. The studies were carried out in trauma centres in the US and Australia.

Vasudeva 2021 assessed the quality of included studies to be moderate, noting that none of the included studies were blinded nor explicitly stated the utilisation of different reviewers for data collection and cross checking. Shih 2019 did not assess risk of bias of included studies. Overall, risk of bias for included observational studies was judged to be moderate due to limited by sample size and confounding.

Haemoglobin

One review (Shih 2019) identified 5 observational studies (Callcut 2013, Paulus 2014, Vandromme 2011, Callcut 2011, Leemann 2010, Schöchl 2011, Schreiber 2007) that assessed the association between haemoglobin and transfusion volume or transfusion requirements in trauma patients with critical bleeding. The studies were carried out in trauma centres in the US, Switzerland, Austria and Iraq. No studies were found that assessed the association between haemoglobin and mortality.

Tran 2018 found the quality of included studies was poor noting the frequent lack of justification, inadequate reporting and suboptimal handling of missing data. Overall, risk of bias for the included observational studies was judged to be moderate to high due to study design and confounding.

Platelet count

Two reviews (Poole 2016, Levy 2017) included data from 9 observational studies in trauma or perioperative surgical patients with critical bleeding that examined the association between platelet count and mortality (2 studies) (Hagemo 2014, Mitra 2010) or transfusion volume (7 studies) (Arnold 2006, Fayed 2013, McGrath 2008, Premaratne 2001, Tanaka 2014, Wu 2014, van Hout 2017). Three studies were carried out in trauma or emergency centres in the US, UK, Norway and Australia. Eight studies were carried out in surgical settings in the US, Canada, Netherlands and Egypt.

Poole 2016 noted the included studies provided very low certainty of evidence, with issues arising due to variables utilised in prediction models and generalisability or results. Overall, the included observational studies were judged to have high risk of bias related to patient selection and confounding.

PT/INR

Five reviews (Lilitsis 2018, Poole 2016, Haas 2015, Tran 2018, Shih 2019) included data from 8 observational studies that assessed the association between PT/INR levels with mortality (5 studies; Macleod 2003, Hess 2009, Mitra 2007, Hagemo 2014, Mitra 2010) or transfusion volume (3 studies; Callcut 2013, Vandromme 2011, Schreiber 2007) in trauma patients with critical bleeding.

All studies were conducted in trauma centres in the US, UK, Norway, Australia and Iraq and typically used an INR value 1.5 times the upper limit of normal as reference. Overall, risk of bias for included observational studies was judged to be high for inadequate control for confounding, study design and reporting.

APTT

Three reviews (Poole 2016, Lilitsis 2018, Haas 2015) identified 7 observational studies that assessed the association between APTT levels with mortality (5 studies; Rourke 2012, Macleod 2003, Sambasivan 2011, Ciavarella 2987, Mitra 2007) or transfusion volume (2 studies; Mannucci 1982, Murray 1998) in trauma patients with critical bleeding.

All studies were conducted in trauma centres in the US, UK, Norway, Italy and Australia. Overall, risk of bias for included observational studies was judged to be unclear to high due to study design, reporting and control for confounding.

Fibrinogen levels

Three reviews (Poole 2016, Abdul-Kadir 2014, Shih 2019) identified 2 observational studies that assessed the association between fibrinogen levels and mortality (Hagemo 2014, Rourke 2012) in critically bleeding trauma patients. The review also included 5 studies that assessed the association between fibrinogen levels and transfusion volumes in trauma and obstetric patients with critical bleeding (Charbit 2007, Cortet 2012, Peyvandi 2012, Rouse 2006, Nakamura 2017).

Four studies were conducted in obstetric settings in the US, France and Italy and 3 studies were carried out in trauma centres in the US, UK, Norway and Japan. Overall, included studies was judged to be high risk of bias due to study design, confounding and reporting biases.

4.2.2.2 Primary studies

There were 3 additional prospective cohort studies (Gaessler 2021, Javali 2017, McQuilten 2017a), 4 additional retrospective cohort studies (McQuilten 2017b, Kawatani 2016, Noorbhai 2016, Sawamura 2009) and 2 secondary analyses of RCTs (Moore 2020, Lester 2019) identified through the systematic review and handsearching process that evaluated one or more of the 8 prognostic factors of interest in this review.

A summary of the characteristics and risk of bias of the additional studies and relevant outcomes assessed are shown in Table 4.3.

Three other retrospective cohort studies (Figueiredo 2018, Verma 2017, Wang 2016) were identified in the literature search but were later excluded as they did not report any data on the prognostic factor or outcomes of interest and were not considered further. A full list of studies that potentially met the inclusion criteria but were not included in the evidence evaluation is provided in **Appendix C** (technical report, volume 2).

Prospective cohort studies

Gaessler 2021 was a prospective observational study conducted at a single centre in Germany that assessed the impact of coagulopathy in 148 injured patients who were medical treated by the Helicopter Emergency Medical Service and transported to Level 1 trauma centres. The study was found to be at moderate risk of bias due related to lack of blinding or outcome assessors.

Javali 2017 was a prospective observational study in 100 trauma patients (nonconsecutive) at risk of haemodynamic compromise in a tertiary care centre emergency department in India. This study was found to be at serious risk of bias due to inadequate control of confounding factors and measurement bias. The study included 92 patients in the analysis of base deficit and did not provide justification for patients lost to follow-up.

McQuilten 2017a was a prospective study that assessed the association of low fibrinogen levels with mortality in all adult trauma patients identified through a statewide trauma registry in Victoria (Australia). Data were available for 4772 patients who presented to the 2 major trauma hospitals between January 2008 and July 2011 and who had a fibrinogen level measured during initial resuscitation. The study had some concerns of bias relating to measurement of outcomes and missing data.

Retrospective cohort studies

McQuilten 2017b was a retrospective cohort study that examined the prognostic value of fibrinogen levels on mortality and transfusion volume in adult trauma patients who received massive transfusion in hospitals across Australia and New Zealand. A total of 2829 patients received massive transfusion between April 2011 and October 2015, which was defined as 5 or more units of RBC within any four-hour period during admission. This study had moderate concerns of bias relating to measurement of the outcome and missing data.

Kawatani 2016 was a retrospective study of the medical records of 25 patients who underwent endovascular aortic repair for ruptured abdominal aortic aneurysms (rAAA) at Chiba-Nishi General Hospital in Japan between October 2013 and December 2015. Major coagulopathy was defined using a PT/INR or APTT ratio greater than 1.5 times the upper limit of normal, or platelet count less than $50 \times 10^{\circ}$ /L. The study was judged to be at serious risk of bias due to patient selection bias and likely confounding.

Noorbhai 2016 was a retrospective cohort study that aimed to assess the correlation between coagulopathy (INR) and mortality in 1000 patients admitted to a level 1 trauma unit in South Africa. INRs were not recorded in 61 patients and were therefore excluded from the analysis to a total of 939 remaining patients. The INR was dichotomised into \leq 1.2 and >1.2, then correlated with ISS and in-hospital mortality. This study was found to have serious risk of bias relating to study design and lack of control for confounding factors.

Sawamura 2009 was a retrospective cohort study conducted in Japan that assessed the impact of disseminated intravascular coagulation on patient outcomes. Data obtained at

4 time points (within 24 hours of arrival to the emergency department) was collected from 314 consecutive severe trauma patients which was further subdivided into 259 survivors and 55 nonsurvivors. This study had serious concerns of bias relating to study design, confounding and inadequate reporting of data.

Secondary analysis of RCTs

Moore 2020 evaluated the association between prehospital plasma and hypocalcaemia with lower survival. To investigate, Moore 2020 used data collected from 2 RCTs, COMBAT (Moore 2018) which included injured adults ≥18 years with acute blood loss and PAMPer (Sperry 2018) which included injured adults at risk of haemorrhagic shock. The authors noted limitations of the studies for the purposes of the secondary analysis acknowledging that these biases can potentially limit the generalisability of the results. These include biases due to outcome data (lack of ionised calcium measurements for all enrolled patients), pre-existing disease severity and survivor bias.

Lester 2019 provided a secondary analysis of data from an RCT to evaluate the association between hypothermia and patient outcomes using the dataset collected during the PROPPR RCT (Holcomb 2015). Hypothermia was defined as a temperature less than 36°C and normothermia was considered to be between ≥36°C and 38.5°C. The study had several limitations relating to measurement of the outcome (no standardised method and variability in devices used), reporting of the outcome (pooling of data across 12 sites) and differences in protocols. Overall, Lester 2019 was judged to be at serious risk of bias due to study design, confounding and reporting.

Review ID Risk of bias	Study design	Population N	Prognostic factor	or Outcomes					
Prospective cohort studies									
Gaessler 2021 (48) <i>Moderate</i>	Prospective cohort, SC	Adult trauma patients medically treated by HEMS enroute to a Level 1 trauma centre N=148	Coagulopathy	Mortality Transfusion volume					
Javali 2017 (49) <i>Critical</i>	Prospective cohort, SC	Trauma patients at risk of hemodynamic compromise N=100	Acid-base status	Mortality Transfusion volumes					
McQuilten 2017a (50) <i>Moderate</i>	Prospective cohort between January 2008 and July 2011 and	Adult trauma patients who had a fibrinogen level measured during initial resuscitation N=4772	Fibrinogen levels	Mortality					
Retrospective c	ohort studies								
Kawatani 2016 (51) <i>Critical</i>	Retrospective cohort	Patients who undergo endovascular aortic repair for rAAA N=25	Measures of coagulopathy	Mortality					
Noorbhai 2016 (52) <i>Critical</i>	Retrospective cohort	Patients admitted to the level 1 trauma unit N=1000	INR	Mortality					

Table 4.3 Characteristics and quality of additional primary studies included in the review

Review ID Risk of bias	Study design	Population N	Prognostic factor	Outcomes
Sawamura 2009 (53) <i>Critical</i>	Retrospective cohort	Severe trauma patients with ISS ≥ 9 N=314	Coagulation and fibrinolytic markers (fibrinogen, PT, lactate, platelet count)	Mortality Transfusion volumes
McQuilten 2017b (54) <i>Moderate</i>	Retrospective cohort	Adult patients who received massive transfusion (≥ 5 units of RBC within any 4 hour period during admission) N=2829	Fibrinogen levels	Mortality Transfusion volumes
Single arm ana	lysis of RCT			
Moore 2020 (55) <i>Moderate</i>	Secondary analysis of RCT data	Adults with traumatic haemorrhagic shock	lonised calcium	Mortality
Lester 2019 (56) PROPPR trial <i>Serious</i>	Secondary analysis of RCT	Severely injured patients ≥ 15 years N=586	Hypothermia ^a	Mortality

HEMS, Helicopter Emergency Medical Service; INR, international normalised ratios; INR, international normalised ratio; ISS, injury severity score; MC, multicentre; PT, prothrombin time; rAAA, ruptured abdominal aortic aneurysm; SC, single centre a: Data sets from RCTs were used to evaluate prognostic markers

4.2.3 Results

4.2.3.1 Mortality

A summary of the evidence relating to mortality in patients with critical bleeding is presented in Table 4.4. Due to the limited evidence and significant heterogeneity among included studies, no meta-analysis was performed.

Temperature

Identified literature suggests hypothermia is independently associated with an increased risk of mortality among critically bleeding patients (*GRADE: Very low*). Four studies in the trauma setting contributed data, with adjusted odds ratios (OR) of around 2.7 observed at 24-hours and adjusted OR ranging from 1.8 to 2.8 observed at 30 days. Hypothermia was generally reported by study authors to be below 35°C.

Acid-base status

Identified literature suggests risk of mortality is significantly increased with increasing lactate levels among patients with critical bleeding (*GRADE: Very low*). Fourteen observational studies in trauma settings contributed mortality data. At high lactate levels (> 4 mmol/L), authors reported OR ranging between 3.8 and 10.58.

lonised calcium

Multiple observational studies have found that hypocalcaemia is common in the context of major bleeding and appears to be associated with mortality, however this may be confounded by increased blood transfusions and injury severity (*GRADE: Very low*).

Identified literature suggests an increased risk of mortality associated with hypocalcaemia. Four studies conducted in trauma patients contributed data, where hypocalcaemia was defined as either < 1.11 or <1.0 mmol/L ionised calcium. Pooled (unadjusted) data suggested the mortality rate to be 24% among those with hypocalcaemia, compared with 15% among those with normocalcaemia (OR 1.87; 95% CI 1.27, 2.75; p = 0.001; random effects, $l^2 = 0\%$) (*GRADE: very low*). After adjustment for confounders (age, ISS, Shock index) in a Cox Proportional Hazards Model, one study (Moore 2020) suggested hypocalcaemia to be independently associated with survival (HR 1.07; 95% CI 1.02, 1.13; p = 0.01).

Haemoglobin

No identified literature reported on the effect of haemoglobin levels and mortality.

Platelet count

The association between platelet count and mortality is unclear (*GRADE: Very low*). Three studies suggested lower platelet counts are not associated with an increased risk of mortality in critically bleeding trauma or surgical patients (adjusted OR ranged between 0.99 and 1.0; p > 0.5). One study (McQuilten 2017b) suggested platelet counts below 100 x 10°/L to be independently associated with survival (adjusted OR 0.50; 95% Cl 0.30, 0.84; p = 0.009) [after adjustment for age, ISS, Shock index). One study (Sawamura 2009) suggested lower platelet counts were associated with increased prediction of death (stepwise logistic regression, OR 1.097; 95% Cl 1.003, 1.116; p = 0.003) [including DIC scores, lactate coagulation and fibrinolysis variables].

PT/INR

Identified literature suggests abnormal PT/INR levels among patients with critical bleeding are associated with an increased risk of mortality (*GRADE: Very low*). Adjusted OR ranged from 1.35 to 3.23 and an adjusted risk ratio (aRR) of 3.68 observed for elevated PT/INR levels compared to normal levels. One study in patients undergoing endovascular aortic repair (rAAA) reported no significant association (p > 0.05) but there were too few patients for any meaningful analysis.

APTT

Identified literature suggests an increased risk of mortality associated with abnormal APTT levels among patients with critical bleeding (*GRADE: Very low*). Six studies in trauma and surgical settings contributed data reporting OR ranges between 1.01 and 4.26 for elevated APTT levels compared to normal APTT levels.

Fibrinogen

Identified literature suggests a significant associated between the risk of mortality and low fibrinogen levels among patients with critical bleeding (*GRADE: Very low*). Definitions of low fibrinogen levels varied across the studies but were generally considered to be levels < 1.5 g/L. Two studies reported an adjusted odds ratio (OR) that ranged between 1.29 and 3.28 for fibrinogen levels lower than 2.0 g/L and 3 studies reported an association with survival (OR ranged between 0.08 to 0.99). One study did not provide usable data.

One study also reported fibrinogen levels above 4 g/L to be associated with an increased risk of mortality (OR 2.03; 95% Cl 1.35, 3.40; p = 0.001) in patients who had received a massive transfusion (compared against fibrinogen levels between 2 to 4 g/L).

Table 4.4Results for physiologic, biochemical and metabolic (including temperature) parameters indicative of critical physiologic derangement:
Patients with critical bleeding – Mortality

Study ID	Sample size	Patient population	Setting	Predictor	Results			
Study design ^a	(no. Of trials) included in analysis		(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Temperature								
Lilitis 2018 SR	N = 702 444 (2 Coh)	Trauma patients	Trauma (US, Netherlands)	Hypothermia vs normothermia	Authors suggest hyp increased mortality r	othermia is associated isk.	l with significant	Significant association Favours normothermia
Critically low quality	N= 701 491 Martin 2005			< 35 °C	Mortality 25.5%	Mortality 3.0%	NR	<i>p</i> = NR
	N = NR Balvers 2016			24-hr mortality 30-day mortality	NR NR	NR NR	OR 2.72 OR 2.82	<i>p</i> = NR <i>p</i> = NR
McQuilten 2017a Prospective Coh Moderate risk of bias	N = 4773	Patient with major trauma	Trauma registry (Australia)	Temperature <35 ℃ 35 to 36.5 ℃ >37.5 ℃ missing	In-hospital mortality n=428 n=1732 n=295 n=536	36.6 to 37.5°C (reference)	Adjusted ^d OR 1.91 (1.28, 2.85) OR 1.11 (0.80, 1.56) OR 0.597 0.72 (0.35, 1.50)	Significant association p = 0.002 p = 0.53 p = 0.38
Lester 2019 Secondary analysis of RCT Serious risk of bias	N = 586	Trauma patients	Level 1 trauma centres (US)	Hypothermia 24-hr mortality 30-day mortality	NR/399	NR/187	Adjusted ^e OR 2.7 (1.7, 4.5) OR 1.8 (1.3, 2.4)	Significant association p < 0.00 p < 0.00
Acid-base status								
Baxter 2016	N = 34 120	Trauma patients	Trauma/		Authors report highe	er odds of mortality as l	actate levels increase.	Favours lactate < 2 mmol/L
SR Critically low quality	(9 Coh) Aslar 2004 Callaway 2009 Duane 2008 Lavery 2000 Mizushima 2011 Neville 2011 Odom 2012 Regnier 2012 Vandromme 2010		Emergency (US, France, Switzerland)	Lactate levels ≥ 2 mmol/L ≥ 4 mmol/L	NR	NR	OR range 1.067 to 1.79 OR range 4.2 to 10.58	Significant association p = NR p < 0.001
Lilitis 2018	N = 6794 (3 Coh)	Trauma patients	Trauma (US,	Acid-base status				Favours lower lactate
SR G Critically low quality	Gale 2016	Gale 2016	South Africa)	Lactate levels Base deficit	A one mmol/L increase in lactate levels is associated with a 17% increased mortality risk. A one mq/L increase in base deficit is associated with approximately 4% increased mortality risk		Significant association p = NR	

Study ID	Sample size	Patient population	Setting	Predictor	Results			
Study design ^a	(no. Of trials) included in analysis		(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
	Heinonen 2014			Lactate levels	High lactate* associated with 54% mortality rate *not normalised within 24 hrs	< 2.5 mmol/L associated with 22% mortality rate	NR	Favours lower lactate Significant association p = NR
	Odom 2013			Lactate levels < 2.5 mmol/L 2.5–3.9 mmol/L > 4 mmol/L	NR	NR	OR 1 (NR) OR 1.5 (NR) OR 3.8 (NR)	Favours lactate < 2.5 mmol/L Significant association p = NR
Javali 2017 Prospective Coh <i>Critical risk of bias</i>	N = 100	Trauma patients at risk of haemodynamic compromise	Tertiary care centre ED (India)	Arterial lactate 24-hour mortality	≥ 4 mmol/L NR (38.1%)	<4 mmol/L NR (0%)	NR	Favours lactate < 2 mmol/L Significant association p < 0.001
				Base deficit 24-hour mortality	≥ 12 mEq/L 30.4% increased risk	< 12 mEq/L NR (1.3%)	NR	Favours base deficit < 12 mEq/L Significant association p = NR
McQuilten 2017b Retrospective Coh Moderate risk of bias	N = 2829	Trauma patients who received massive transfusion	Trauma registry (Australia, New Zealand)	Base deficit -29 to -8.7 -8.6 to -5 -4.9 to -1.5	In hospital mortality	≥ -1.4 (reference)	Adjusted OR ^r OR 3.68 (2.70, 5.03) OR 1.33 (0.95, 1.86) OR 0.94 (0.66, 1.33)	Adjusted <i>p</i> < 0.001 <i>p</i> = 0.10 <i>p</i> = 0.72
Sawamura 2009 Retrospective Coh <i>Critical risk of bias</i>	N = 314	Severe trauma patients with an ISS of ≥ 9	ED (Japan)	Lactate (mmol/L)	NR	NR	OR 1.236 (1.016, 1.502)	Favours lower lactate Significant association p = 0.034
Ionised calcium								
Vasudeva 2021 SR Critically low quality	N = 1213 (3 Coh) Cherry 2006 Magnotti 2011 Vasudeva 2020	Trauma patients not receiving blood transfusion	Trauma centres (US, Australia)	Hypocalcaemia (< 1.11 mmol/L) prior to transfusion	NR 24/91 (26.4) NR/332 (15.5) 29/113 (25.6)	NR 48/305 (15.7) NR/259 (8.7) 17/113 (15.0)	NR ^c OR 1.92 (NR) NR OR 1.95 (1.00, 3.80)	NR p < 0.05 p = 0.036 p = 0.047
Moore 2020	N = 160	Patients with	Trauma centres	Hypocalcaemia	13/70 (18.6)	11/90 (12.2)	OR 1.64 (0.68, 3.92) ^c	<i>p</i> = 0.26
Secondary analysis of RCTs <i>Moderate risk of</i> <i>bias</i>		traumatic haemorrhagic shock from blunt or penetrating injuries	(US)	(i-Ca < 1.0 mmol/L) vs normocalcaemia (i-Ca > 1.0 mmol/L)	Hypocalcaemia independently associated with survival after adjustment for confounders (age, ISS, Shock index) in a Cox Proportional Hazardds Model.		HR 1.07 (1.02, 1.13)	<i>p</i> = 0.01

Study ID	Sample size	Patient population	Setting	Predictor	Results			
Study design ^a	(no. Of trials) included in analysis		(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Haemoglobin – no	studies identified					·		
Platelet count								
Poole 2016	N = 1650	Patients with non-TBI	Trauma (MC [US,	Platelet count			Adjusted	No significant association
SR	(2 Coh)	trauma	UK, Norway] and					p = NR
Critically low quality	Hagemo 2014 Mitra 2010		Australia)		99/1133 (8.7) 99/331 (29.9)		OR 1 (1.0, 1.0) OR 0.99 (0.99, 0.99)	
McQuilten 2017a	N = 4773	Patient with major	Trauma registry	Platelet count	In-hospital mortality	> 150 x10 ⁹ /L	Adjusted OR ^d	Adjusted:
Prospective Coh		trauma	(Australia)	< 100 x 10º / L 100 to 150 x 10º / L		(reference)	OR 0.50 (0.30, 0.84)	p = 0.009
Moderate risk of bias				100 10 130 x 10 7 E			OK 0.98 (0.09, 1.40)	μ - 0.91
Sawamura 2009	N = 314	Severe trauma	ED (Japan)	Platelet count	Survivors (n=259)	Non-survivors (n=55)	NR	No significant association
Retrospective coh		patients with an ISS of			159 ±79	147 ±82		<i>p</i> = 0.182
Critical risk of bias		29			Stepwise logistic regr of death.	ression for prediction	OR 1.097 (1.003, 1.116)	<i>p</i> = 0.003
Kawatani 2016	N= 25	Patients undergoing	Surgical (Japan)	Preoperative platelet	Survivors	Non-survivors	NR	NR
Retrospective coh		endovascular aortic		count (10⁴/uL)				
Critical risk of bias		repair (rAAA)		24-hour 30-day	22/25 (88) 16.1 +/- 5.4 20/25 (80) 16.2 +/- 5.54	3/25 (12) 17.3 +/- 3.0 5/25 (20) 16.8 +/- 2.7		p = 0.616 p = 0.767
				Postoperative platelet count (10 ⁴ /uL)	Survivors	Non-survivors	NR	NR
				24-hour	22/25 (88) 10.2 +/- 5.0	3/25 (12) 7.7 +/- 1.9		<i>p</i> = 0.558
				30-day	20/25 (80) 10.4 +/- 5.0	5/25 (20) 7.2 +/- 1.9		<i>p</i> = 0.299
Lilitis 2018	N = 7638	Trauma patients	Trauma (US)	PT	NR	NR	Abnormal PT	NR
SR	(1 Coh)		indumid (00)				associated with 35%	
Critically low	MacLeod 2003						increased risk of	
quality							mortality	
Poole 2016	N = 7638	Patients with non-TBI	Trauma (MC [UK,	PT	NR	NR	OR 1.35 (1.11–1.68)	NR
SR	(1 Coh)	trauma	US, Norway],					
Critically low	MacLeod 2003		Australia)					
quality	N = 1650 (2 Coh)			INR	00/1122 (0.7)			NR
	Hagerno 2014 Mitra 2010	-			99/331 (29.9)		OR 1.65 (0.65-4.18) OR 1.43 (1.02-2.01)	
Haas 2015	N= 35 441	Patients with critical	Trauma (US,	INR	Hess 2009 reported a	in increase in INR	NR	NR
эк	(2 CON)	Dieeding	Australia)		in-hospital mortality	ise relationship with		
	1633 2009							

Study ID	Sample size	Patient population	Setting	Predictor	Results			
Study design ^a	(no. Of trials) included in analysis		(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Critically low quality	Mitra 2007				Mitra 2007 reported a between high INR an	an association Id mortality	OR 1.62 (1.18, 2.24)	<i>p</i> < 0.01
McQuilten 2017a Prospective Coh Moderate risk of bias	N = 4773	Patient with major trauma	Trauma registry (Australia)	INR 1.5 to 1.9 >2.0	In-hospital mortality	< 1.5 (reference)	Adjusted OR OR 3.23 (2.12, 4.92) OR 3.02 (1.82, 5.03)	Adjusted: <i>p</i> = 0.009 <i>p</i> = 0.91
Noorbhai 2016 Retrospective Coh <i>Critical risk of bias</i>	N = 939	Trauma patients	Level I trauma centre (South Africa)	INR All patients External admissions Interhospital transfers	High (>1.20) 74/482 (15.4) 15/121 (12.4) 59/361 (16.3)	Low (<1.20) 132/457 (28.9) 44/107 (41.1) 88/350 (28.9)	Adjusted RR aRR 1.92 (1.49, 2.48) aRR 3.68 (2.11, 6.44) aRR 1.54 (1.15, 2.05)	<i>p</i> < 0.001 p < 0.001 p = 0.004
Kawatani 2016 Retrospective Coh <i>Critical risk of bias</i>	N = 25	Patients undergoing endovascular aortic repair (rAAA)	Surgical (Japan)	PT-INR, pre-operative 24-hour 30-days	Survivors N = 22/25 1.2 +/- 0.16 N = 20/25 1.2 +/- 0.16	Non-survivors N = 3/25 1.2 +/- 0.2 N = 5/25 1.23 +/- 0.19	NR	NR p = 0.802 p = 0.0767
				PT-INR, post-operative 24-hour 30-day	Survivors N = 22/25 1.3 +/- 0.2 N = 20/25 1.4 +/- 0.2	Non-survivors N = 3/25 1.5 +/- 0.28 N = 5/25 1.5 +/- 0.2	NR	NR p = 0.295 p = 0.148
APTT								
Lilitis 2018 SR <i>Critically low</i> <i>quality</i>	N = 7638 (1 Coh) MacLeod 2003	Trauma patients	Trauma (US)	APTT	Elevated APTT was associated with 326% increased risk of mortality			
Poole 2016 SR Critically low quality	N = 9336 (3 Coh) Rourke 2012 MacLeod 2003 Sambasivan 2011	Patients with non-TBI trauma patients	Trauma (UK, US, Norway)	APTT	62/517 (12.0) NR 173/1181 (14.6)	NR	OR 1.05 (1.01–1.09) OR 4.26 (3.23–5.62) OR 1.015 (1.01–1.02)	NR
Haas 2015N = 155 (2 Coh)SRCiavarella 1987Critically lowHess 2000Mitra 2001		Patients with critical bleeding	Trauma (US, Australia)	APTT Ciavarella 1987	Microvascular bleedir coagulation factor lev control).	NR		
quality	With a 2007			Hess 2009	An increase in aPTT d hospital mortality			
				Mitra 2007	Higher APTT is a pred	lictor of mortality	OR 1.01 (1.01, 1.02).	<i>p</i> < 0.01
Kawatani 2016 Retrospective Coh <i>Critical risk of bias</i>	N = 25	Patients undergoing endovascular aortic repair (rAAA)	Surgical (Japan)	APTT, seconds (pre- opoerative) 24-hour	Survivors N = 22/25 27.0 +/- 4.3	Non-survivors N = 3/25 33.6 +/- 8.4	NR	NR p = 0.21
				30-days	N = 20/25 26.8 +/- 4.3	N = 5/25 32 +/- 7.0		<i>p</i> = 0.119

Study ID Study design ^a	Sample size (no. Of trials) included in analysis	Patient population	Setting (Location)	Predictor	Results				
					Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance p</i> -value Heterogeneity ^b	
				APTT, seconds (post- operative) 24-hour 30-day	Survivors N = 22/25 38.9 +/-8.7 N = 20/25 38.1 +/- 7.9	Non-survivors N = 3/25 108.7 +/- 63.4 N = 5/25 95.7 +/- 57.9	NR	NR <i>p</i> = 0.006 <i>p</i> = 0.002	
Fibrinogen level									
Poole 2016 SR Critically low quality	N = 1650 (2 Coh)	Patients with non-TBI trauma	Trauma (UK, US, Norway)	Fibrinogen Hagemo 2014 Low level <2.29 High level >2.29 Rourke 2012	28-day survival 99/1133 (8.7) 62/517 (12.0)		Adjusted OR 0.08 (0.03, 0.20) OR 1.77 (0.94, 3.32) OR 0.22 (0.10, 0.47)	<i>p</i> < 0.001 <i>p</i> = 0.076 <i>p</i> < 0.001	
Gaessler 2021 Prospective Coh Moderate risk of bias	N = 148	Trauma patients	Level 1 trauma centres (Germany)	Fibrinogen	A correlation between prognostic indicators and mortality could not be determined. No data reported on prognostic factors and their association with outcomes of mortality or transfusion requirements. However, TIC and TIC with hyperfibrinolysis resulted in worse prognosis for mortality compared to those without coagulopathy.			t be determined. utcomes of mortality or sis resulted in worse	
McQuilten 2017a Prospective Coh <i>Moderate risk of</i> <i>bias</i>	N = 4773	Patient with major trauma	Trauma registry (Australia)	Fibrinogen < 1.0 g/L 1.0-1.5 g/L 1.6-1.9 g/L > 4.0 g/L	In-hospital mortality 54/114 (47.4) 71/283 (25.1) 77/617 (12.5) 53/735 (7.2)	2.0-4.0 g/L (reference) 186/3024 (6.2)	Adjusted OR ⁴ OR 3.28 (1.71, 6.28) OR 2.08 (1.36, 3.16) OR 1.39 (0.97, 2.00) OR 1.04 (0.70, 1.52)	Adjusted <i>p</i> < 0.001 <i>p</i> = 0.001 <i>p</i> = 0.08 <i>p</i> = 0.86	
McQuilten 2017b Retrospective Coh Moderate risk of bias	N = 2829	Trauma patients who received massive transfusion	Trauma registry (Australia, New Zealand)	Fibrinogen < 1.0 g/L 1.0-1.5 g/L 1.6-1.9 g/L > 4.0 g/L	In-hospital mortality 91/198 (46) 163/622 (26) 103/532 (19) 56/244 (23)	2.0-4.0 g/L (reference) 200/1233 (16)	Adjusted OR [†] OR 2.31 (1.48, 3.60) 1.0-1.9: OR 1.29 (0.99, 1.67) OR 2.03 (1.35, 3.04)	<i>p</i> <0.001 <i>p</i> = 0.056 <i>p</i> = 0.001	
Sawamura 2009 Retrospective Coh <i>Critical risk of bias</i>	N = 314	Severe trauma patients with an ISS of ≥ 9	ED (Japan)	Fibrinogen (g/L), mean (SD)	Survivors (n = 259) 2.53 (0.9)	Non-survivors (n=55) 1.44 (0.8)	Stepwise logistic regression OR 0.989 (0.979, 0.998)	<i>p</i> = 0.015	

APTT, activated partial thromboplastin time; aRR, adjusted relative risk; CI, confidence interval; Coh, cohort study; ED, emergency department; i-Ca; ionised calcium; INR, international normalised ratio; ISS, injury severity score; MC, multicentre; NR, not reported; OR, odds ratio; PT; prothrombin time; RCT, randomised controlled trial; SR, systematic review; TBI, traumatic brain injury; TIC, trauma-induced coagulopathy; UK, United Kingdom; US, United States

a. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if Phet >0.1 and I2 <25%; (ii) mild heterogeneity if I2 <25%; moderate heterogeneity if I2 between 25–50%; substantial heterogeneity I2 >50%.

c. Calculated post-hoc using RevMan 5.4.

d. Adjusted for age, gender, ISS, injury type, pH, temperature, Glasgow Coma Score (GCS), initial international normalised ratio and platelet count.

e. Adjusted for number of RBC units used in 24 hours, need for emergent OR (within 90 minutes of arrival), ISS, mechanism of injury (blunt versus penetrating), weight, age, sex, and initial pulse and systolic blood pressure on arrival

f. Adjusted for hospital, age, gender, clinical context, CCI, Hb, platelet count, APTT, INR and base excess at massive transfusion commencement.

4.2.3.2 Transfusion volume

A summary of the evidence reported in the identified literature relating to mortality in patients with critical bleeding is presented in Table 4.5. Due to the limited evidence and significant heterogeneity, no meta-analysis was performed.

Temperature

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Among trauma patients, one study reported an increased risk of massive transfusion (\geq 10 units in 6 hours) (OR 4.0; 95% Cl1.6, 10.1) to be associated with hypothermia in patients with critical bleeding and one study reported no important association between hypothermia and the volume of RBC transfused (RR 0.90; 95% Cl 0.89, 0.92).

Acid-base status

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Included studies were in trauma settings and reported an increased risk of higher RBC transfusion requirements associated with increased lactate levels in patients with critical bleeding. Two studies reported OR of 3.13 and 5.20 (not reported for other studies). High lactate levels were reported above 2.9 mmol/L.

lonised calcium

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Observational studies in trauma settings reporting a significant association between hypocalcaemia and increased transfusion requirements in critically bleeding patients.

Haemoglobin

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Seven observational studies in trauma settings contributed data, reporting an association between low haemoglobin levels and increased risk of transfusion requirements.

Platelet count

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Included studies were in surgical settings and reported an association between low platelet count and increased transfusion requirements (*GRADE: Very low*). Studies included varying measurements of platelet count to trigger transfusion requirements, making it difficult to draw conclusions.

PT/INR

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Included studies were in the trauma setting, reporting increased PT/INR levels were associated with an increased an increased risk of massive transfusion (10 or more units of RBC) (OR ranges between 2.1 and 5.9).

APTT

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*), with studies in trauma and surgical settings reporting an association between increased APTT levels and increased risk of massive transfusion in patients with critical bleeding.

Fibrinogen

Only limited conclusions can be drawn from the available evidence (*GRADE: Very low*). Six observational studies in the trauma and obstetrics settings contributed data, with 5 studies reporting a significant association between low fibrinogen levels and increased RBC transfusion volumes in patients with critical bleeding. Definitions of low fibrinogen levels were commonly considered less than 2 g/L.

Table 4.5Results for physiologic, biochemical and metabolic (including temperature) parameters indicative of critical physiologic derangement:Patients with critical bleeding – Transfusion volume

Study ID Study designª	Sample size (no. of trials) included in analysis	Patient population	Setting (Location)	Predictor	Results					
					Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b		
Temperature										
Shih 2019 SR <i>Critically low quality</i>	N = 170 (1 Coh) Callcut 2011	Trauma patients	Trauma (US)	Hypothermia (< 35.5°C) ≥10 units RBC / 6hrs	NR	NR	OR 4.0 (1.6, 10.1)	NR		
Lester 2019 Secondary analysis of RCT Serious risk of bias	N = 586	Trauma patients	Level 1 trauma centre (US)	Hypothermia (< 35.5°C) RBC units required	N = 399 9.9 (11.4)	N = 187 6.3 (7.9)	RR 0.90 (0.89, 0.92)	No significant difference <i>p</i> = 0.00		
Acid-base status	Acid-base status									
Tran 2018 SR <i>Critically low quality</i>	N = NR (2 Coh) Vandromme 2010 Vandromme 2011	Adult patients with traumatic torso injuries (civilian and military)	Trauma (US)	Lactate risk of massive transfusion	NR	NR	OR 4.10 (2.50, 6.74) OR 5.20 (3.51, 7.71) OR 3.13 (1.96, 5.00)	Favours severe lactic acidosis p < 0.0001 Substantial heterogeneity $l^2 = 62\% (p < 0.10)$		
Baxter 2016 SR <i>Critically low quality</i>	N = 1093 (3 Coh) Regnier 2012 Baron 2004 Ipekci 2013	Trauma patients	Trauma/ED (US, France)	Lactate (mmol/L)	In all trauma patients, increased lactate and lactate clearance were found to predict major haemorrhage, defined as blood transfusion of more than 6 red cell units within 24 hours and/or death from haemorrhagic shock. Increased lactate was also found to be associated with increased blood loss in penetrating torso trauma patients. Two studies found that raised lactate was associated with blood component requirements, but this was not significant in a study that only looked at patients with isolated extremity injuries.					
Javali 2017 Prospective Coh <i>Critical risk of bias</i>	N = 100	Trauma patients at risk of haemodynamic compromise	Tertiary care centre emergency department (India)	Arterial lactate (mmol/L) Transfusion requirement	lactate < 2.9 mmol/L 24.6%	lactate ≥2.9 mmol/L 85.7%		p < 0.001		
				Base deficit (mEq/L)	< 12 mEq/L 36.4%	≥ 12 mEq/L 78.3%		P = NR		
Ionised calcium										
Vasudeva 2021 SR <i>Critically low quality</i>	N = 817 (2 Coh)	Trauma patients	Trauma centres (US, Australia)	Hypocalcaemia (< 1.11 mmol/L) Magnotti 2011 ≥5 U	NR NR/332 (17.1)	NR NR/259 (7.1)	NR	NR p = 0.005		
				≥10 U Vasudeva 2020	NR/332 (8.2) 75/113 (62.5)	NR/259 (2.2) 45/113 (37.5)	NR NR	<i>p</i> = 0.017 <i>p</i> < 0.001		
Study ID	Sample size	Patient	Setting	Predictor	Results					
---	--	---	---	---	--	--	---	---	--	--
Study design ^a	(no. of trials) included in analysis	population	(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b		
Shih 2019 SR <i>Critically low quality</i>	N = 591 (1 Coh) Magnotti 2011	Civilian trauma patients	Regional trauma centre (US)	Hypocalcaemia (i-Ca < 1.0 mmol/L) Multiple transfusions Massive transfusion	NR/332 (17.1) NR/332 (8.2)	NR/259 (7.1) NR/259 (2.2)	OR 2.294 (1.053, 4.996) NR	p = 0.005 ° p = 0.017 °		
Moore 2020 Secondary analysis of RCTs <i>Moderate risk of</i> <i>bias</i>	N = 160	Patients with traumatic haemorrhagic shock from blunt or penetrating injuries	2 trauma centres (US)	Hypocalcaemia (i-Ca < 1.0 mmol/L) RBC in 24 hrs Plasma in 24 hrs Platelet in 24 hrs CRYO in 24 hrs	Median (IQR) N = 70 2040 (2-10) 2 (1-7) 0 (0-1) 0 (0-1)	Median (IQR) N = 90 1 (0-5) 2 (0-4) 0 (0-0) 0 (0-0)	NR	p = 0.0002 p = 0.007 p = 0.30 p = 0.0003		
Haemoglobin										
Tran 2018 SR <i>Critically low quality</i>	N = NR (3 Coh) Callcut 2013 Paulus 2014 Vandromme 2011	Adult patients with traumatic torso injuries (civilian and military)	Trauma (US)	Haemoglobin (< 110 to 115 g/L) Significant bleeding	NR	NR	OR 3.78 (1.97, 7.26) OR 2.40 (1.82, 3.16) OR 2.56 (2.02, 3.25) OR 1012 (6.01 17.05)	Favours low haemoglobin p < 0.0001 Substantial heterogeneity $l^2 = 92\%$ ($p < 0.00001$)		
Shih 2019 SR <i>Critically low quality</i>	N = 2349 (5 Coh) Callcut 2011 Callcut 2013 Leemann 2010 Schochl 2011 Schreiber 2007	Trauma patients	Trauma (US, Iraq, Switzerland, Austria)	Haemoglobin (< 11 g/dL) RBC transfusion volume (≥ 10 units in 6 hrs)	NR	NR	NR OR 3.1 (1.2, 8.4) OR 1.8 (1.3, 2.5) OR 18.18 (2.73, 125.00) ROC AUC 0.87 (0.83, 0.91) OR 7.7 (5.0, 11.9)	NR		
Platelet count				·						
Levy 2017 SR <i>Critically low quality</i>	N = 30 735 (7 Coh) Arnold 2006 Fayed 2013 McGrath 2008 Premaratne 2001 Tanaka 2014 Wu 2014 van Hout 2017	Adult perioperative surgical patients receiving platelet transfusion (cardiac, acute aortic dissection, liver transplant)	Surgical (US, Canada, Egypt, Netherlands)	Platelet count	Included studies used transfusion, including viscoelastic measured between the 2 public ×10 ⁹ /L for interventio mixed medical/surgid <100 ×10 ⁹ /L accompa Hout 2017). Different platelet dos studies, ranging from Fayed 2013). Wu 2014 measurement for trig administered.	d different measureme g platelet count, bleed s. The platelet counts u ations, ranging from a nal treatment in a stud cal intensive care unit inied by bleeding in ca ses per transfusion wer n one to 6 to 12 units (va and McGrath 2008 dia ggering transfusion or	ents to trigger platelet ng (visual measure), and ised as triggers varied median of 51 (IQR 26–68) dy evaluating patients in a (Arnold 2006) to a trigger of rdiac surgery patients (van e administered in all an Hout 2017, Tanaka 214, d not report a dose of transfusion	Heterogeneity between studies was so substantial that quantitative synthesis was not possible.		

Study ID	Sample size	Patient	Setting	Predictor	Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b	
					Premaratne 2001 observed a change in bleeding time (NR) between cardiopulmonary bypass patients who received less than 10 units or or more units of platelet transfusions.				
PT/INR									
Tran 2018 SR <i>Critically low quality</i>	N = NR (2 Coh) Callcut 2013 Vandromme 2011	Adult patients with traumatic torso injuries (civilian and military)	Trauma (US)	INR and risk of MT	NR	NR	OR 4.16 (2.57, 6.73) OR 3.40 (2.48, 4.66) OR 5.61 (2.57, 6.73)	Favours coagulopathy p < 0.00001 Substantial heterogeneity $l^2 = 60\% (p < 0.11)$	
Shih 2019 SR <i>Critically low quality</i>	N = 1803 (2 Coh) Callcut 2013 Schreiber 2007	Trauma patients	Trauma (US, Iraq)	INR (> 1.5) ≥ 10 units RBC / 24 hr	NR	NR	OR 2.1 (1.4, 3.1) OR 5.9 (3.5, 10.2)	NR	
APTT									
Haas 2015 SR <i>Critically low quality</i>	N = NR (2 Coh) Mannucci 1982 Murray 1998	Patients with critical bleeding	Trauma (US, Italy)	APTT	Mannucci 1982 repor patients who underv recommended FFP t	rted PT > 1.2 times norm went major surgery and transfusion if PT or aPT	nal or APTT > 1.25 times norm d received massive transfusio T is >1.5 times prolonged dur	al were found in 93% of on. Murray 1998 ing massive transfusion.	
Fibrinogen level									
Shih 2019 SR <i>Critically low quality</i>	N = 625 (1 Coh) Nakamura 2017	Trauma patients	Trauma (Japan)	Fibrinogen (≤ 190 mg/dL)	NR	NR	OR 0.931 (0.898, 0.963)	NR	
Abdul-Kadir 2014 SR <i>Critically low quality</i>	N = NR (4 Coh) Charbit 2007 Cortet 2012 Peyvandi 2012 Rouse 2006	Obstetrics	Obstetrics (US, France, Italy)	Fibrinogen (> 2 g/L)	Three studies assess Two studies (Charbit women who develop decreased fibrinoger measure of blood los concentrate, which h	ed the association betw 2007, Cortet 2012) rep bed more severe PPH. I n is an independent an ss. Rouse 2006 notes th nas been used in obste	ween PPH requiring transfus orted a lower (≤ 2 g/L) mean Peyvandi 2012 however was u Ind measurable predictor of se nat low fibrinogen may requir trics for the management of	ion and fibrinogen levels. plasma fibrinogen level in unable to determine if evere PPH or simply a re transfusion of fibrinogen PPH since 1948.	
McQuilten 2017 Retrospective Coh Moderate risk of	N = 2829	Patients who received massive transfusion	Hospitals (Australia, New Zealand)	Fibrinogen <1g/L 1.0-1.9g/L >4g/L	RBC transfusion 11 (8-18) 9 (7-13) 7 (6-9)	2.0-4.0g/L (reference) 8 (6-11)	NR	p < 0.001	
DIAS				Fibrinogen <1g/L 1.0-1.9g/L >4g/L	FFP transfusion 8 (4-14) 2040 (4-10) 4 (2-6)	2.0-4.0g/L (reference) 5 (3-8)	NR	<i>p</i> < 0.001	

Study ID	Sample size	Patient	Setting	Predictor	Results						
Study design ^a	(no. of trials) included in analysis	population	ppulation (Location) F		Predictor n/N (%)	No predictor n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b			
				Fibrinogen <1g/L 1.0-1.9g/L >4g/L	PLT transfusion 2 (1-4) 2 (1-3) 0 (0-1)	2.0-4.0g/L (reference) 1 (0-2)	NR	<i>p</i> < 0.001			
				Fibrinogen <1g/L 1.0-1.9g/L >4g/L	CRYO or FC 4.2 (2.1-8.5) 3.8 (0-6.8) 0.0 (0.0-1.9)	2.0-4.0g/L (reference) 1.7 (0.0-4.2)	NR	<i>p</i> < 0.001			

APTT, activated partial thromboplastin time; AUC, area under the curve; Cl, confidence interval; Coh, cohort; ED, emergency department; FFP, fresh frozen plasma; hrs, hours; INR, international normalised ratio; IQR, interquartile range; MT, massive transfusion; N, number; NR, not reported; OR, odds ratio; PPH, postpartum haemorrhage; PT, prothrombin time; RBC, red blood cells; RCT, randomised controlled trial; ROC, receiver operating characteristic; RR, relative risk; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if Phet >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

4.3 Defined major haemorrhage protocol (Question 2)

Question 2 - (Interventional)

In patients with critical bleeding, what is the effectiveness of major haemorrhage protocols?

4.3.1 Methods

This review examined the effects of defined major haemorrhage protocols (MHPs) versus no defined MHPs in patients with critical bleeding (i.e. major haemorrhage that is lifethreatening and is likely to result in the need for massive transfusion) as outlined in Figure 4.5 below.

Figure 4.5 PICO criteria: Question 2 – defined MHPs



CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; MHP, major haemorrhage protocol; PLT, platelets; RBC, red blood cells

Notes:

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. c. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2013, noting primary studies published prior to 2013 that had been included in a systematic review were also eligible for inclusion. There were no restrictions applied to study sample size. Screening of RCTs and nonrandomised studies (with concurrent or noncurrent controls) and observational cohort studies was performed.

Studies of lower-level evidence were only screened for primary outcomes insufficiently addressed in systematic reviews. Assuming all relevant primary studies had been identified in the included systematic review; screening for lower-level evidence was not conducted. An updated literature search was conducted in August 2019 and again in September 2021 to identify any new systematic reviews meeting the eligibility criteria.

Assuming all relevant primary studies have been identified in the included systematic review studies; the systematic screen for RCTs was limited to studies published from September 2019. This is based on the literature search date of the most recent identified systematic review (Consunji 2020), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting.

4.3.2 Summary of evidence

4.3.2.1 Systematic reviews

There were 8 systematic reviews (37, 57-63) identified in the systematic review and handsearching process that assessed the effects of a defined MHP versus no defined MHP in patients with critical bleeding. Two other systematic reviews (Hallet 2013, Johansson 2012) were identified in the search, but these did not provide any additional data to that provided by the included reviews, thus are not discussed further in this report (see **Appendix B** for a complete list of excluded studies that met the PICO criteria for this question).

Four reviews (Cannon 2017, Mitra 2013, Vogt 2012, Consunji 2020) we focused on critical bleeding in the trauma setting, 3 reviews (Maw 2018, Kamyszek 2019, Kinslow 2020) included paediatric trauma patients and one review (Sommer 2019) included both trauma and non-trauma patients.

The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.6. A matrix illustrating the overlap of studies included in each review is provided in Table 4.7.

Review ID Quality	Study design	Population	Intervention	Comparator	Outcomes
Trauma sett	ing				
Consunji 2020 (57) <i>High</i>	SR / MA of cohort and observational studies (17 studies)	Trauma patients receiving or anticipated to receive a massive blood transfusion	MTP	No MTP	Mortality
Cannon 2017 (61) <i>Moderate</i>	SR / MA of RCTs and cohort studies (11 studies)	Severely injured adult patients requiring blood transfusion and/or ISS >25	MT/DCR protocol	no MT/DCR protocol	Mortality Transfusion volumes
Mitra 2013 (62) <i>Moderate</i>	SR / MA of clinical trials, clinical studies, guidelines and meta-analyses (10 studies)	Adult trauma patients in the initial trauma resuscitation phase	After implementation of a predefined MHP	Before implementation of a predefined MHP	Mortality

Table 4.6 Characteristics and quality of systematic reviews by clinical setting: defined MHPs versus no defined MHPs

Review ID Quality	Study design	Population	Intervention	Comparator	Outcomes
Vogt 2012 (63) <i>Moderate</i>	SR / MA of comparative observational studies (7 studies)	Adult civilian trauma patients expected to require MT	Formal TTP	no formal TTP	Mortality Transfusion volumes
Non-trauma	setting				
Sommer 2019 (59) <i>Critically</i> <i>low</i>	SR / MA of RCTs, observational studies and retrospective studies (12 studies)	Adult non-trauma (perioperative, obstetric, gastrointestinal bleeding and vascular emergencies) patients with massive bleeding	MHP	No MHP	Mortality Transfusion volume Wastage of blood components
Paediatric s	etting				
Kinslow 2020 (58) <i>Critically</i> <i>low</i>	SR of observational studies (3 studies)	Paediatric trauma patients	MTP	No MTP	Mortality
Kamyszek 2019 (37) <i>Critically</i> <i>low</i>	SR / MA of clinical trials, clinical studies, guidelines and meta-analyses (3 studies)	Paediatric patients receiving an MT	Formal MHP implementation	Before MTP implementation	Mortality Transfusion volume
Maw 2018 (60) Critically Iow	SR / MA of RCTs, observational studies and retrospective studies (5 studies)	Paediatric patients, younger than 18 years, with traumatic injury requiring blood transfusion	Predetermined ratios of blood component therapy, including MHP	Transfusion at physician discretion	Mortality

DCR, damage control resuscitation; FFP, fresh frozen plasma; ICU, intensive care unit; ISS, injury severity score; LOS, length of stay; MA, meta-analysis; MHP, major haemorrhage protocol; MT, massive transfusion; MTP, massive transfusion protocol; PLT, platelet; RBC, red blood cells; RCTs, randomised controlled trials; SR, systematic review; TTP, trauma transfusion pathway

		Trau	ıma																											
	Study ID	Campion 2014	Cinat 1999	Cotton 2009	Cotton 2011	Duchesne 2010	Fox 2008	Kahn 2008	Nascimento 2013	0'Keefe 2008	Riskin 2009	Shaz 2010	Cotton 2008	Dirks 2010	Johansson 2008	Simmons 2010	Sinha 2013	Sisak 2012	Dente 2009	Johansson 2009	Vogt 2009	Ball 2013	Brinck 2016	Hwang 2018	Maciel 2015	Noorman 2016	Nunn 2017	Soderlund 2017	van der Meij 2019	Zaydfudim 2009
۵	Consunji 2020			×						×	~	~		~		~	~	×				Х	~	×	×	~	~	Х	×	Х
≥	Cannon 2017	~	Х	~	Х	~	~	X	X	~	~	~																		
svie	Mitra 2013			~						~	~	~	Х	~	Х	~	~	~												
Å	Vogt 2012			~						~	1		~						~	~	~									

Table 4.7 Overlap table primary studies included in the identified systematic reviews: defined MHPs versus no defined MHPs

		Mixed	lixed trauma and non-trauma											Paediatrics				
	Study ID	Bauman Kreuziger 2014	Balvers 2015	Chay 2016	Dutta 2017	Goodnough 2011	Gutierrez 2012	Johansson 2007	Martinez-Calle 2016	McDaniel 2013	Morse 2012	Sinha 2013	Wijaya 2016	Chidester 2012	Edwards 2015	Hendrickson 2012	Hwu 2016	Nosanov 2013
۵	Kinslow 2020													~		~	~	
N	Sommer 2019	Х	~	Х	~	Х	Х	~	~	~	Х	Х	Х					
svie	Kamyszek 2019													~		~	~	
R	Maw 2018													~	Х	~		Х

✓ = study included in this review

X = study did not meet the inclusion criteria for this review

--- = study identified by the systematic review authors but not included (no usable data)

4.3.2.2 Randomised controlled trials

No RCTs were found in the systematic review and handsearching process that examined the effects of defined MHPs versus no defined MHPs in patients with critical bleeding.

4.3.2.3 Observational and cohort studies

There were 29 nonrandomised studies identified in the included systematic reviews that examined the effects of defined MHPs versus no defined MHPs on mortality and transfusion volumes in patients with critical bleeding and were considered relevant to this review. The main characteristics and quality of included cohort studies and relevant outcomes assessed are summarised in Table 4.8.

Table 4.8	Characteristics and quality of observational and cohort studies by clinical setting:
	defined MHPs versus no defined MHPs

Review ID Risk of bias	Study design	Population N	Intervention	Comparator	Outcomes
Mixed traum	a and non-traun	na setting			
Balvers 2015 <i>Moderate</i>	Retrospective cohort	Adult trauma and non- trauma patients with massive bleeding ^a N=547	MTP	Pre-MTP *historical control	Mortality Transfusion volume
Trauma setti	ing				
van der Meij 2019 <i>Moderate</i>	Retrospective cohort, SC	Civilian trauma patients with haemorrhage requiring MT N=101	MHP	No MHP	Mortality MOF Ventilator days LOS Transfusion volume
Hwang 2018 <i>Moderate</i>	Retrospective cohort	Civilian trauma patients with haemorrhage requiring MT N=190	MHP	No MHP	Mortality Transfusion volume
Nunn 2017 <i>Moderate</i>	Retrospective cohort	Civilian trauma patients with haemorrhage requiring MT N=447	MHP	No MHP	Mortality Ventilator days LOS
Brink 2016 <i>Moderate</i>	Retrospective cohort	Trauma patients ≥16 years with ISS >15 requiring MT N=352	MHP	No MHP	Mortality Transfusion volume
Noorman 2016 <i>Moderate</i>	Retrospective cohort, SC	Trauma patients requiring MT N=201	МНР	No MHP	Mortality Transfusion volume LOS
Maciel 2015 <i>Moderate</i>	Retrospective cohort, SC	Patients with abdominal aorta injuries N=46	MHP	No MHP	Mortality Duration on mechanical ventilator LOS Complications
Campion 2013 <i>High</i>	Retrospective cohort, SC	Adult patients requiring surgery within 24 hours of admission to Level I trauma centre	DCR	Pre-DCR	Mortality Transfusion volume

Review ID Risk of bias	Study design	Population N	Intervention	Comparator	Outcomes
		N=216			
Sinah 2013 <i>Moderate</i>	Prospective cohort	Civilian trauma patients with haemorrhage requiring MT N=152	MHP	No MHP	Mortality Transfusion volume
Sisak 2012 <i>Moderate</i>	Retrospective cohort	Civilian trauma patients with haemorrhage requiring MT N=58	MHP	No MHP	Mortality
Dirks 2010 <i>Moderate</i>	Before-after with historical control	Civilian trauma patients with haemorrhage requiring MT N=66	MHP	No MHP	Mortality Transfusion volume
Duchesne 2010 <i>High</i>	Retrospective cohort, SC	Trauma patients requiring transfusion during surgery N=196	DCR	Pre-DCR	Mortality
Shaz 2010 <i>Moderate</i>	Prospective cohort	Civilian trauma patients with haemorrhage requiring MT N=224	MHP	No MHP	Mortality Transfusion volume
Simmons 2010 <i>Moderate</i>	Retrospective cohort	Military trauma patients with haemorrhage requiring MT N=777	MHP	No MHP	Mortality Transfusion volume
Riskin 2009 <i>Moderate</i>	Retrospective cohort	Civilian trauma patients with haemorrhage requiring MT N=77	MHP	No MHP	Mortality Transfusion volume
Cotton 2008 <i>Moderate</i>	Prospective cohort	Civilian trauma patients with haemorrhage requiring MT N=211	MHP	No MHP	Mortality Transfusion volume
Cotton 2009 <i>Moderate</i>	Prospective cohort	Civilian trauma patients with haemorrhage requiring MT N=264	MHP	No MHP	Mortality Transfusion volume
O'Keefe 2008 <i>Moderate</i>	Retrospective before-after study	Civilian trauma patients with haemorrhage requiring MT N=178	MHP	No MHP	Mortality Transfusion volume
Dente 2009 <i>High</i>	Prospective cohort, SC	Trauma patients requiring MT N=157	МНР	No MHP	Mortality Transfusion volume
Vogt 2009 <i>High</i>	Retrospective cohort	Trauma patients requiring MT N=46	МНР	No MHP	Mortality Transfusion volume
Johansson 2009 <i>High</i>	Before-after with historical control	Trauma patients requiring MT N=832	МНР	No MHP	Mortality Transfusion volume
Fox 2008 High	Retrospective 2 cohort case- control study	Military patients with life-threatening haemorrhage N=40	DCR	Pre-DCR	Transfusion volume

Review ID Risk of bias	Study design	Population N	Intervention	Comparator	Outcomes
Non-trauma	setting				
Dutta 2017 <i>Moderate</i>	Retrospective cohort	Obstetric patients with massive bleeding N=62	MTP	No MTP	Mortality Transfusion volume
Martinez- Calle 2016 <i>Moderate</i>	Retrospective cohort	Adult non-trauma patients with massive bleeding (perioperative, obstetric, gastrointestinal bleeding and vascular emergencies) N=304	MTP	No MTP	Mortality Transfusion volume
McDaniel 2013 <i>Moderate</i>	Retrospective cohort	Adult non-trauma (perioperative, obstetric, gastrointestinal bleeding and vascular emergencies) patients with massive bleeding N=64	MTP	No MTP	Mortality Transfusion volume Wastage of blood components
Johansson 2007 <i>Moderate</i>	Retrospective cohort	Adult patients with ruptured abdominal aortic aneurysm N=132	MTP	No MTP	Mortality Transfusion volume
Paediatric se	etting				
Hwu 2016 <i>High</i>	Retrospective cohort, SC	Paediatric trauma patients receiving MT N=43	MHP	Pre-MHP	Mortality Transfusion LOS
Chidester 2012 <i>High</i>	Prospective cohort, SC	Paediatric trauma patients receiving MT N=55	MHP	Pre-MHP	Mortality Morbidity
Hendrickson 2012 <i>High</i>	Retrospective cohort, SC	Paediatric trauma patients receiving MT N=102	MHP	Pre-MHP	Mortality LOS

DCR, damage control resuscitation; LOS, length of stay; MHP, major haemorrhage protocol; MOF, multiple organ failure; MT, massive transfusion; MTP, massive transfusion protocol; SC, single centre

a. Patients from a variety of settings including surgery (63%), internal medicine (13%), other (11%), trauma (9%), obstetric (4%)

Trauma setting

The 4 systematic reviews (Consunji 2020, Cannon 2017, Mitra 2013, Vogt 2012) identified a total of 21 observational studies in adult trauma patients relevant to this review question.

Consunji (2020) identified 14 relevant studies (Brink 2016, Cotton 2009, Dirks 2010, Shaz 2010, Hwang 2018, Maciel 2015, Noorman 2016, Riskin 2009, O'Keefe 2008, Nunn 2017, Simmons 2010, Sinah 2013, Sisak 2012, van der Meij 2019). Cannon 2017 identified 3 additional studies (Champion 2013, Duchesne 2010, Fox 2008) not identified elsewhere. Mitra (2013) identified 8 studies, all of which were also identified in Consunji (2020). Vogt 2012 identified 4 additional studies (Cotton 2008, Dente 2009, Johansson 2009, Vogt 2009) not identified elsewhere.

Most studies were carried out in single and multicentre trauma centres in the US, Canada, Denmark and Australia. Overall, the systematic reviews judged included observational

studies to be moderate to high risk of bias due to study design, data collection and adjustments for confounding.

The 14 studies identified in Consunji 2020 and Mitra 2013 evaluated the effect of implementing defined MHPs in adult trauma patients, assessing patient outcomes before and after implementation. The 3 studies identified in Cannon 2017 evaluated the implementation of an MHP with or without a formal damage control resuscitation protocol in adult trauma patients. The 4 studies identified in Vogt 2012 only included studies in adult civilian trauma patients. Identified systematic reviews did not define MHPs and most acknowledged differences between included observational studies in the definitions and triggers of MHPs.

All systematic reviews aimed to evaluate the association between MHPs and mortality as a patient outcome, pooling analysis of included studies. Cannon 2017 and Vogt 2012 reported pooled volume of red blood cells (RBC) transfused as a surrogate endpoint for the total blood components transfused. Cannon 2017, Mitra 2013 and Vogt 2012 reported volume of other blood components transfused.

Mixed trauma and non-trauma setting

One systematic review (Sommer 2019) assessing the effects of MHPs in both trauma and non-trauma settings was identified in the literature. Sommer 2019 reported outcome data of one observational study (Balvers 2015) including a mixture of trauma (8.8%) and non-trauma patients requiring massive transfusion (defined as the administration of 5 of more units of RBC within 12-hours).

Balvers 2015 was carried out in an academic medical centre in the Netherlands and was judged by review authors to have high risk of bias due to study design, data collection and adjustments for confounding.

Sommer 2019 also identified 12 observational studies in the non-trauma settings. Of these, only 4 were included in the quantitative analysis. Studies were in the non-trauma setting and included patients with bleeding due to obstetric complications (Dutta 2017), ruptured abdominal aortic aneurysm (Johansson 2007) or perioperative surgery (McDaniel 2013, Martinez-Calle 2016).

Studies included patients with major bleeding who required transfusion, which was defined as 4 or more units of RBC (Dutta 2017), 10 or more units of RBC (McDaniel 2013, Johansson 2007) or the replacement of whole blood volume in 24-hours, 50% of volume in 3-hours or blood loss of more than 1500 mL in 10 minutes (Martinez-Calle 2016).

The 4 included observational studies (Dutta 2017, McDaniel 2013, Martinez-Calle 2016, Johansson 2007) were carried out in single centre settings in the US, Denmark and Spain. Overall, included observational studies were judged by review authors to be high risk of bias due to study design and confounding.

Paediatric setting

Three reviews (Kinslow 2020, Kamyszek 2019, Maw 2018) assessing MHPs in paediatric patients were identified in the literature. All reviews identified several observational studies, however, only 3 met our inclusion criteria. Two systematic reviews (Kinslow 2020 and Kamyszek 2019) identified all 3 studies. One review (Maw 2018) only identified 2 of the 3 studies as one was published after the search date.

All 3 included observational studies (Chidester 2013, Hendrickson 2012, Hwu 2016) were carried out in single paediatric trauma centres in the US, Afghanistan and Iraq. Overall, risk of bias for included studies was judged by review authors to be high risk due to study design and selection bias.

4.3.3 Results

4.3.3.1 MHP compared to no MHP

Mortality

A summary of the evidence relating to mortality in patients with critical bleeding treated with an MHP is presented in Table 4.9.

All identified systematic reviews reported a weak association between defined MHPs and mortality (latest timepoint) in trauma patients (Consunji 2020, Cannon 2017, Mitra 2013, Vogt 2012, Sommer 2019). For all other subgroups, the identified systematic reviews reported no association between a defined MHPs and mortality (Kamyszek 2019, Kinslow 2020, Maw 2018, Sommer 2019).

Pooled data from observational studies included in this review (Figure 4.6) showed the mortality rate at 24-hours in patients with critical bleeding to be no different among those who were managed using an MHP (192/1114, 17.2%) compared with those who did not (158/777, 20.3%) (OR 0.88; 95% CI 0.61, 1.27; p = 0.09; random effect, $l^2 = 42\%$).

A meta-analysis of data from observational studies included in this review (Figure 4.7) showed the mortality rate (latest timepoint) in patients with critical bleeding to be lower among those who were managed using an MHP (926/2927, 31.6%) compared with those who were not (977/2492, 39.2%) with the odds ratio (OR) of 0.71 observed (95% CI 0.57, 0.87; p = 0.001; random effect, I² = 62%).

Among patients with blunt and penetrating trauma, a meta-analysis of data from observational studies included in this review (Figure 4.6) showed little to no difference in mortality (24 hours) among those who were managed using an MHP (131/618, 21.2%) compared with those who were not managed using an MHP (122/412, 29.6%) with the odds ratio (OR) of 0.79 observed (95% CI 0.56, 1.11; p = 0.17; random effect, $l^2 = 15\%$). However, mortality at the latest timepoint reported (typically up to 30-days or upon hospital discharge) was lower among patients who were managed using an MHP (717/2278, 31.5%) compared with those who were not (786/1948, 40.3%) with the OR of 0.67 observed (95% CI 0.53, 0.85; p = 0.001; random effect, $l^2 = 63\%$).(*GRADE: Very low*)

Among paediatric trauma patients, the mortality rate of 41.7% (43/103) who were managed using an MHP was not significantly different from the mortality rate of 36.1% (35/97) observed among those who did not receive MHPs. This corresponded to an OR of 1.31 (95% CI 0.71, 2.42; p = 0.38; random effect, I² = 5%). (*GRADE: Very low*)

Among non-trauma patients who were managed using an MHPs, the mortality rate 22% (42/191) was not significantly different from the mortality rate of 35.7% (91/255) observed among those who were not managed using an MHPs. This corresponded to an OR of 0.54 (95% CI 0.25, 1.15; p = 0.11; random effects, $I^2 = 61\%$). Including the hospital-wide study that assessed trauma and non-trauma patients, the mortality rate (latest timepoint) of 30.4% (166/546) was slightly lower than the mortality rate of 34.9% (156/447) observed among patients whose transfusions were not guided by an MHP, but the effect estimates were inconsistent and the lower bound of the confidence interval suggests no important association (OR 0.67; 95% CI 0.35, 1.29; p = 0.23; $I^2 = 74\%$).(*GRADE: Very low*)

Figure 4.6 Forest plot of comparison: MHPs vs no MHPs, outcome: Mortality, 24 hours

	MHF	•	no M⊦	IP		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
1.1.1 Tauma setting							
Cotton 2009 (Coh, trauma)	39	125	55	141	18.9%	0.71 [0.43, 1.18]	
Noorman 2016 (Coh, trauma)	3	144	6	57	5.4%	0.18 [0.04, 0.75]	
O'Keeffe 2008 (Coh, trauma)	27	132	9	46	11.5%	1.06 [0.46, 2.45]	_
Shaz 2010 (Coh, trauma)	38	142	27	84	16.7%	0.77 [0.43, 1.39]	
Sisak 2012 (Coh, trauma)	10	28	9	30	8.1%	1.30 [0.43, 3.89]	
van der Meij 2019 (Coh, trauma)	14	47	16	54	11.3%	1.01 [0.43, 2.37]	
Subtotal (95% CI)		618		412	71.9%	0.79 [0.56, 1.11]	•
Total events	131		122				
Heterogeneity: $Tau^2 = 0.03$; $Chi^2 = 5.86$, $df = 5$ (P = 0.32);	l² = 15%						
Test for overall effect: Z = 1.37 (P = 0.17)							
1.1.2 Mixed trauma and non-trauma setting							
Balvers 2015 (RCoh, 9% trauma, 63% surgical)	52	355	23	192	18.4%	1.26 [0.75, 2.13]	- -
Subtotal (95% CI)		355		192	18.4%	1.26 [0.75, 2.13]	•
Total events	52		23				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.86 (P = 0.39)							
1.1.3 Non-trauma setting							
Dutta 2017 (RCoh. Obstetrics)	0	23	0	39		Not estimable	
Martinez-Calle 2016 (RCoh. surgical & nonsurgical) (1)	1	92	7	96	2.7%	0.14 [0.02, 1.16]	
McDaniel 2013 (RCoh, non-trauma)	8	26	6	38	7.0%	2.37 [0.71, 7.92]	+
Subtotal (95% CI)		141		173	9.7%	0.66 [0.04, 11.26]	
Total events	9		13				
Heterogeneity: Tau ² = 3.47; Chi ² = 5.50, df = 1 (P = 0.02);	l² = 82%						
Test for overall effect: $Z = 0.29$ (P = 0.77)							
Total (95% CI)		1114		777	100.0%	0.88 [0.61, 1.27]	•
Total events	192		158			• • •	
Heterogeneity: $Tau^2 = 0.12$: $Chi^2 = 13.68$, df = 8 (P = 0.09)	: ² = 42%	'n					
Test for overall effect: $Z = 0.66$ (P = 0.51)	,,	-					0.01 0.1 1 10 100
Test for subaroup differences: $Chi^2 = 2.21$, df = 2 (P = 0.33)	3), ² = 9.5	5%					Favours MHP Favours no MHP
Footnotes							

(1) Data reported from most recent protocol updates (i.e. Group 2B) used for MHP group.

	MHF		no Mł	ΗP		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% CI
1.2.1 Trauma setting							
Brinck 2016 (Coh, truama)	35	206	39	146	4.8%	0.56 [0.34, 0.94]	
Campion 2013 (Coh, trauma)	27	99	42	117	4.5%	0.67 [0.37, 1.20]	
Cotton 2008 (Coh, trauma)	77	117	48	94	4.6%	1.84 [1.06, 3.22]	
Cotton 2009 (Coh, trauma)	54	125	88	141	5.0%	0.46 [0.28, 0.75]	
Dente 2009 (Coh, trauma)	25	73	46	84	4.1%	0.43 [0.23, 0.82]	_ - _
Dirks 2010 (Coh, trauma)	47	156	24	97	4.5%	1.31 [0.74, 2.33]	- -
Duchesne 2010 (Coh, trauma)	19	72	56	124	4.2%	0.44 [0.23, 0.82]	
Hwang 2018 (Coh, trauma)	43	126	35	64	4.3%	0.43 [0.23, 0.79]	
Johansson 2009 (Coh, trauma)	17	50	46	82	3.7%	0.40 [0.19, 0.84]	
Maciel 2015 (Coh, trauma)	9	17	25	29	1.6%	0.18 [0.04, 0.75]	
Noorman 2016 (Coh, trauma)	10	144	13	57	3.0%	0.25 [0.10, 0.62]	
Nunn 2017 (Coh, trauma)	83	208	113	239	5.6%	0.74 [0.51, 1.08]	
O'Keeffe 2008 (Coh, trauma)	69	132	23	46	4.0%	1.10 [0.56, 2.14]	_
Riskin 2009 (Coh, trauma)	7	37	18	40	2.6%	0.29 [0.10, 0.80]	
Shaz 2010 (Coh, trauma)	63	132	42	84	4.7%	0.91 [0.53, 1.58]	
Simmons 2010 (Coh, trauma)	81	426	84	351	5.8%	0.75 [0.53, 1.05]	
Sinha 2013 (Coh, trauma)	24	83	16	69	3.7%	1.35 [0.65, 2.81]	
Sisak 2012 (Coh, trauma)	13	28	12	30	2.5%	1.30 [0.46, 3.68]	
van der Meij 2019 (Coh, trauma)	14	47	16	54	3.2%	1.01 [0.43, 2.37]	
Subtotal (95% CI)		2278		1948	76.7%	0.67 [0.53, 0.85]	\blacklozenge
Total events	717		786				
Heterogeneity: Tau ² = 0.16; Chi ² = 48.19, df = 18 (P = 0.0	001); l² =	63%					
Test for overall effect: Z = 3.30 (P = 0.0010)							
1.2.2 Mixed trauma and non-trauma setting							
Balvers 2015 (RCoh, 9% trauma, 63% surgical)	124	355	65	192	5.7%	1.05 [0.72, 1.52]	Ť
Subtotal (95% CI)		355		192	5.7%	1.05 [0.72, 1.52]	•
Total events	124		65				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.25$ (P = 0.80)							
1.2.3 Non-trauma softing							
	0	00	0	00		Net estimable	
Dutta 2017 (RCon, Obstetrics)	0	23	0	39	0.70/		
Jonansson 2007 (RCon, ruptured AAA)	17	50	46	82	3.7%	0.40 [0.19, 0.84]	
Manunez-Calle 2016 (RCon, surgical & nonsurgical) (1)	12	92	29	90	3.1% 2.7%	0.30 [0.10, 0.73]	
Subtotal (05% CI)	13	20 101	10	აი 255	2.7% 101%	0.54 [0.50, 3.75]	
	40	131	01	200	10.170	0.04 [0.20, 1110]	-
10 (a) events	42		91				
Therefore overall effect: $7 = 1.60 (P = 0.11)$	1 0170						
Test for overall effect. $\Sigma = 1.00 (F = 0.11)$							
1.2.4 Paediatric setting							
Chidester 2012 (Coh. paediatric trauma)	15	33	10	22	2.4%	1.00[0.34, 2.95]	
Hendrickson 2012 (Coh. paediatric trauma)	20	53	11	49	3.1%	2.09 [0.88, 5.00]	
Hwu 2016 (Coh, paediatric trauma)	-0	17	14	26	2.0%	0.76 [0.22, 2.59]	
Subtotal (95% CI)	•	103		97	7.6%	1.31 [0.71, 2.42]	◆
Total events	43		35				-
Heterogeneity: Tau ² = 0.01; Chi ² = 2.10, df = 2 (P = 0.35);	l ² = 5%						
Test for overall effect: $Z = 0.88$ (P = 0.38)							
· · · · ·							
Total (95% CI)		2927		2492	100.0%	0.71 [0.57, 0.87]	♦
Total events	926		977				
Heterogeneity: Tau ² = 0.16; Chi ² = 66.10, df = 25 (P < 0.0	001); l² =	62%					
Test for overall effect: Z = 3.26 (P = 0.001)							U.U.I U.I I IU 100 Favours MHP Favours on MHP
Test for subgroup differences: $Chi^2 = 7.66$, df = 3 (P = 0.0	5), l² = 60	.8%					

Figure 4.7 Forest plot of comparison: MHPs vs no MHPs, outcome: Mortality, latest timepoint

(1) Data reported from most recent protocol updates (i.e. Group 2B) used for the MHP group.

Footnotes

Table 4.9 Results for defined MHP versus no defined MHP: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study designª	(no. of trials) included in analysis	population	(Location)			MHP n/N (%)	No MHP n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b		
Mixed trauma	and non-trauma settir	ngs									
Sommer 2019	N = 547	Adult patients with	Trauma and non-	MHP vs pre-	Mortality, 24-hour	52/355 (15)	23/192 (12)	OR 1.26 (0.75, 2.13) ^c	p = 0.39°		
SR	(1 Coh)	major bleeding	trauma (surgical,	MHP	Mortality, 30-days	124/355 (35)	62/192 (34)	OR 1.05 (0.72, 1.52)°	<i>p</i> = 0.80 ^c		
Critically low quality	Balvers 2015		general medicine). SC (Netherlands)								
Trauma setting	3										
Consunji 2020	N = 1030	Various patients	Trauma, MC and	MHP vs no	Mortality, 24-hour	131/618 (21.2)	122/412 (29.6)	OR 0.79 (0.56, 1.11) ^c	No significant difference		
SR	(6 Coh)	with haemorrhage	SC (US, Australia)	MHP					<i>p</i> = 0.17 ^c		
High quality		requiring							Mild heterogeneity		
		transfusion							I ² = 15%		
					Noorman 2016	3/144 (2)	6/57 (11)	OR 0.18 (0.04, 0.75)	Favours MHP, p = 0.004		
					O'Keeffe 2008	27/132 (20.5)	9/46 (19.6)	OR 1.06 (0.46, 2045)	μ = 0.185 p > 0.05		
					Shaz 2010	38/142 (29)	27/84 (32)	OR 0.85 (0.47, 1.54)	p = 0.28		
					Sisak 2012	10/28 (35.7)	9/30 (30)	OR 1.30 (0.43, 3.89)	p = 1.00		
	NL 0014	Maria	Traverse MO and	MUDIE	van der Meij 2019	14/47 (29.8)	16/54 (29.6)	OR 1.01 (0.43, 2.37)	p = 0.99		
	N = 3314	various patients	Frauma, MC and	MHP vs no	Mortality, 30-days	552/1867 (29.6)	548/1447 (37.9)	OR 0.69 (0.53, 0.89)°	Favours MHP		
	(14 Con)	requiring	Australia)	IVITE	or latest timepoint				$p = 0.004^{\circ}$		
		transfusion	Australia						Substantial		
		a anorasion							recerogeneity		
					Princk 2016	25/200 (10.0)	20/140 (20 5)	00050/004 004	F = 57%		
					Cotton 2009	35/206 (16.9) 54/125 (43.2)	39/146 (26.5) 88/141 (62.4)	OR 0.56 (0.34, 0.94) OR 0.46 (0.28.0.75)	Favours MHP, $p = 0.032$		
					Dirks 2010	47/156 (30.1)	24/97 (24.7)	OR 1.31 (0.74, 2.33)	p = 0.382		
					Shaz 2010	63/132 (48)	42/84 (50)	OR 0.91 (0.53, 1.58)	<i>p</i> = 0.47		
					Hwang 2018 Maciel 2015	43/126 (34.1)	35/64 (54.7)	OR 0.48 (0.26, 0.88)	Favours MHP, p = 0.007		
					Noorman 2016	10/144 (7)	13/57 (23)	OR 0.25 (0.06, 0.91) OR 0.25 (0.10, 0.62)	Favours MHP, p = 0.002		
					Riskin 2009	7/37 (19)	18/40 (45)	OR 0.29 (0.00, 0.80)	Favours MHP, p = 0.02		
					O'Keeffe 2008	69/132 (52.3)	23/46 (50.0)	OR 1.10 (0.56, 2.14)	<i>p</i> = <i>NR</i>		
					Simmons 2010	83/208 (40.1)	113/239 (47.2)	OR 0.77 (0.53, 1.12)	p = 0.1/32 p = 0.115		
					Sinha 2013	24/83 (29)	16/69 (23)	OR 0.75 (0.33, 1.05) OR 0.77 (0.16, 3.75)	p = 0.43		
					Sisak 2012	13/28 (46)	12/30 (40)	OR 1.30 (0.46, 3.68)	p = 0.791		
					van der Meij 2019	14/47 (29.8)	16/54 (29.6)	OR 1.16 (0.53, 2.58)	p = 0.99		

Study ID	y ID Sample size Patient Setting Comparison Ou				Outcome	Results					
Study designª	(no. of trials) included in analysis	population	(Location)			MHP n/N (%)	No MHP n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b		
Cannon 2017 SR <i>Moderate</i> quality	N = 412 (2 Coh) Campion 2013 Duchesne 2010	Adult patients with severe trauma at risk of death from haemorrhage (civilian and military)	SC (US)	MHP vs pre- MHP	Mortality, 30-days or in hopsital	46/171 (26.9) 27/99 (27.3) 19/72 (26.4)	98/241 (40.7) 42/117 (35.9) 56/124 (45.2)	OR 0.55 (0.36, 0.84) ^c 0.67 (0.37, 1.20) 0.44 (0.23, 0.82)	Favours MHP p = 0.006 No heterogeneity $l^2 = 0\% (p = 0.33)$		
Vogt 2012 SR Moderate quality	N = NR (4 Coh)	Adult trauma patients (civilian) requiring massive transfusion	Trauma (Canada, Denmark, US)	MHP vs no MHP	Mortality, 30-day or in-hospital Cotton 2008° Dente 2009 Johansson 2009 Voqt 2009	48/94 (51.1) 25/73 (34.2)' 17/50 (34)' NR	77/117 (65.8) 46/84 (55)' 46/82 (56)' NR	Adjusted RR estimate 0.51 (0.29, 0.90) Unadjusted RR estimates 0.69 (0.52, 0.91) ^r 0.65 (0.51, 0.82) ^r 0.64 (0.32, 1.27)	NR		
Non-trauma se	etting		1								
Sommer 2019 SR Critically low quality	N = 314 (3 Coh)	Adult patients with major bleeding	Non-trauma (surgical, obstetric). SC (Denmark, Spain, US)	MHP vs pre- MHP	Mortality, 24-hour McDaniel 2013 Martinez -Calle 2016 ^d Dutta 2017	9/141 (6.4)° 8/26 (30.8) 1/92 (0.5) 0/23 (0)	13/173 (7.5) ^c 6/38 (15.8) 7/96 (7.3) 0/39 (0)	OR 0.66 (0.04, 11.26) ^c 2.37 (0.71, 7.92) ^c 0.14 (0.02, 1.16) ^c NE	No significant difference $p = 0.77^{\circ}$ Substantial heterogeneity $l^2 = 82\% (p = 0.02)^{\circ}$		
	N = 446 (4 Coh)				Mortality, 30-days Johansson 2007 McDaniel 2013 Martinez-Calle 2016 ^a Dutta 2017	42/191° 17/50 (34) 13/26 (50.0) 12/92 (13) 0/23 (0)	91/255 ^c 46/82 (56) 16/38 (42.1) 29/96 (30.2) 0/39 (0)	OR 0.54 (0.25, 1.15) ^c 0.40 (0.19, 0.84) ^c 1.38 (0.50, 3.75) ^c 0.35 (0.16, 0.73) ^c NE	No significant difference $p = 0.11^{\circ}$ Substantial heterogeneity $l^2 = 61\% (p = 0.08)^{\circ}$		
Paediatrics, tra	auma setting										
Kinslow 2020 SR Critically low quality	N = 328 (3 Coh) Hwu 2016 Chidester 2012 Hendrickson 2012	Paediatric trauma patients with various ISS.	Paediatric trauma (US, Iraq, Afghanistan)	MHP vs no MHP	Mortality, in- hospital	43/103 (41.7) ^c 8/17 (47.1) 15/33 (45) 20/53 (38)	35/97 (36.1) ^c 14/26 (53.8) 10/22 (45) 11/49 (23)	OR 1.31 (0.71, 2.42) ^c 0.76 (0.22, 5.29) ^c 1.00 (0.34, 2.95) ^c 2.09 (0.88, 5.00) ^c	p = 0.38 ^c I ² = 5% (p = 0.35) ^c		

Cl, confidence interval; Coh, cohort study; ISS, injury severity score; MC, multicentre; MHP, major haemorrhage protocol; MT, massive transfusion; NR, not reported; OR, odds ratio; RR, relative risk; SC single centre; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4.

d. Martinez–Calle 2016 report MHP mortality for group A and B based on protocol updates in different years. For this review, data reported from the most recent protocol updates (i.e. Group 2B) are used and compared to pre-MHP. Where necessary, data from primary study was sourced.

e. Cotton 2008 reported estimate adjusted for age, gender, mechanism of injury, TRISS, and 24-hour transfusion requirements.

f. Data sourced from primary study.

Transfusion volume

Red blood cells

A summary of the evidence reported in the identified systematic reviews relating to transfusion volume of other blood components in patients with critical bleeding is presented in Table 4.10.

The systematic reviews (Sommer 2019, Cannon 2017, Mitra 2013, Vogt 2012) suggest only limited conclusions can be drawn from the available evidence, with a nonsignificant reduction in the volume of RBC transfused (less than one red cell unit saved).

A meta-analysis of data from observational studies included in this review (see Figure 4.8) revealed no important difference in the volume of RBC transfusion in patients with critical bleeding who received MHPs (n=1909) compared with those who did not (n=1593), with around 1.2 units of RBC saved. The overall standardised mean difference (SMD) was -0.12 (95% CI -0.31, 0.07; p = 0.22; random effect, I² = 74%).

Among patients with blunt and penetrating trauma, no important difference in the volume of RBC transfusion was observed among those who received MHPs compared with those who did not, around 1.2 units of RBC saved (SMD–0.13; 95% CI –0.33, 0.07; p = 0.10; random effect, I² = 77%).

Among non-trauma patients, only one study contributed data, which reported no difference in the volume of RBC transfused among those who were managed using an MHP compared with those who were not (less than one unit saved) (SMD 0.04; 95% CI – 0.46, 0.54; p = 0.88). These data are consistent with the studies that reported median volumes that could not be included in the meta-analysis.

		MHP	MHP no MHP Std. Mean		Std. Mean Difference	Std. Mean Difference			
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% C	I IV, Random, 95% CI
1.3.1 Trauma setting									
Cotton 2008 (Coh, trauma)	18.8	11.2	94	18.8	11.2	117	12.3%	0.00 [-0.27, 0.27]	-+-
Fox 2008 (Coh, trauma)	23	18	16	12	6.4	24	5.4%	0.87 [0.21, 1.54]	
Johansson 2009 (Coh, trauma)	18	12.6	442	19.2	15.8	390	15.2%	-0.08 [-0.22, 0.05]	
O'Keeffe 2008 (Coh, trauma)	11.8	11.8	132	15.5	15.5	46	10.8%	-0.29 [-0.62, 0.05]	
Riskin 2009 (Coh, trauma)	20.5	2.6	37	23.9	2.7	40	7.7%	-1.27 [-1.76, -0.78]	
Shaz 2010 (Coh, trauma)	24	14	132	23	14	84	12.2%	0.07 [-0.20, 0.34]	
Simmons 2010 (Coh, trauma)	17	12	426	19	11	351	15.1%	-0.17 [-0.31, -0.03]	
Sinha 2013 (Coh, trauma) (1)	0	0	28	0	0	30		Not estimable	
Sisak 2012 (Coh, trauma)	19.8	8.5	28	19.6	9.7	30	7.4%	0.02 [-0.49, 0.54]	
Vogt 2009 (Coh, trauma)	23	10.7	23	25	15.2	23	6.4%	-0.15 [-0.73, 0.43]	
Subtotal (95% CI)			1358			1135	92.4%	-0.13 [-0.33, 0.07]	•
Heterogeneity: Tau ² = 0.06; Chi ² = 34.19, df = 8 (P < 0.00) Test for overall effect: Z = 1.27 (P = 0.20)	001); l² =	: 77%							
1.3.2 Non-trauma setting (or mixed)									
Balvers 2015 (RCoh, 9% trauma, 63% surgical) (2)	0	0	355	0	0	192		Not estimable	
Dutta 2017 (RCoh, Obstetrics) (3)	0	0	23	0	0	39		Not estimable	
Johansson 2007 (RCoh, ruptured AAA) (4)	0	0	55	0	0	93		Not estimable	
Martinez-Calle 2016 (RCoh, surgical & nonsurgical) (5)	0	0	92	0	0	96		Not estimable	
McDaniel 2013 (RCoh, non-trauma)	12.6	11.5	26	12.2	9	38	7.6%	0.04 [-0.46, 0.54]	
Subtotal (95% CI)			551			458	7.6%	0.04 [-0.46, 0.54]	\bullet
Heterogeneity: Not applicable									
Test for overall effect: $Z = 0.15$ (P = 0.88)									
Total (95% CI)			1909			1593	100.0%	-0.12 [-0.31, 0.07]	•
Heterogeneity: Tau ² = 0.06; Chi ² = 34.57, df = 9 (P < 0.00	001); l² =	74%							
Test for overall effect: Z = 1.22 (P = 0.22)									
Test for subgroup differences: $Chi^2 = 0.38$, df = 1 (P = 0.5)	54), l² = (0%							

Figure 4.8 Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, red blood cells

Footnotes (1) Median RBCs transfused: MHP 14 (11-21) vs pre-/no MHP 16 (12-20) (2) Median RBCs transfused: MHP 8 (7-13) vs pre-/no MHP 8 (6-12)

(3) Median RBCs transfused: MHP 7 (5-9) vs pre-/no MHP 6 (5-8)

(4) Median RBCs transfused for ICU patients: MHP 2 (0-30) vs pre-/no MHP 6 (0-54)
(5) Median RBCs transfused: MHP 10 (18-12) vs pre-no/MHP 9 (8-14)

Table 4.10	Results for defined MHP versus no defined MHP: Patients with critical bleeding - Transfusion volume, re	d blood cells
------------	---	---------------

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study designª	(no. of trials) included in analysis	population	(Location)			MHP Mean ± SD (n)	No MHP Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b		
Mixed trauma a	and non-trauma se	ettings									
Sommer 2019 SR <i>Critically low</i> <i>quality</i>	N = 1261 (6 Coh)	Adult patients (aged 18 years or older) with major bleeding in trauma and non- trauma setting	Trauma and non- trauma (surgical, obstetrics, general). Single centres (Australia, The Netherlands, US)	MHP vs pre- MHP	RBC transfusion volume, units McDaniel 2013 Balvers 2015 Sinha 2013 Dutta 2017 Martinez-Calle 2016 ^a Johansson 2007 (operating room) (intensive care unit)	Mean ± SD (n 12.6 ± 11.5 (26) Median (IQR) 8 (7-13) (n355) 14 (11-21) (n83) 7 (5-9) (n23) 10 (8-12) (n92) NR 2 (0-30)(n50)	Mean ± SD (n 12.2 ± 9.0 (38) Median (IQR) 8 (6–12) (n192) 16 (12–20) (n69) 2040 (5–8) (n39) 9 (8–14) (n96) NR	MD 0.40 (–4.87, 5.67) ^c	No significant difference p = 0.864 p = 0.279 p = NR p = 0.85 p = NR Favours MHP, $p < 0.05$		
						2 (0-30)(1130)	6 (0–54)(n82)				
Trauma setting	1										
Cannon 2017 SR <i>Moderate</i> <i>quality</i>	N = 551 (4 Coh)	Adult patients with severe trauma at risk of death from haemorrhage (civilian and military)	SC (Iraq, US)	MHP vs pre- MHP	RBC transfusion volume, units, 24 hours Fox 2008 O'Keeffe 2008 Riskin 2009 Shaz 2010	n=217 23 ± 18 (16) 11.8 ± 11.8 (132) 20.5 ± 2.6 (37) 24 ± 14 (132)	n=194 12 ± 6.4 (24) 15.5 ± 15.5 (46) 23.9 ± 2.7 (40) 23 ± 14 (84)	MD 0.36 (-4.54,3.83) 11.00 (1.82, 20.18) -3.70 (-8.61, 1.21) -3.40 (-4.58, -2.22) 1.00 (-4.54, 3.83)	No significant difference p = 0.87 Substantial heterogeneity I² = 78% (p = 0.004)		
Mitra 2013 SR Moderate quality	N = 835 (2 Coh)	Adult trauma patients (civilian and military) in the initial trauma resuscitation phase	SC (US, Australia)	MHP vs pre- MHP	RBC transfusion volume Simmons 2010 Sisak 2012	NR 17 ± 12 (426) 19.8 ± 8.5 (28)	NR 19 ± 11 (351) 19.6 ± 9.7 (30)	NR	NR		
Vogt 2012 SR Moderate quality	N = 1267 (4 Coh) Cotton 2008 Johansson 2009 Vogt 2009	Adult patients requiring massive transfusion due to civilian trauma	SC, trauma (Canada, US, Denmark)	MHP vs no MHP	RBC transfusion volume, units at 24 hours	NR 18.8 ± 11.2 (94) 18 ± 12.6 (442) 23 ± 10.7 (23)	NR 18.8 ± 11.2 (117) 19.2 ± 15.8 (390) 25 ± 15.2 (23)	MD -0.90 (-2.51, 0.71) ^c MD 0.00 (-3.04, 3.04) MD -1.20 (-3.16, 0.76) -2.00 (-9.60, 5.60)	No significant difference p = 0.27° No heterogeneity I² = 0% (p = 0.78)°		

Study ID Study	Sample size	Patient	Setting	Comparison Outcome	Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			MHP Mean ± SD (n)	No MHP Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b		
Non-trauma se	Non-trauma setting – no identified studies reported on outcome of interest										
Paediatrics, trauma setting - no identified studies reported on outcome of interest											
Cl. confidence interval: Coh. cohort: MD, mean difference: MHP, maior haemorrhage protocol: NR, not reported: RBC, red blood cell: SC, single centre: SD, standard deviation: SR, systematic review: US, United											

States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4.

d. Martinez-Calle 2016 report MHP mortality for group A and B based on protocol updates in different years. For this review, data reported from the most recent protocol updates (i.e. Group 2B) are used and compared to pre-MHP. Where necessary, data from primary study was sourced.

Fresh frozen plasma

A summary of the evidence reported in the identified systematic reviews relating to transfusion volume of FFP in patients with critical bleeding is presented in Table 4.11.

Included systematic reviews (Sommer 2019, Mitra 2013, Vogt 2012) suggest only limited conclusions can be drawn from the available evidence, reporting slight nonsignificant reduction in the volume of FFP transfused (less than one FFP unit saved).

A meta-analysis of data from observational studies included in this review (see Figure 4.9) shows no difference in the volume of FFP transfused in patients with critical bleeding who had an MHP (n=1314) compared with those who did not (n=1081) (less than one unit save). The overall standardised mean difference (SMD) was –0.09 (95% CI –0.41, 0.23; p = 0.57; random effect, I² = 93%). Heterogeneity was substantial with differences in triggers activating MHPs varying between studies.

Among non-trauma patients, only one study contributed data, which reported no difference in the volume of FFP transfused among those who were managed using an MHP compared with those who were not (less than one unit saved) (SMD 0.04; 95% CI – 0.46, 0.53; p = 0.89).

	l	MHP		no MHP			;	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.4.1 Trauma setting									
Johansson 2009 (Coh, trauma)	13.5	12.3	442	12.1	15.2	390	14.5%	0.10 [-0.03, 0.24]	-
Simmons 2010 (Coh, trauma)	8	8	426	14	11	351	14.4%	-0.63 [-0.78, -0.49]	+
Cotton 2008 (Coh, trauma)	9.9	7	94	12.4	12.5	117	13.4%	-0.24 [-0.51, 0.03]	
Shaz 2010 (Coh, trauma)	13	12	132	8	7	84	13.4%	0.48 [0.20, 0.76]	
O'Keeffe 2008 (Coh, trauma)	5.7	5.4	132	8.7	6.9	46	12.7%	-0.51 [-0.85, -0.17]	
Sisak 2012 (Coh, trauma)	9.4	5.8	28	8.1	6.2	30	10.7%	0.21 [-0.30, 0.73]	
Vogt 2009 (Coh, trauma)	14	8	23	15	10.1	23	10.0%	-0.11 [-0.69, 0.47]	
Riskin 2009 (Coh, trauma) (1)	10.7	0	37	12.3	0	40		Not estimable	
Subtotal (95% CI)			1314			1081	89.1%	-0.11 [-0.46, 0.24]	•
Heterogeneity: Tau ² = 0.19; Chi ² = 82	2.01, df =	6 (P <	0.0000	01); l² =	93%				
Test for overall effect: Z = 0.61 (P = 0).54)								
1.4.2 Non-trauma setting (or mixed	l)								
McDaniel 2013 (RCoh, non-trauma)	9.2	8	26	8.9	8.7	38	10.9%	0.04 [-0.46, 0.53]	<u> </u>
Subtotal (95% CI)			26			38	10.9%	0.04 [-0.46, 0.53]	•
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.14 (P = 0).89)								
Total (95% CI)			1340			1119	100.0%	-0.09 [-0.41, 0.23]	· · · •
Heterogeneity: Tau ² = 0.18; Chi ² = 82	2.70, df =	7 (P <	0.0000	01); l² =	92%				-2 -1 0 1 2
Test for overall effect: Z = 0.57 (P = 0).57)								Favours MHP Favours pre-MHP
Test for subgroup differences: Chi ² =	0.22, df =	= 1 (P	= 0.64)	, l² = 0%	þ				
Footnotes									

Figure 4.9 Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, FFP

(1) Standard deviation and patient numbers not reported

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results			
Study	(no. of trials)		(Location)			МНР	No MHP	Risk estimate	Statistical significance
designa	analysis					Mean ± SD (n)	Mean ± SD (n)	(95% CI)	<i>p</i> -value
	anarysis								Heterogeneity ^b
Mixed trauma	and non-trauma set	tings							
Sommer 2019	N = 1261	Adult patients (≥18	SC	MHP vs pre-	FFP transfusion				
SR	(6 Coh)	years) with major	(Australia, The	MHP	volume, units				
Critically low		bleeding in trauma	Netherlands, US)		McDaniel 2013	9.2 ± 8.0 (26)	8.9 ± 8.7 (38)	MD 0.30 (-3.84, 4.44) ^c	No significant difference
quality		and non-trauma							<i>p</i> = 0.631
		setting			Balvers 2015	Median (IQR)	Median (IQR)		n - 0.224
					Sinha 2013	6 (4-11)(355)	6 (3–9) (192)		p = 0.224 p = NR
					Dutta 2017	10 (7-17)(83)	6 (5-10)(69)		p = 0.28
					Martinez-Calle 2016d	5 (3-9)(92)	5 (3-9) (96)		<i>p</i> = 0.376
					Johansson 2007	- (/(/	- (/ (/		
					(intensive care unit)	4 (2-16)(50)	0 (0-3)(82)		Favours MHP, $p < 0.05$ Eavours MHP, $p < 0.05$
Troumo cottine	7					0 (0-4)	T (U-6)		
Trauma securi	y								1
Mitra 2013	N = 1532	Adult trauma patients	SC	MHP vs pre-	FFP transfusion	NR	NR	NR	
SR	(7 Coh)	(civilian) in the initial	(US, Australia,	MHP	volume				
Moderate		trauma resuscitation	Denmark)		Riskin 2009	10.7 ± NR	12.3 ± NR		No significant difference
quality		pnase			O'Keeffe 2008	9.4 ± 5.8 (28) 5.7 ± 5.4 (132)	8.1 ± 6.2 (30) 8.7 ± 6.9 (46)		p = NR Eavours MHD p = ND
					Simmons 2010	8 + 8 (426)	14 + 11 (351)		Favours no MHP $\rho = NR$
					Shaz 2010	13 ± 12 (132)	8 ± 7 (84)		Favours no MHP, $p = NR$
					Dirks 2010	NŔ	NR		p = NR
					Sinha 2013	NR	NR		p = NR
Vogt 2012	N = 1089	Adult trauma patients	SC, trauma	MHP vs no	FFP transfusion	NR	NR	MD -0.50 (-3.37,	No significant difference
SR	(3 Coh)	(civilian) requiring	(Canada, US,	MHP	volume, unit at 24			2.37)	$p = 0.22^{c,e}$
Moderate		massive transfusion	Denmark)		hours				Substantial heterogeneity
quality					Cotton 2008	9.9 ± 7 (94)	12.4 ± 12.5 (117)	-2.50 (-5.17, 0.17)	$l^2 = 0\% (p = 0.06)^{c, e}$
					Johansson 2009	13.5 ± 12.3 (442)	12.1 ± 15.2 (390)	1.40 (-0.49, 3.29)	
					Vogt 2009	14 ± 8 (23)	15 ± 10.1 (23)	-1.00 (-6.27, 4.27)	
Non-trauma se	etting – no identified	studies reported on outco	ome of interest						

Table 4.11 Results for defined MHP versus no defined MHP: Patients with critical bleeding – Transfusion volume, fresh frozen plasma

Paediatrics, trauma setting - no identified studies reported on outcome of interest

Cl, confidence interval; Coh, cohort; FFP, fresh frozen plasma; IQR, interquartile range; MD, mean difference; MHP, major haemorrhage protocol; NR, not reported; SC, single centre; SD, standard deviation; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. Martinez-Calle 2016 report MHP mortality for group A and B based on protocol updates in different years. For this review, data reported from the most recent protocol updates (i.e. Group 2B) are used and compared to pre-MHP. Where necessary, data from primary study was sourced.

e. Data from Cotton 2008 not included in the meta-analysis due to substantial heterogeneity (I2=64%, p = 0.06)

Platelets

A summary of the evidence reported in the identified systematic reviews relating to transfusion volume of platelets in patients with critical bleeding is presented in Table 4.12.

The systematic reviews (Sommer 2019, Mitra 2013, Vogt 2012) suggest only limited conclusions can be drawn from the available evidence, with a nonsignificant increase in the volume of platelets transfused (more than one plasma unit wasted).

A meta-analysis of data from observational studies included in this review (see Figure 4.10) revealed an increase in the volume of platelet transfusion in patients with critical bleeding who were managed using an MHP (n=2049) compared with those who were not (n=1666) (more than 3.5 units) (SMD 0.54; 95% CI –0.26, 1.33; p = 0.19; random effect, I² = 99%). Heterogeneity was substantial with effect estimate likely to be largely influenced by differences between studies for MHP activation.

Figure 4.10 Forest plot of comparison: MHPs vs no MHPs, outcome: Transfusion volume, platelets

	I	MHP		nc	MHP			Std. Mean Difference		Std. Mean	Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI		IV, Rando	om, 95% Cl
1.5.1 Trauma and non-trauma setting											
Cotton 2008 (Coh, trauma) (1)	31	3.7	94	6.8	7.2	117	12.3%	4.08 [3.61, 4.56]			
Dirks 2010 (Coh, trauma) (2)	0	0	156	0	0	97		Not estimable			
Johansson 2009 (Coh, trauma) (3)	5	4.2	442	1.7	2	390	12.9%	0.98 [0.84, 1.13]			+
O'Keeffe 2008 (Coh, trauma) (4)	1.1	1.3	132	3.8	5.2	46	12.6%	-0.94 [-1.29, -0.59]			
Riskin 2009 (Coh, trauma) (5)	2.3	0	37	2.8	0	40		Not estimable			
Shaz 2010 (Coh, trauma)	2	2	132	2	1	84	12.7%	0.00 [-0.27, 0.27]		-	+ -
Simmons 2010 (Coh, trauma)	1	2	426	2	3	351	12.9%	-0.40 [-0.54, -0.26]			
Sinha 2013 (Coh, trauma) (6)	0	0	28	0	0	30		Not estimable			
Sisak 2012 (Coh, trauma)	10.1	6.5	28	5.8	6.8	30	12.2%	0.64 [0.11, 1.17]			
Vogt 2009 (Coh, trauma) (7)	14	8	23	15	10.1	23	12.1%	-0.11 [-0.69, 0.47]		_	+
Subtotal (95% CI)			1498			1208	87.7%	0.60 [-0.27, 1.47]		-	◆
Heterogeneity: Tau ² = 1.34; Chi ² = 475.71, df = 6 (P < 0.00	001); l²	= 999	6								
Test for overall effect: Z = 1.35 (P = 0.18)											
1.5.2 Non-trauma setting (or mixed)											
Balvers 2015 (RCoh, 9% trauma, 63% surgical) (8)	0	0	355	0	0	192		Not estimable			
Dutta 2017 (RCoh, Obstetrics) (9)	0	0	23	0	0	39		Not estimable			
Johansson 2007 (RCoh, ruptured AAA) (10)	0	0	55	0	0	93		Not estimable			
Martinez-Calle 2016 (RCoh, surgical & nonsurgical) (11)	0	0	92	0	0	96		Not estimable			
McDaniel 2013 (RCoh, non-trauma)	7.2	6.7	26	6.5	8.6	38	12.3%	0.09 [-0.41, 0.59]		-	<u>+</u>
Subtotal (95% CI)			551			458	12.3%	0.09 [-0.41, 0.59]		•	•
Heterogeneity: Not applicable											
Test for overall effect: Z = 0.34 (P = 0.73)											
Total (95% CI)			2049			1666	100.0%	0.54 [-0.26, 1.33]			
Heterogeneity: Tau ² = 1.27; Chi ² = 476.43, df = 7 (P < 0.00	1001); l²	= 999	6					-	-4	-2	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
Test for overall effect: Z = 1.32 (P = 0.19)										Favours MHP	Favours pre-MHP
Test for subgroup differences: $Chi^2 = 1.00$, df = 1 (P = 0.32)), l ² = 0 ⁶	%									
Footnotes											
(1) Missing SD data											
(2) Median (IQR): MHP 0 (0-0) vs pre-no MHP 1 (0-4)											
(3) SD data missing											
(4) SD data missing											
(5) Standard devation and patient numbers not reported											
(6) Median (IQR): MHP 3 (2-4) vs pre-/no MHP 2 (1-3)											
(7) SD data missing											
(8) Median (IQR): MHP 2 (0-4) vs pre-/no MHP 2 (1-3)											
(9) Median (IQR): MHP 0 (0-0.6) vs pre-/no MHP 0 (0-0.6)											
(10) Median (IQR) in operating room: MHP 11 (2-42) vs pr	e-/no Mł	HP 7	(0-46): ;	and in IO	CU: M	HP 2 (0)-12) vs pr	e-/no MHP 4 (0-32)			
(11) Median (IQR): MHP 1 (0-2) vs pre-/no MHP 1 (0-2)	(11) Median (IQR): MHP 1 (0-2) vs pre-/no MHP 1 (0-2)										

Table 4.12	Results for defined MHP versus no defined MHP: Patients with critical bleedin	g – Transfusion volume, platelets
------------	---	-----------------------------------

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study	(no. of trials)	population	(Location)			MHP	No MHP	Risk estimate	Statistical significance
design ^a	included in					Mean ± SD (n)	Mean ± SD (n)	(95% CI)	<i>p</i> -value
	analysis								Heterogeneity ^b
Mixed trauma a	and non-trauma settin	igs							
Sommer 2019	N = 863 (6 Coh)	Adult patients	SC	MHP vs pre-MHP	PLT transfusion			NR	NR
SR		with major	(Australia, The		volume, units				
Critically low		bleeding in	Netherlands,		McDaniel 2013	7.2 ± 6.7 (26)	6.5 ± 8.6 (38)		
quality		trauma and non-	US)			Median (IOR)	Median (IOR)		
		trauma setting			Balvers 2015	2 (0-4) (n355)	2 (1-3) (n192)		
					Sinha 2013	3 (2–4) (n83)	2 (1–3) (n69)		
					Dutta 2017 Martinoz Callo 2016c	0 (0–0.6) (n23)	0 (0–0.6) (n39)		
					Johansson 2007	1 (0–2) (n)	1 (0–2)		
					(operating room)	11 (2-42)	7 (0-46)		
					(intensive care unit)	2 (0–12)	4 (0-32)		
Trauma setting	1		-		-		-	-	
Mitra 2013	N = 1532 (7 Coh)	Adult trauma	SC	MHP vs pre-MHP	PLT transfusion			NR	NR ^d
SR	Riskiin 2009	patients (civilian)	(US, Australia,		volume	2.3 ± NR	2.8 ± NR		
Moderate	O'Keefe 2008	in the initial	Denmark)			2 ± 2 (132) 11 ± 13(132)	2 ± 1 (84) 3 8 ± 5 2 (46)		
quality	Sisak 2012	trauma				10.1 ± 6.5 (28)	5.8 ± 6.8 (30)		
	Simmons 2010	resuscitation				1 ± 2 (426)	2 ± 3 (351)		
		phase				Madian (IOD)	Madian (IOD)		
	Dirks 2010 (total)								
	Sinha 2013 (24 hours)					3 (2-4) ^r	2 (1-3)		
Mart 2012	N 425 (2 Cab)				DI Through using			ND	NDd
vogt 2012	N = 435 (3 CON)	Adult trauma	SC, trauma	WIHP VS NO MIHP	Volume units at	21. 27(/04)	CO. 70(417)	NR	NR ^s
SR	Johansson 2009	requiring massive	(Canada, Dopmosk, US)		24 hours	5.0 ± 4.2^{t} (94)	$6.8 \pm 7.2^{\circ} (117)$ $1.7 \pm 2.0^{\circ} (390)$		
moderate	Vogt 2009	transfusion	Denmark, US)		24 110015	3 ± NR (23)	2 ± NR (23)		
Non trauma co	tting no identified at	udias reported an a	uteeme of intere	ot					

Paediatrics, trauma setting - no identified studies reported on outcome of interest

Cl, confidence interval; Coh, cohort; IQR, interquartile range; MD, mean difference; MHP, major haemorrhage protocol; NR, not reported; PLT, platelets; SC single centre; SR, systematic review; US, United States a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been

considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. Meta-analysis not conducted due to substantial heterogeneity (I2=89%-100%)

e. The MHP implemented in Martinez-Calle 2016 was updated during the study period (MHP 1: 2007–2009 and MHP 2: 2010–2012). The p-value is pre-MHP vs MHP 1 vs MHP 2 f. Sourced from primary study

Wastage of blood components

A summary of the evidence reported in the identified systematic reviews relating to wastage of blood components in patients with critical bleeding is presented in Table 4.13.

Systematic review suggests only limited conclusions can be drawn from the available evidence, reporting significant wastage of platelets following MHP implementation. As noted by Sommer 2018, MHP termination may influence wastage of blood components. Furthermore, overactivation of MHPs can also lead to wastage of blood components. Due to the limited data, no meta-analysis was performed.

Time to delivery of blood components

A summary of the evidence reported in the identified systematic reviews relating to time to delivery of blood components in patients with critical bleeding is presented in Table 4.14.

Systematic review suggests only limited conclusions can be drawn from the available evidence, reporting significant reduction in the time to first FFP administration following MHP implementation. Due to the limited data, no meta-analysis was performed.

Table 4.13 Results for defined MHP versus no defined MHP: Patients with critical bleeding – Wastage of blood components

Study ID	Sample size	Patient population	Setting (Location)	Comparison	Outcome	Results					
Study design ^a	(no. of trials) included in analysis					MHP units/total issued (%)	No MHP units/total issued (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b		
Mixed trauma and non-trauma settings – not identified studies reported on outcome of interest											
Trauma setting – no identified studies reported usable data											
Non-trauma settin	ng										
Sommer 2019	N = 164 (1 Coh)	Adult patients with	Non-trauma	MHP vs pre-	Wastage of RBC	3/613 (0.5)	3/848 (0.35)	OR 1.39 (0.28, 6.89) ^c	No significant difference		
SR	McDaniel 2013	major bleeding due to	(surgical, general	MHP					p = 0.69 ^c		
Critically low		non-trauma	medicine) SC (US)		Wastage of FFP	1/406 (0.25)	4/553 (0.72)	OR 0.34 (0.04, 3.04)°	No significant difference		
quality									<i>p</i> = 0.33 ^c		
					Wastage of PLT	39/304 (12.8)	29/358 (8.1)	OR 1.67 (1.01, 2.77) ^c	Favours no MHP		
									<i>p</i> = 0.05 ^c		
Prodictrice trauma softing not identified studies reported on outcome of interest											

Paediatrics, trauma setting - not identified studies reported on outcome of interest

Cl, confidence interval; Coh, cohort; FFP, fresh frozen plasma; MHP, major haemorrhage protocol; OR, odds ratio; PLT, platelet; RBC, red blood cell; SC, single centre; SR, systematic review; US, United States a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been

considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4.

Table 4.14 Results for defined MHP versus no defined MHP: Patients with critical bleeding – Time to delivery of blood components

Study ID	Sample size	Patient	Setting Comparison Outcome Results								
Study design ^a	(no. of trials) included in analysis	population	(Location)			MHP Mean ± SD (n)	No MHP Mean ± SD (n)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b		
Mixed trauma and non-trauma settings - no identified studies reported on outcome of interest											
Trauma settings -	no identified studie	es reported on outco	me of interest								
Non-trauma settings – no identified studies reported on outcome of interest											
Paediatrics, traun	na setting										
Karnyszek 2019	N = 43 (1 Coh)	Paediatric trauma	MC	MHP vs pre-	Hours to first blood	0.9 ± NR (17)	0.8 ± NR (26)	NR	No significant difference		
SR	Hwu 2016	patients receiving	(US, Iraq,	MHP	component				<i>p</i> = 0.688		
Critically low		MT	Afghanistan)		Hours to first RBC	1.4 ± NR (17)	0.8 ± NR (26)	NR	No significant difference		
quality									<i>p</i> = 0.180		
					Hours to first PLT	4.4 ± NR (17)	6 ± NR (26)	NR	No significant difference		
								<i>p</i> = 0.421			
	N = NR (2 Coh)				Hours to first FFP			NR	Favours MHP		
	Hwu 2016					1.0 ± NR	2.7 ± NR		<i>p</i> = 0.005		
	Henderickson 2012					0.8 ± NR	3.3 ± NR		p <0.001		

Cl, confidence interval; Coh, cohort; FFP, fresh frozen plasma; MHP, major haemorrhage protocol; MC, multicentre; MT, massive transfusion; NR, not reported; PLT, platelet; RBC, red blood cell; SD, standard deviation; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if Phet >0.1 and I2 <25%; (ii) mild heterogeneity if I2 <25%; moderate heterogeneity if I2 between 25–50%; substantial heterogeneity I2 >50%.

4.4 Dose, timing and ratio (algorithm) of RBC to blood component therapy (Question 3)

Question 3 - (Interventional)

In patients with critical bleeding, what is the optimal dose, timing and ratio (algorithm) of RBC to blood component therapy (FFP, platelets, cryoprecipitate or fibrinogen concentrate) to reduce morbidity, mortality and transfusion?

4.4.1 Methods

This question investigated the optimal dose, timing and ratio of different ratios of red blood cells (RBC) to blood component therapy including fresh frozen plasma (FFP), platelets (PLT) and cryoprecipitate (CRYO) in patients with critical bleeding, outlined in Figure 4.11 below.

Figure 4.11 PICO criteria: Question 3 – dose, timing and ratio of different ratios of red blood cells



ARDS, acute respiratory distress syndrome; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; MHP, major haemorrhage protocol; PLT, platelets; RBC, red blood cells; TE, thromboembolic event

Notes:

a. 1 vs 1; 2 vs 2; etc.

b. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months).

c. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. d. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

This question was included in the previous version of the guidelines, which investigated literature published up to 2009. Further, a comprehensive systematic review conducted by Monash included a search of the literature published between 2009 and 2015. However, this comprehensive review only included RCTs. Hence, for this question, evidence published after June 2015 was considered but any articles published prior to 2015 that had been identified in a systematic review were also included. An updated literature search was conducted in September 2021 to identify any new systematic reviews meeting the eligibility criteria. Assuming all relevant primary studies have been identified in the included systematic review studies; the systematic screen for RCTs was limited to studies published from January 2021. This is based on the literature search dates of the most recent identified systematic review (Rijnhout 2021), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting.

Overall, the systematic review and handsearching process identified 16 systematic reviews that included 3 RCTs and 22 nonrandomised cohort studies that met the criteria and were relevant to the research question. The systematic review process also identified one additional nonrandomised cohort study relevant to this research question that was not included in the systematic reviews.

4.4.2 Summary of evidence

4.4.2.1 Systematic reviews evidence

Sixteen systematic reviews (42, 58, 60, 61, 64-75) were identified that assessed the effect of different ratios of RBC to blood component therapy including FFP, PLT and CRYO in patients with critical bleeding. The main characteristics and quality of these systematic reviews and relevant outcomes assessed are summarised in Table 4.15.

McQuilten 2018, Kleinveld 2021 and Richie 2020 restricted their search to RCTs only. All other systematic reviews included RCTs and observational studies as part of their search criteria. Most reviews (da Luz 2019, Rahouma 2017, Cannon 2017, Jones 2016, Poole 2016, Tapia 2013, Kinslow 2020, Maw 2018, Meneses 2020, Rijnhout 2021, Ritchie 2020, Rodriguez 2020, Wirtz 2020) included trauma patients. One review (Rahouma 2017) included trauma and perioperative patients and one review (Phillips 2021) included patients with diagnosed ruptured abdominal aortic aneurysms.

Identified systematic reviews investigated various ratios of blood components. For this review, a high ratio was defined as 1:1:1 and was compared to lower ratios of blood components. Overall, 12 systematic reviews (Kleinveld 2021, Rijnhout 2021, Meneses 2020, Richie 2020, Rodriguez 2020, Wirtz 2020, da Luz 2019, McQuilten 2018, Cannon 2017, Rahouma 2017, Poole 2016, Tapia 2013) identified 3 RCTs (Holcomb 2015, Nascimento 2013, Galganski 2016) relevant to this research question (see <u>'RCT evidence'</u>). A matrix illustrating the overlap of RCTs included in the reviews is provided in Table 4.16.

Twelve systematic reviews (da Luz 2019, Rahouma 2017, Rijnhout 2021, Rodriguez 2020, Meneses 2020, Cannon 2017, Rahouma 2017, Poole 2016, Jones 2016, Tapia 2013, Phillips 2021, Kinslow 2020, Maw 2018) identified 22 nonrandomised cohort studies that met inclusion criteria for this question (see <u>'Observational and cohort studies</u>'). A matrix illustrating the overlap of cohort studies included in the reviews is provided in Table 4.18.

Eight other systematic reviews (McQuilten 2015, Bhangu 2013, Hallet 2013, Johansson 2012, Rajasekhar 2011, Murad 2010, Phan 2010, Zehtabchi 2009, Kozek-Langeneck 2011) were identified in the literature but these reviews did not conduct pooled analyses or provide any additional evidence relevant to this question and are therefore not discussed further in this report. A complete list of excluded studies that met the PICO criteria for this question but were later excluded is provided in **Appendix B** (technical report, volume 2).

Review ID <i>Review quality</i>	Study designPopulationuality(No. of studies)		Intervention	Comparator	Outcomes						
Trauma setting	Trauma setting										
Kleinveld 2021 (64) <i>Moderate</i>	SR / MA of RCTs (5 studies)	Trauma patients	High PLT:RBC ratio	Low PLT:RBC ratio	Mortality MOF TE						
Rijnhout 2021 (66) <i>Moderate</i>	SR / MA of RCTs and observational studies (12 studies)	Military and civilian adult trauma patients	High PLT:RBC ratio	Low PLT:RBC ratio	Mortality Transfusion volumes						
Meneses 2020 (67) <i>Critically low</i>	SR of observational studies (11 studies)	Adult trauma patients	High ratio of blood components	Low ratio of blood components	Mortality						
Ritchie 2020 (68) <i>Low</i>	SR of RCTs (7 studies)	SR of RCTs Trauma patients (7 studies)		gher ratio of Lower ratio of blood blood components components							
Rodriguez 2020 (69) <i>Moderate</i>	020 SR / MA of Adult trauma observational patients studies (33 studies)		High Low FFP:RBC ratio FFP:RBC ratio		Mortality						
Wirtz 2020 (70) <i>Moderate</i>	020 (70) SR of RCTs and observational studies Patients ≥16 years with severe trauma (ISS ≥16) resulting (40 studies)		High FFP:RBC ratio PLT:RBC ratio	High FFP:RBC ratio PLT:RBC ratio	TE						
da Luz 2019 (71) <i>Moderate</i>	Ja Luz 2019 (71) SR / MA of RCTs Civilian and military traum observational studies (54 studies)		High FFP:PLT:RBC ratio	Low FFP:PLT:RBC ratio	Mortality						
McQuilten 2018 SR / MA of RCTs (72) (16 studies) Moderate		Trauma patients	Higher ratio of RRP, PLT, CRYO or FC to RBC	Lower ratio of RRP, PLT, CRYO or FC to RBC	Mortality MOF TE Transfusion volumes						
Cannon 2017 (61) <i>Moderate</i>	n 2017 SR / MA of RCTs Severely injured and adult patients observational requiring studies transfusion and/or (19 studies) ^a ISS >25		High Low FFP:RBC ratio FFP:RBC ratio PLT:RBC ratio PLT:RBC ratio		Mortality Transfusion volume						
Rahouma 2017 (73) <i>Critically low</i>	SR / MA of RCTs and observational studies (36 studies)	Trauma and surgical patients requiring transfusion	High FFP:RBC ratio	Low FFP:RBC ratio	Mortality						
Jones 2016 (74) Critically low	SR of observational studies (25 studies) ^b	Civilian or military trauma patients	High FFP:RBC ratio PLT:RBC ratio	Different FFP:RBC ratio PLT:RBC ratio	Mortality MOF						

Table 4.15 Characteristics and quality of systematic review evidence: ratio of RBC to blood component therapy

Review ID <i>Review quality</i>	Study design (No. of studies)	Population	Intervention	Comparator	Outcomes		
Poole 2016 (42) Critically Low	SR of RCTs and observational studies (10 studies)	Adult trauma patients (excluding TBI) in non-military setting	High ratio FFP:RBC FFP:PLT:RBC	Lower ratio FFP:RBC FFP:PLT:RBC	Mortality		
Tapia 2013 (75) <i>Critically low</i>	SR / MA of RCTs, Coh & IV studies (20 studies)	Military and civilian trauma patients with at least 30% penetrating injury receiving MT	High FFP:RBC ratio PLT:RBC ratio	Low FFP:RBC ratio PLT:RBC ratio	Mortality		
Paediatric, trau	ma setting						
Kinslow 2020 (58) <i>Critically low</i>	SR of observational studies (33 studies)	Paediatric trauma patients	Higher ratio of RBC to blood components (PLT or FFP)	Lower ratio of RBC to blood components (PLT or FFP)	Mortality Adverse outcomes		
Maw 2018 (60) Critically low	SR of observational studies (4 studies)	Paediatric patients (<18 years) with traumatic injury requiring blood transfusion	MHP High ratios: FFP:RBC Plasma/PLT:RBC	MHP Low ratios: FFP:RBC Plasma/PLT:RBC	Mortality ICU LOS Unnecessary transfusion		
Surgical setting							
Phillips 2021 (65) <i>Low</i>	SR of observational studies (7 studies)	Surgical patients diagnosed with AAA	High FFP:RBC ratio	Low FFP:RBC ratio	Mortality		

AAA, abdominal aortic aneurism; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ICU, intensive care unit; ISS, injury severity score; LOS, length of stay; MHP, major haemorrhage protocol; MOF, multiple organ failure; MT, massive transfusion; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; SR, systematic review; TBI, traumatic brain injury; TE, thromboembolic event

a. 15 studies (N=5292) assessed RBC:FFP and 4 studies (N=1607) assessed RBC:PLT.

b. 17 studies (N=6801) assessed RBC:FFP and 8 studies (N=3960) assessed RBC:PLT.

4.4.2.2 Randomised controlled trials

A matrix illustrating the RCTs identified in the included systematic reviews is provided in Table 4.16. Among the 12 RCTs identified by the included systematic reviews, 3 were considered relevant to this review because they examine the effect of high (1:1:1) ratios in patients with critical bleeding.

One review (Kleinveld 2021) identified 3 RCTs that did not meet inclusion criteria for this research question and were not included. One study (Sperry 2018) assessed prehospital plasma resuscitation compared to standard care resuscitation. Two studies (Baksaas-Aasen 2020, Gonzalez 2016) investigated the addition of viscoelastic haemostatic assays to MHPs on patient outcomes and are considered in Section 4.9 (question 8).

One other review (McQuilten 2018) identified 13 RCTs, of which 10 were ongoing trials and did not have any data available for inclusion in the analysis. Three RCTs did not meet the inclusion criteria for this question. One RCT (Cotton 2013) compared treatment with whole blood to treatment with blood components in adult trauma patients, the other 2 studies (Curry 2015, Nascimento 2016) compared the use of CRYO or fibrinogen concentrate (FC) versus no CYRO or FC in trauma patients. Both Curry 2015 and Nascimento 2016 are considered in Section 4.7 (question 6).

	Trauma												
	Study ID	Holcomb 2015	Nascimento 2013	Galganski 2016	Gonzalez 2016	Sperry 2018	Baksaas-Aasen 2020	Curry 2015	Cotton 2013	Nascimento 2016	Rahbar 2015	Curry 2018	Akbari 2018
	Kleinveld 2021	✓	~		Х	X	Х						
	Rijnhout 2021	~	~										
	Meneses 2020	✓											
	Richie 2020	~	~					Х	Х		Х	Х	Х
۵	Rodriguez 2020	~	~										
Ň	Wirtz 2020	~											
svie	da Luz 2019	~	~										
ž	McQuilten 2018	~	~	~				Х	Х	Х			
	Cannon 2017	~											
	Rahouma 2017	~	~										
	Poole 2016	~											
	Tapia 2013	✓											

Table 4.16 Overlap table of RCTs identified by included systematic reviews: RBC:FFP and RBC:PLT ratios

FFP, fresh frozen plasma; PLT, platelet; RBC, red blood cell; RCT, randomised controlled trial

✓ = study included in this review;

X = study did not meet the inclusion criteria for this review;

-- = study identified by the systematic review authors but not included (no usable data)

No additional RCTs were identified in the systematic review and handsearching process. The main characteristics and quality of eligible RCTs and the relevant outcomes assessed are detailed in Table 4.17.

Two RCTs (Holcomb 2015, Nascimento 2013) compared the effect of high (1:1:1) RBC:FFP:PLT transfusion ratios to lower transfusion ratios on the 28-day mortality in trauma patients (aged 15 years or older) requiring massive transfusion. One RCT (Galganski 2016) compared the effect of a high (1:1) RBC:FFP ratio versus low (1:4) ratio in 45 children (aged younger than 18 years) who had a third degree burn injury of greater than 20% of the total body surface area. The study reported no deaths at hospital discharge and is not included in the meta-analysis for this research question. Transfusion requirements reported did not meet criteria for this question and are not reported further.

All 3 included RCTs were carried out in the US; 2 in trauma centres and one in a children's hospital. Overall, systematic review authors judged included RCTs to be high risk of bias with blinding being the main sources of bias. Holcomb 2015 was the only RCT that attempted to minimise bias from lack of blinding by having each death adjudicated by a clinician blinded to group assignment (McQuilten 2018).
Study ID <i>Risk of bias</i>	Study design	Population N	Intervention	Comparator	Outcomes
Trauma settin	g				
Holcomb 2015 (PROPPR trial) <i>Moderate</i>	RCT, multicentre	Severely injured patients predicted to require MT N=680	FFP:PLT:RBC 1:1:1	FFP:PLT:RBC 1:1:2	Mortality Morbidity
Nascimento 2013 <i>High</i>	RCT, SC, unblinded	Patients with hypotension and bleeding expected to need MT N=78	Fixed FFP:PLT:RBC ratio 1:1:1	Transfusion guided by the institution's usual MT protocol	Mortality Transfusion volume
Paediatric, tra	uma setting				
Galganski 2016 <i>High</i>	RCT, SC	Paediatric patients (<18 years) admitted to ICU *with a third degree burn injury of ≥ 20% TBSA N=45	FFP:RBC 1:1	FFP:RBC 1:4	Mortality Transfusion volume

Table 4.17	Characteristics and quality of RCT evidence: ratio of RBC to blood component
	therapy

FFP, fresh frozen plasma; ICU, intensive care unit; MC, multicentre; MT, massive transfusion; PLT, platelet; RBC, red blood cells; RCT, randomised controlled trial; SC, single centre; TBSA, total body surface area

4.4.2.3 Observational and cohort studies

Twenty-two observational studies identified by the included systematic reviews met the inclusion criteria and were considered relevant to this question. A matrix illustrating the overlap of nonrandomised cohort studies identified in the included systematic reviews that met our inclusion criterion is provided in Table 4.18. The main characteristics of these studies is provided in Table 4.20.

Ten reviews (da Luz 2019, Rahouma 2017, Rijnhout 2021, Rodriguez 2020, Meneses 2020, Wirtz 2020, Cannon 2017, Poole 2016, Jones 2016, Tapia 2013) identified 47 cohort studies reporting outcome data for mortality, morbidity and transfusion volume but the studies assessed ratios of blood components that did not meet the high ratio criteria of 1:1:1 and were therefore not included in this review. An overlap table listing the observational and cohort studies that did not meet the high ratio criterion is provided in Table 4.19.

One other review (Tanaka 2017) searched for evidence regarding the ratio of RBC to FFP used in patients with major obstetric bleeding. The authors identified 5 retrospective noncontrolled observational studies that reported RBC:FFP ratios use in the obstetric setting. As these were noncomparative case series, they were not eligible for inclusion and were not considered further.

One additional cohort study (Peralta 2016) (76) was identified in the literature search that investigated the effect of high PLT:RBC ratio defined as 1:1.5 or more within 4 hours postinjury and their impact on the outcomes of trauma patient receiving MHP. As this ratio does not meet the set criteria of 1:1:1 for this question, this study was not considered further.

		Traur	Trauma											Surgi	cal						Paedi	atrics,	traum	a	
	Study ID	Balvers 2017	Duchesne 2008	Duchesne 2009	Haltmeier 2017	Holcomb 2011	Maegele 2008	Perkins 2009	Sambasivan 2011	Vulliamy 2017	Wafaisade 2011	Zink 2009	Holcomb 2013	Mell 2010	Mazzeffi 2016	Johansson 2007	Johansson 2008	Henriksson 2012	Hall 2013	Tadlock 2010	Butler 2019	Cunningham 2019	Edwards xxx	Noland 2018	Nosanov 2013
	da Luz 2019	✓	~	~	×	~	×	~	~	~	~	~													
	Rijnhout 2021	~				~																			
	Rodriguez 2020		~				~				~	~													
	Meneses 2020		~																						
۵	Cannon 2017			~				~																	
Ň	Rahouma 2017		~	~			~				~	~		~	1										
svie	Poole 2016					~			~		~		~												
å	Jones 2016			~		~	~	~				~													
	Tapia 2013		1						~			~													
	Phillips 2021													~	~	~	~	~	~	~					
	Kinslow 2020																				~	~		~	~
	Maw 2018																								

Table 4.18 Overlap table of observational and cohort primary studies identified by included systematic reviews that meet the 1:1:1 ratio inclusion criterion: RBC:FFP and RBC:PLT ratios

FFP, fresh frozen plasma; PLT, platelet; RBC, red blood cell

✓ = study included in this review

X = study did not meet the inclusion criteria for this review

-- = study identified by the systematic review authors but not included (no usable data)

		Tra	aun	na																																												
	Study ID	gman 2007	aman 201	vn 2011	vn 2012	2016	2012	on 2008	3iasi 2011	te 2009	thesne 2010	hesne 2013	o 2018	dry 2013	ter 2008	nin 2013	din 2014	comb 2008 ^ª	comb 2012 ^b	a 2010	huk 2008	2014	o 2014	cher 2014	enberger 2011	jnotti 2011	a 2010	di 2015	1ger 2011	alta 2015	kins 2011	coke 2012	/ell 2011	ea 2008	rpe 2012	z 2010	ms 2014	mons 2011	rry 2008	iella 2011	erke 2011	worth 2016	der 2009	eira 2009	urraga 2015	2010	g 2015	nski 2013
		Bor	Bor	Bro	Bro	Bui	Cap	Cott	de E	Den	Duc	Duc	End	Guio	Gun	Halr	Har	Ĕ	Ĕ	Inat	Kas	Kim	Kud	Kut	Lust	Maç	Mitr	Nar	Peir	Pera	Per	Pido	Roy	Scal	Sha	Sha	Sim	Sim	Spe	Spir	Spo	Star	Syn	Teix	Dnd	Van	Yan	Ziel
	da Luz 2019	X	X			X											X	X				X	X		X	X		X	Х				X		X				X			X	X	X				
	Rahouma 2017	x	x	x	x				x	x					x			x			x	x			x	x	x		х				x		x	x			x	х	x	x	x	x	x	x	x	
	Rijnhout 2021				Х		Х													Х					Х						Х		Х			X	Х											
₽	Rodriguez 2020		x		x				x	x		х	х	x	x			x			x	x	x		x	x	x		х	x			x		x	x	x		x	х	x	х	х	x				
eview	Meneses 2020	x								x								x			x													x		x			x					x				
ž	Wirtz 2020													Х																																	2	X
	Cannon 2017	X														Х		Х				Х		Х		Х	Х		Х							X			Х				Х	Х				
	Poole 2016		X																	Х							Х							Х										х				
	Jones 2016	Х			X		X											Х					Х				Х		Х										Х				Х			Х		
	Tapia 2013	Х		X			X	Х	Х	Х	Х				Х			Х	Х	Х												Х	Х		Х	Х		Х										

Table 4.19 Overlap table of observational and cohort primary studies identified by included systematic reviews that do not meet the 1:1:1 ratio inclusion criterion: RBC:FFP and RBC:PLT ratios

FFP, fresh frozen plasma; PLT, platelet; RBC, red blood cell

study included in this review; X = study did not meet the inclusion criteria for this review; -- = study identified by the systematic review authors but not included (no usable data)

^a Superseded by Holcomb 2009

^b Superseded by Holcomb 2013

Review ID Risk of bias	Study design	Population N	Intervention	Comparator	Outcomes
Trauma sett	ing			1	1
Balvers 2017 High	Prospective, MC	Trauma patients receiving 4 or more units of RBC in 24 hours N=385	High ≥1:1 ratio Plasma:RBC	High <1:1 ratio Plasma:RBC	Transfusion volume Waste of blood components
Haltmeier 2017 <i>High</i>	Retrospective	Patients with isolated severe blunt TBI N=242	High ratio Plasma/PLT:RBC	Low ratio Plasma/PLT:RBC	Mortality
Vulliamy 2017 <i>High</i>	Prospective	Patients who received at least 4 units RBC N=161	High ratio FFP:RBC	Low ratio FFP:RBC	Mortality
Holcomb 2011 <i>High</i>	Retrospective, MC	Adult trauma patients who had received at least one unit of RBC in the ED N=427	High 1:1 ratio PLT:RBC	Low >1:20 ratio PLT:RBC	Mortality Transfusion volume
Sambasivan 2011 <i>High</i>	Retrospective, MC	Trauma patients N=1181	High 1:1 ratio FFP/PLT:RBC	Low <1:1 ratio FFP/PLT:RBC	Mortality
Wafaisade 2011 <i>High</i>	Retrospective	Trauma patients 16 years or older with an ISS \geq 16 who had received multiple transfusion but not MT N=970	High >1:1 ratio FFP:RBC	Low <1:1 ratio FFP:RBC	Mortality Morbidity Transfusion volume
Duchesne 2009 High	Retrospective	Trauma patients with initial ED diagnosis of TIC who required transfusion during initial surgical intervention N=89	High 1:1 ratio FFP:RBC	Low 1:4 ratio FFP:RBC	Mortality
Perkins 2009 High	Retrospective	Trauma patients receiving MT (10 or more units of blood within 24 hours) N=310	High PLT:RBC	Low PLT:RBC	Mortality
Zink 2009 <i>Moderate</i>	Retrospective	Trauma patients requiring massive transfusion N=153	High ≥1:1: ratio FFP:RBC	Low <1:4: ratio FFP:RBC	Mortality Transfusion volume

Table 4.20 Characteristics and quality of observational and cohort evidence: RBC:FFP, RBC:PLT or RBC:CRYO ratio

Review ID <i>Risk of bias</i>	Study design	Population N	Intervention	Comparator	Outcomes
Duchesne 2008 <i>High</i>	Retrospective	Adult trauma patients (blunt and penetrating) requiring surgical intervention and > 10 units of RBC transfusion N=135	High 1:1 ratio FFP:RBC	Low 1:4 ratio FFP:RBC	Mortality
Maegele 2008 <i>High</i>	Retrospective	Trauma patients with an ISS >16 and who received ≥10 RBC N=713	High 1:1 ratio FFP:RBC	Low <1:1 ratio FFP:RBC	Mortality Morbidity Transfusion volume
Paediatric, t	rauma setting				
Cunningha m 2019 <i>High</i>	Retrospective	Paediatric trauma patients (≤18 years) requiring MT N=465	High ratio Plasma:RBC	Low ratio Plasma:RBC	Mortality Transfusion volume
Butler 2019 <i>High</i>	Retrospective	Paediatric trauma patients (≤14 years) receiving MT N=583	High ratio FFP/PLT:RBC	Low ratio FFP/PLT:RBC	Mortality
Noland 2018 <i>High</i>	Retrospective	Paediatric trauma patients (≤18 years) requiring MT N=110	High ratio FFP:RBC	Low ratio FFP:RBC	Mortality
Nosanov 2013 High	Retrospective	Paediatric trauma patients (≤18 years) requiring MT (transfusion of ≥50% total blood volume) N=105	High ratio Plasma/PLT:RBC	Low ratio Plasma/PLT:RBC	Mortality
Surgical set	ting				
Hall 2013 Serious	Pre-post *2005 to 2008, 2010	Patients with rAAA (nonconsecutive) N=89	High ratio RBC:FFP (4:4:1)	Low ratio FFP:RBC (>1)	Mortality *not clear if in-hospital or 30-day
Henriksson 2012 <i>Serious</i>	Retrospective *1992 to 1999, 2000 to 2008	Patients with rAAA (consecutive) N=174 *trauma registry	High ratio RBC:FFP (1:1)	Low ratio RBC:FFP (>1:1)	Mortality, 30-day
Kauvar 2011 <i>Serious</i>	Retrospective *1992 to 2008 (16-yr period)	Patients with rAAA who received >10U of blood or blood products N=87	High ratio RBC:FFP (≤2)	Low ratio RBC:FFP (>2)	Mortality, in hospital
Mell 2010 <i>Serious</i>	Retrospective *1987 to 2007 (20-yr period)	Patients with rAAA who received >10U of blood or blood products N=128	High ratio RBC:FFP (≤2)	Low ratio RBC:FFP (>2)	Mortality, 30-day
Tadlock 2010 <i>Serious</i>	Retrospective *2002 to 2008	Patients with rAAA (consecutive) N=12	High ratio RBC:FFP (1:1)	Low ratio RBC:FFP (>1:1)	Mortality *not clear if in-hospital or 30-day

Review ID	Study design	Population	Intervention	Comparator	Outcomes
Risk of bias		N			
Johansson	Pre-post I	Patients with rAAA	High ratio	Low ratio	Mortality, 30-day
2008	*2002 to 2004,	(non-consecutive)	FFP:RBC:PLT	RBC:FFP (>1)	
Serious	2006 to 2007	N=146	(5:5:2)		
Johansson	Pre-post	Patients with rAAA	High ratio	Low ratio	Mortality, 30-day
2007	* 2002-2005	(consecutive)	RBC:FFP:PLT	RBC:FFP (>1)	
Serious		N=132	(5:5:2)		

ED, emergency department; FFP, fresh frozen plasma; ISS, injury severity score; MC, multicentre; MT, massive transfusion; PLT, platelet; rAAA, ruptured abdominal aortic aneurysm; RBC, red blood cells; TBI, traumatic brain injury; TIC, trauma-induced coagulopathy

Trauma setting

There were 11 observational studies conducted in adult trauma patients identified by 9 systematic reviews (42, 61, 66, 67, 69, 71, 73-75) that met the PICO criteria for this question (inclusive of a high 1:1:1 ratio).

Five studies (Vulliamy 2017, Wafaisade 2011, Duchesne 2009, Duchesne 2008, Maegele 2008) assessed RBC:FFP ratios, 2 studies (Holcomb 2011, Perkins 2009) assessed RBC:PLT ratios and 4 studies (Hatimeier 2017, Balvers 2017, Sambasivan 2011, Zink 2009) assessed both RBC:FFP and RBC:PLT ratios.

The cohort studies were carried out in trauma centres in Denmark, Germany, Iraq, the Netherlands, UK or the US. Overall, risk of bias of included studies was judged to be moderate with concerns arising due to confounding.

Surgical setting

There were 7 cohort studies (Hall 2013, Henriksson 2012, Kauvar 2011, Mell 2010, Tadlock 2010, Johansson 2008, Johansson 2007) identified by one systematic review (65) that evaluated whether a higher RBC:FFP ratio improves patient outcomes in the surgical setting. All studies included patients with ruptured abdominal aortic aneurysms.

Five studies (Hall 2013, Henriksson 2012, Johansson 2007, Johansson 2008, Tadlock 2010) defined a 1:1 ratio of RBC:FFP as high and 2 studies (Kauvar 2011, Mell 2010) did not define a high ratio.

Six studies were carried out in single-centre surgical settings in North America and one study involved patients in a trauma registry in Denmark. Overall, review authors judged included studies to be at serious risk of bias, with a significant amount of bias arising from confounding and patient selection related to the retrospective study design or the use of historical control groups.

Paediatric setting

Six cohort studies (total 1025 patients) were identified by one systematic review (Kinslow 2020 (58) that assessed blood component ratios in paediatric patients undergoing massive transfusion. Four studies (Butler 2019, Cunningham 2019, Noland 2018, Nosanov 2013) reported data relevant to our review.

All 4 studies included paediatric trauma patients with predominantly blunt injuries and compared a high ratio of blood components (1:1) to lower ratios. Three studies (Butler 2019, Noland 2018, Nosanov 2013) assessed RBC:FFP and one study (Cunningham 2019) assessed RBC:Plasma.

Included cohort studies were carried out in paediatric trauma centres in the US. Review authors did not conduct a risk of bias assessment of included studies but noted there was significant heterogeneity throughout due to adherence to ratio targets and differences in activation of major haemorrhage protocols. For these reasons, authors were unable to conduct a formal meta-analysis.

4.4.3 Results

4.4.3.1 High transfusion (1:1:1) ratio vs lower transfusion ratio

Mortality

A summary of the evidence reported in the identified systematic reviews relating to mortality in patients with critical bleeding is presented in Table 4.21.

The identified systematic reviews suggest there is a significant survival benefit for patients who receive a high (1:1:1) blood component to RBC ratio compared with those who receive a low (2:1:1) blood component to RBC ratio, regardless of clinical setting. However, the evidence is largely based on observational studies that are heterogeneous and at risk of bias. The RCT evidence suggests no difference in mortality between a 1:1:1 or a 2:1:1 transfusion strategy.

A meta-analysis of data from RCTs included in this review (see Figure 4.12) showed the mortality rate (latest timepoint) in the trauma setting to be comparable among those who received high transfusion ratios of blood components compared to those who received lower transfusion ratios with the relative risk (RR) of 1.26 observed (95% CI 0.49, 3.22; p = 0.64; random effect, I² = 75%). Neither of the included RCTs were powered to detect differences in mortality.

In contrast, a meta-analysis of data from nonrandomised cohort studies included in this review showed a significant difference in mortality rate of 24% (474/1978) observed among patients who received a high transfusion ratio of blood components compared to 33.1% (1219/3686) among patients who did not (RR 0.60; 95% CI 0.48, 0.74; p < 0.00001, random effect, $I^2 = 79\%$).

Among patients with blunt and penetrating trauma, a total of 308 patients received a high transfusion ratio (1:1:1) of blood components compared with 922 patients who received lower ratios. A significant difference in mortality was observed between groups (24.3% vs 31.4%, RR 0.58; 95% CI 0.41, 0.82; p = 0.002, random effect, $l^2 = 88\%$).

Among patients with ruptured abdominal aortic aneurysms, the observed mortality rate of 23.6% (88/373) among patients receiving a high transfusion ratio was significantly different to the mortality rate of 46.4% (143/308) among patients receiving lower transfusion ratios (RR of 0.56; 95% CI 0.43, 0.72; p < 0.0001; random effect, $l^2 = 15\%$).

Among paediatric trauma patients, a total of 78 patients received a high transfusion ratio of blood components compared to 154 patients who received lower transfusion ratios, with no significant difference in mortality observed (23.3% vs 34.6%, RR 0.70; 95% CI 0.47, 1.04; random effect, $I^2 = 40\%$).

Figure 4.12 Forest plot of comparison: high ratio vs low ratio blood components, outcome: Mortality, latest timepoint

	High ratio ((1:1:1)	Low ratio ((2:1:1)		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl
1.1.1 Trauma (RCTs)							
Holcomb 2015 (RCT)	75	338	89	342	5.9%	0.85 [0.65, 1.11]	
Nascimento 2013 (RCT)	13	40	5	35	2.9%	2.27 [0.90, 5.74]	
Subtotal (95% CI)		378		377	8.8%	1.26 [0.49, 3.22]	\bullet
Total events	88		94				
Heterogeneity: Tau ² = 0.36	; Chi ² = 3.99,	df = 1 (P	P = 0.05); l ² =	= 75%			
l est for overall effect: $Z = 0$	0.47 (P = 0.64	+)					
1.1.2 Trauma (observation	nal)						
Duchesne 2008	, 18	71	56	64	5.2%	0.29 [0.19, 0.44]	
Duchesne 2009	13	46	22	43	4.5%	0.55 [0.32, 0.95]	
Haltmeier 2017	53	156	46	86	5.8%	0.64 [0.47, 0.85]	
Holcomb 2011	65	216	101	211	6.0%	0.63 [0.49, 0.81]	
Maegele 2008	28	115	220	484	5.6%	0.54 [0.38, 0.75]	
Perkins 2009	15	96	86	150	4.8%	0.27 [0.17, 0.44]	
Sambasivan 2011	47	202	126	979	5.8%	1.81 [1.34, 2.44]	
Vulliamy 2017	25	107	15	54	4.5%	0.84 [0.49, 1.46]	
Wafaisade 2011	31	210	194	760	5.5%	0.58 [0.41, 0.82]	
Zink 2009	13	51	56	102	4.8%	0.46 [0.28, 0.77]	
Subtotal (95% CI)		1270		2933	52.6%	0.58 [0.41, 0.82]	\bullet
Total events	308		922				
Heterogeneity: Tau ² = 0.26	; Chi² = 77.28	3, df = 9 (P < 0.00001); l ² = 88	%		
Test for overall effect: Z = 3	3.11 (P = 0.00	02)					
1.1.3 Surgical (observation	onal)						
Hall 2013	21	68	6	21	3.5%	1.08 [0.50, 2.32]	
Henriksson 2012	20	100	23	74	4.7%	0.64 [0.38, 1.08]	
Johansson 2007	17	50	46	82	5.1%	0.61 [0.39, 0.93]	
Johansson2008	16	64	46	82	4.9%	0.45 [0.28, 0.71]	
Mell 2010	13	87	16	41	4.1%	0.38 [0.20, 0.72]	
l adlock 2010 Subtotal (95% CI)	1	373	6	308	1.2% 23.5%	0.33 [0.06, 1.91]	
Total overts	88	5/5	1/3	500	20.070	0.50 [0.45, 0.72]	•
Heterogeneity: $T_{2}u^2 = 0.02$	00 • Chi2 – 5 91	df = 5 (P	143 - 0 32)· 12 -	- 15%			
Test for overall effect: $7 - 7$, 0111 = 0.01, 1 37 (P < 0.00	01 – 5 (i 001)	- 0.52), 1 -	- 1370			
1.1.4 Paediatrics (observa	ational)						
Butler 2019	46	136	104	232	5.9%	0.75 [0.57, 0.99]	
Cunningham 2019	15	126	38	163	4.5%	0.51 [0.29, 0.89]	
Noland 2018	6	39	10	35	3.0%	0.54 [0.22, 1.33]	
Nosanov 2013	11	34	2	15	1.7%	2.43 [0.61, 9.63]	
Subtotal (95% CI)		335		445	15.1%	0.70 [0.47, 1.04]	•
Total events	78		154				
Heterogeneity: Tau ² = 0.06	; Chi ² = 4.98,	df = 3 (P	' = 0.17); l ² =	= 40%			
i est for overall effect: Z = 1	I.77 (P = 0.08	3)					
Total (95% CI)		2356		4063	100.0%	0.63 [0.51, 0.78]	•
Total events	562		1313				
Heterogeneity: Tau ² = 0.17	; Chi ² = 101.1	I3, df = 2	1 (P < 0.000	001); l ² = ⁻	79%		
Test for overall effect: Z = 4	4.26 (P < 0.00	001)					U.U1 U.1 1 10 100
Test for subgroup difference	es: Chi ² = 3.2	24. df = 3	(P = 0.36), I	l² = 7.4%			

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			High ratio n/N (%)	Low ratio n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting	7	1	1	1	1	1	1		
da Luz 2019 SR <i>Moderate</i> <i>quality</i>	N = 2414 (5 Coh)	Adult trauma patients	Trauma centres (UK, Denmark, Netherlands, Germany, Iraq)	FFP:RBC High ratio (1:1) vs Iower ratio (<1:1)	Mortality, 24 hours Balvers 2017 Maegele 2008 Perkins 2009 Vulliamy 2017 Wafaisade 2011	126/738 (17.1) 89/210 (42.4) 13/115 (11.3) 5/96 (5.2) 8/107 (7.5) 11/210 (5.2)	420/1676 (25.1) 65/169 (38.5) 158/484 (32.6) 75/209 (35.9) 9/54 (16.7) 113/760 (14.9)	OR 0.34 (0.14, 0.82) 1.18 (0.78, 1.78) 0.26 (0.14, 0.48) 0.10 (0.04, 0.25) 0.40 (0.15, 1.12) 0.32 (0.17, 0.60)	Favours high ratio p = 0.02 Substantial heterogeneity $I^2 = 88\% (p < 0.00001)$
	N = 749 (2 RCTs)	Adult trauma patients	Trauma centres (US, Canada)	FFP:PLT:RBC High (1:1:1) vs lower ratio (1:1:2)	Mortality, 28-30 days Holcomb 2015 Nascimento 2013	88/378 (23.3) 75/338 (22.2) 13/40 (32.5)	94/377 (25) 89/342 (26) 5/35 (14.3)	OR 1.35 (0.40, 4.59) 0.81 (0.57, 1.15) 2.89 (0.91, 9.17)	No significant difference p = 0.63 Substantial heterogeneity $l^2 = 76\% (p = 0.04)$
	N = 4203 (10 Coh)	Adult trauma patients	Trauma centres (UK, Germany, US, Iraq)	FFP:RBC High ratio (1:1) vs Iower ratio (<1:1)	Mortality, 30 days Duchesne 2008 Duchesne 2009 Haltmeier 2017 Holcomb 2011 Maegele 2008 Perkins 2009 Sambasivan 2011 Vulliamy 2017 Wafaisade 2011 Zink 2009	308/1270 (24.3) 18/71 (23.4) 13/46 (28.3) 53/156 (34) 65/216 (30.1) 28/115 (24.3) 15/96 (15.6) 47/202 (23.3) 25/107 (23.4) 31/210 (14.8) 13/51 (25.5)	922/2933 (31.4) 56/64 (87.5) 22/43 (51.2) 46/86 (53.5) 101/211 (47.9) 220/484 (45.5) 86/150 (57.3) 126/979 (12.9) 15/54 (27.8) 194/760 (25.5) 56/102 (54.9)	OR 0.38 (0.22, 0.68) 0.05 (0.02, 0.12) 0.38 (0.16, 0.90) 0.45 (0.26, 0.77) 0.47 (0.32, 0.70) 0.39 (0.24, 0.61) 0.14 (0.07, 0.26) 2.05 (1.41, 299) 0.79 (0.38, 1.67) 0.51 (0.33, 0.76) 0.28 (0.13, 0.59)	Favours high ratio p = 0.001 Substantial heterogeneity l ² = 91% (p < 0.0001)
Kleinveld 2021 SR <i>Moderate</i> <i>quality</i>	N = 1757 (2 RCTs)	Trauma patients (≥16 years)	Trauma centres (US, Canada)	PLT:RBC High ratio vs lower ratio	Mortality, 24-hours Nascimento 2013 Holcomb 2015	8/37 (21.6) 43/338 (12.7)	3/32 (9.4) 58/342 (17.0)	OR 2.67 (0.64, 11.07) OR 0.71 (0.47, 1.09)	NR
Cannon 2017 SR <i>Moderate</i> <i>quality</i>	N = 927 (3 Coh)	Adult trauma patients	Trauma centres (UK, Denmark, Netherlands, Germany, Iraq)	PLT:RBC High ratio (1:1) vs Iower ratio (<1:1)	Mortality, overall Holcomb 2008 Perkins 2008 Shaz 2010	163/505 (32.3) 67/234 (28.6) 49/145 (33.8) 47/126 (37.3)	239/764 (31.3) 94/184 (51.1) 86/150 (57.3) 59/88 (67)	OR 0.36 (0.27, 0.47) 0.38 (0.26, 0.58) 0.38 (0.24, 0.61) 0.29 (0.16, 0.52)	Favours high ratio p < 0.00001 No heterogeneity $l^2 = 0\% (p = 72\%)$

Table 4.21 Results for high ratio of blood components versus low ratio of blood components: Patients with critical bleeding - Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			High ratio n/N (%)	Low ratio n/N (%)	Risk estimate (95% CI)	Statistical significance p-value
									Heterogeneity ^b
Paediatrics, tra	auma setting								
Kinslow 2020	N = 804	Paediatric trauma	Paediatric trauma	FFP:RBC or PLT:RBC	Mortality, overall ^d	NR	NR	NR	NR
SR	(4 Coh)	patients with	(US, Iraq,	High vs lower ratios	Noland 2018	6/39 (15)	10/35 (29)		
Critically low		various ISS	Afghanistan)		Cunningham 2019	15/126 (12)	38/163 (23)		
quality					Butler 2019 Nosanov 2013	46/136 (33.8)	104/232 (44.8)		
Surgical setting	a		1	1	140501104 2013	1004 (02.0)	2/10 (10.0)		I
Dhilling 2021	N C01	Adults with a	Surgical SC	EED DRC	Mortality 20 days or	ND	ND	ND	ND
F111111p3 2021	(C C c b)		(North Amorica	Lligh up lower ration	latost timopoint	INK	INK	INIX	NR
5K	(6 CON)		(North America,	Fight vs lower ratios	Mail 2010	10/07 (15)	10/41/00)		- 0.00
Low quality			Deninark)		Johansson 2007	13/87 (15)	16/41 (39)	NR	p<0.03
					Johansson 2008	16/64 (25)	46/82 (56)	NR	p = 0.02
					Henriksson 2012	20/100 (20)	23/74 (31)	NR	p = 0.111
					Hall 2013	21/68 (31)	6/21 (28)	NR	<i>p</i> > 0.05
					Tadlock 2010	1/4 (25)	6/8 (75)	NR	<i>p</i> = 0.222
	N = 87				Mortality, in-hospital	19/39 (49)	19/48 (40)	NR	<i>p</i> = 0.39
	(1 Coh)								
	Kauvar 2011								

AAA, abdominal aortic aneurysm; CI, confidence interval; Coh, cohort; FFP, fresh frozen plasma; ISS, injury severity score; NR, not reported; OR, odds ratio; PLT, platelet; RBC, red blood cell; RCT, randomised controlled trial; SC, single centre; SR, systematic review; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. Mortality data for identified by Kinslow 2020 sourced from primary study. No meta-analysis conducted by Kinslow 2020.

Morbidity

A summary of the evidence reported in the identified systematic reviews relating to morbidity (e.g. thromboembolic events, multiple organ failure [MOF], acute respiratory distress syndrome [ARDS]) in patients with critical bleeding is presented in Table 4.22 and Table 4.23.

Thromboembolic events

One RCT (Holcomb 2015) in the trauma setting and one cohort study (Butler 2019) in the paediatric setting reported on the outcome of thromboembolic events. Combined data (see Figure 4.13) suggest no important difference between groups (RR 1.46; 95% CI 0.64, 3.32; p = 0.37; random effect; I²=59%).

Holcomb 2015 suggested no significant difference in thromboembolic events (deep vein thrombosis, pulmonary embolus) between patients who received high ratio of blood components (39/338, 11.5%) compared with those who did not (37/342, 10.8%) (RR 1.07; 95% Cl 0.64, 3.32; p = 0.37; random effect; l²=59%).

Butler 2019 suggested a nonsignificant increased risk in thromboembolic events (deep vein thrombosis) between paediatric patients who received a high ratio of blood components (9/136, 6.6%) compared to paediatric patients who received a low ratio (6/323, 2.6%) (RR 2.56; 95% CI 0.93, 7.03; p = 0.07).

	Experim	ental	Contr	ol		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	I M-H, Random, 95% CI	
1.2.1 Trauma								
Holcomb 2015 (RCT)	39	338	37	342	64.3%	1.07 [0.70, 1.63]	-#-	
Subtotal (95% CI)		338		342	64.3%	1.07 [0.70, 1.63]	•	
Total events	39		37					
Heterogeneity: Not app	licable							
Test for overall effect: Z	2 = 0.30 (P	= 0.77)						
1.2.2 Paediatrics								
Butler 2019 (1)	9	136	6	232	35.7%	2.56 [0.93, 7.03]		
Subtotal (95% CI)		136		232	35.7%	2.56 [0.93, 7.03]		
Total events	9		6					
Heterogeneity: Not app	licable							
Test for overall effect: Z	. = 1.82 (P	= 0.07)						
Total (95% CI)		474		574	100 0%	1 46 [0 64 3 32]		
Total (00% CI)	40	717	13	014	100.070	1.40 [0.04, 0.02]		
Hotorogonoity: Tau ² – C	40 1 22: Chi2 -	2 /5 di	+	0 12)+	2 - 50%			-
Tost for overall offect: 7	- 0 00 /P	2.40, ui - 0 37)	- 1 (1 -	0.12), 1	- 5570		0.01 0.1 1 10 10)0
Test for subgroup diffor	. = 0.90 (F oncos: Chii	= 0.37	df _ 1 (D	_ 0 12	0/_	Favours high (1:1) ratio Favours lower ratio		
		- 2.4J,	ui – i (r	- 0.12), 1- – 39.1	/0		
$\frac{FUUIIIUIUS}{(4) I i ch rotio} (1, 4, 4) vol$	au (1 0)							
(1) mign ratio (>1.1) VS	IOW (<1.2)							

Figure 4.13 Forest plot of comparison: high ratio vs low ratio blood components, outcome: Morbidity, thromboembolic events

Multiple organ failure

Two RCTs reported on the outcome of MOF in the trauma setting. A meta-analysis of the 2 RCTs (Figure 4.14) found no significant difference in MOF between patients who received a high ratio of blood components (21/375, 5.6%) compared with patients who received a low ratio (15/374, 4%) (RR 1.39, 95% CI 0.73, 2.63; p = 0.32; random effects; $I^2 = 0$ %).

Figure 4.14 Forest plot of comparison: high ratio vs low ratio blood components, outcome: Morbidity, multiple organ failure



Other adverse events

Two RCTs and 6 cohort studies were found in the trauma setting that reported on the outcome of other adverse events.

For the outcome of ARDS, one RCT (Holcomb 2015) reported no significant difference between patients who received high ratio of blood components (46/338, 13.6%) compared with those who did not (48/342, 14%). Another RCT (Nascimento 2013) reported a difference between patients who received high ratio of blood components (17/37, 46%) compared with patients who received a low ratio (7/32, 21.9%), however, sample sizes were small and not powered to inform the outcome of ARDS.

One systematic review (Rahouma 2017) using data from 6 cohort studies reported no significant difference on the outcome of ARDS between patients who received a high ratio of blood components (133/833, 16%) and patients who received a low ratio (199/1165, 17.1%).

One RCT (Holcomb 2015) reported no significant difference in acute kidney injury between patients who received high ratio of blood components (74/338, 21.9%) compared with those who did not (85/342, 24.9%). Similarly, one cohort study (Kim 2014) reported no difference in acute lung injury between patients who received a high ratio of blood components (1/68, 1.5%) compared to patients who received a low ratio (0/32, 0%).

One RCT (Holcomb 2015) reported no significant difference in sepsis between patients who received a high ratio of blood components (99/338, 28.9%) compared with patients who received a low ratio (91/342, 26.6%).

One RCT (Holcomb 2015) reported no significant difference in myocardial infarction between patients who received a high ratio of blood components (0/338, 0%) compared with those who did not (2/342, 0.6%).

Table 4.22 Results for high ratio of blood components versus low ratio of blood components: Patients with critical bleeding – Morbidity, thromboembolic events

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			High ratio n/N (%)	Low ratio n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting	7								
Kleinveld 2021 SR <i>Moderate</i> <i>quality</i>	N = 1187 (1 RCT) Holcomb 2015	Trauma patients (≥16 years)	Trauma centres (US, Canada)	PLT:RBC High (1:1:1) vs lower ratios	TE (DVT & symptomatic pulmonary embolus)	39/338 (11.5)	37/342 (10.8)	OR 1.08 (0.67, 1.73)	p = 0.77°
McQuilten	N = 680	Trauma patients	Trauma centres	Transfusion ratio 1:1:1	DVT	25/338 (7.4)	24/342 (7.0)	RR 1.05 (0.61,1.81)	p = 0.85°
2018 SR	(1 RCT) Holcomb 2015		(US, Canada)	versus 1:1:2	Pulmonary embolus, symptomatic	14/338 (4.1)	13/342 (3.8)	RR 1.09 (0.52,2.28)	p = 0.82°
Moderate quality					Stroke	8/338 (2.4)	11/342 (3.2)	RR 0.74 (0.30,1.81)	p = 0.50°
Paediatrics, tra	auma setting								
Kinslow 2020	N = 583 ^d	Paediatric trauma	Paediatric trauma	FFP:RBC or PLT:RBC	DVT	9/136 (6.6)	6/232 (2.6)	NR	<i>p</i> = 0.07 ^c
SR Critically low quality	(1 Coh) Butler 2019	patients with various injury severity scores	database (US)	High vs lower ratios		2:1 FFP:RBC as development	sociated with 6.9x i of DVT compared t	ncreased risk for o lower ratios	
Surgical setting	g – not identified s	tudies reported on ou	Itcome of interest						

Cl, confidence interval; Coh, cohort; DVT, deep vein thrombosis; FFP, fresh frozen plasma; NR, not reported; OR, odds ratio; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; RR, relative risk; SR, systematic review; TE, thromboembolic event; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. Information sourced from primary study

Table 4.23 Results for high ratio of blood components versus low ratio of blood components: Patients with critical bleeding – Morbidity, critical complications

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			High ratio n/N (%)	Low ratio n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
Trauma settin	g								
Kleinveld 2021 SR <i>Moderate</i> <i>quality</i>	N = 749 (2 RCTs) Nascimento 2013 Holcomb 2015	Trauma patients (paediatric and adult)	Trauma centres (US, Canada)	High PLT:RBC ratio v Iow PLT:RBC ratio	MOF	1/37 (2.7) 20/338 (5.9)	0/32 (0) 15/342 (4.4)	OR 2.67 (0.11, 67.89) OR 1.37 (0.69, 2.73)	<i>p</i> = 0.32 ^c
McQuilten	N = 680	Trauma patients	Trauma centres	High (1:1:1) vs lower	ARDS	46/338 (13.6)	48/342 (14)	RR 0.97 (0.67,1.41)	p = NR
2018	(1 RCT)	(paediatric and	(US, Canada)	(1:1:2) ratios	acute kidney injury	74/338 (21.9)	85/342 (24.9)	RR 0.88 (0.67,1.16)	p = NR
SR	Holcomb 2015	adult)			Sepsis	99/338 (28.9)	91/342 (26.6)	RR 1.10 (0.86,1.40)	p = NR
quality					myocardial infarction	0/338 (0)	2/342 (0.6)	RR 0.20 (0.01,4.20)	p = NR
Rahouma 2017 SR Critically low quality	N = 1998 (1 RCT, 6 Coh)	Trauma patients	Trauma centres (US, Canada)	High vs lower ratio	ARDS Nascimento 2013 (RCT) Brown 2012 Kim 2014 Lustenberger 2011 Sperry 2008 Undurraga 2015 Van 2010	133/833 (16.0) 17/37 (46) 47/116 (40.5) 4/68 (5.9) 10/177 (5.6) 24/102 (23.5) 22/174 (12.6) 9/159 (5.7)	199/1165 (17.1) 7/32 (21.9) 133/476 (27.9) 1/32 (3.1) 1/52 (1.9) 24/313 (7.7) 19/172 (11) 14/88 (15.9)	NR OR 0.33 (0.11, 0.95) OR 0.57 (0.37, 0.87) OR 0.52 (0.06, 4.81) OR 0.33 (0.04, 2.62) OR 0.27 (0.15, 0.50) OR 0.86 (0.45, 1.65) OR 3.15 (1.30, 7.62)	NR
	N = 100 (1 Coh) Kim 2014	Trauma patients	Trauma centres (US, Canada)		acute lung injury	1/68 (1.5)	0/32 (0)	NR	NR
Paediatrics, tr	auma setting								
Kinslow 2020 SR Critically low quality	N = 583 ^d (1 Coh) Butler 2019	Paediatric trauma patients with various ISS	Paediatric trauma database (US)	FFP:RBC or PLT:RBC High vs lower ratios	Pneumonia	NR Authors conclude >2 increased risk for de lower ratios	NR 2:1 PLT:RBC associat velopment of pneu	NR red with 23.6x monia compared to	NR

ARDS, acute respiratory distress syndrome; CI, confidence interval; Coh, cohort; FFP, fresh frozen plasma; ISS, injury severity score; MOF, multiple organ failure; NR, not reported; OR, odds ratio; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; RR, relative risk; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4 d. Information sourced from primary study.

Transfusion volumes

A summary of the evidence reported in the identified systematic reviews relating to transfusion of red blood cells and other blood components in patients with critical bleeding is presented in Table 4.24 and Table 4.25.

Red blood cells

A meta-analysis of data from 2 RCTs in the trauma setting (see Figure 4.15) showed no significant difference in median volume of RBC transfused in the first 24-hours between patients receiving a high ratio of blood components compared to patients receiving a low ratio (SMD –0.1; 95% CI –0.24, 0.05; p = 0.18, random effect, I² = 0%).

Similarly, a meta-analysis of data from nonrandomised cohort studies in the trauma setting showed showed no significant difference in median volume of RBC transfused in the first 24-hours between patients receiving a high ratio of blood components compared to patients receiving a low ratio (SMD –0.26; 95% CI –0.68, 0.15; p = 0.21, random effect, $I^2 = 92\%$).

Figure 4.15 Forest plot of comparison: high ratio vs low ratio blood components, outcome: Transfusion volume, red blood cells

	:1:1)	Low r	atio (2:	1:1)		Std. Mean Difference	Std. Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.4.1 Trauma, total units									
Holcomb 2015 (RCT) (1)	9.7	7.4	338	10.3	7.4	342	16.4%	-0.08 [-0.23, 0.07]	-
Nascimento 2013 (RCT)	7.7	3.1	37	9	6.2	32	11.4%	-0.27 [-0.74, 0.21]	
Subtotal (95% CI)			375			374	27.8%	-0.10 [-0.24, 0.05]	◆
Heterogeneity: Tau ² = 0.00	; Chi² = 0	.54, df	= 1 (P =	= 0.46); I	² = 0%				
Test for overall effect: Z = 1	I.34 (P =	0.18)							
1.4.2 Trauma, units 24 ho	urs								
Guidry 2013	193	14.8	194	13.9	11	81	14 9%	0 39 [0 13 0 65]	
Holcomb 2015 (RCT)	13.5 Q	7.4	338	10.0 Q	74	341	16.4%	0.00[-0.15, 0.05]	
Kim 2014	26	19.8	66	31	17.8	32	12.2%	-0.26 [-0.68, 0.17]	_
Kutcher 2014	7	1 7	01	10	3 75	52	13.2%	-1 13 [-1 50 -0 77]	(
Sperry 2008	16	۱. <i>۲</i>	102	22	17	313	15.5%	-0.39[-0.61 -0.16]	
Subtotal (95% CI)	10	5	791	22		819	72.2%	-0.26 [-0.68, 0.15]	
Heterogeneity: Tau ² = 0.20	; Chi² = 5	2.41, d	f = 4 (P	< 0.000	01); l² =	= 92%			
Test for overall effect: Z = 1	.24 (P =	0.21)			,.				
Total (95% CI)			1166			1193	100.0%	-0.23 [-0.50, 0.05]	
Heterogeneity: $Tau^2 = 0.11$: Chi ² = 5	3.07. d	f = 6 (P	< 0.000	01): l ² =	- 89%		• / •	
Test for overall effect: $Z = 1$.62 (P =	0.11)			,, .	2270			-2 -1 0 1 2
Test for subgroup difference	es: Chi ² =	= 0.54.	df = 1 (I	P = 0.46). $I^2 = 0$	%			Favours high (1:1) ratio Favours lower ratio
Footnotes		,			,, -				

(1) Authors reported no significant difference in median [IQR] volume of RBCs in 24 hours between high and low ratio groups (9 [5-15] vs 9 [9-16], respectively).

Other blood components

A meta-analysis of data from 2 RCTs in the trauma setting (see Figure 4.16) showed a significant increase in the volume of FFP transfused in the first 24-hours among patients who receiving a high ratio (1:1:1) of blood components compared to patients receiving a lower ratio (2:1:1) (SMD 0.3; 95% CI 0.15, 0.44; p < 0.0001, random effect, $I^2 = 0$ %).

Holcomb (2015) also suggested an increase in the volume of PLT (median 12 units vs 6 units) and CRYO (median 0 units vs 0 units) transfused among patients who received high ratio of blood components compared with those who did not, but data were skewed and the true difference is unclear.

Figure 4.16 Forest plot of comparison: high ratio vs low ratio blood components, outcome: Transfusion volume, FFP

	High ra	atio (1:	1:1)	Low ra	atio (2:	1:1)	:	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.5.4 Trauma, total plasm	a (unit)								
Holcomb 2015 (RCT) (1)	7.7	7.4	338	5.7	6	342	90.8%	0.30 [0.15, 0.45]	-
Nascimento 2013 (RCT) Subtotal (95% CI)	6	3.1	37 375	5	3.9	32 374	9.2% 1 00.0%	0.28 [-0.19, 0.76] 0.30 [0.15, 0.44]	•
Heterogeneity: $Tau^2 = 0.00$ Test for overall effect: Z = $\frac{1}{2}$); Chi² = 0. 4.02 (P < 0	00, df : 0.0001)	= 1 (P =)	= 0.96); l ²	² = 0%				
Total (95% CI)			375			374	100.0%	0.30 [0.15, 0.44]	◆
Heterogeneity: $Tau^2 = 0.00$ Test for overall effect: $Z = -$ Test for subgroup difference); Chi² = 0. 4.02 (P < 0 ces: Not ap	00, df : 0.0001) oplicab	= 1 (P =) le	= 0.96); l ^a	2 = 0%				-2 -1 0 1 2 Favours high (1:1) ratio Favours lower ratio

Footnotes (1) Authors reported median volume of other blood products in 24 hours was higher (range 0-12) for patients receiving lower ratios compared to higher ratios (range...

Table 4.24 Results for high ratio of blood components versus low ratio of blood components: Patients with critical bleeding – Transfusion volumes, red blood cells

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			High ratio Median (SD)	Low ratio Median (SD)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
Trauma setting	1								
Rijnhout 2021 SR <i>Moderate</i> quality	N = 749 (2 RCTs) Holcomb 2015 Nascimento 2013	Trauma patients with an ISS ranging between 26 and 37	Trauma, military and civilian (North America, UK, Iran)	PLT:RBC ≥1 vs PLT:RBC 0.6 or <1	RBC transfusion	NR 9.7 (7.4) (n=338) 7.7 (3.1) (n=37)	NR 10.3 (7.4) (n=342) 9 (6.2) (n=32)	MD -0.73 (-1.73, 0.28) MD -0.60 (-1.71, 0.51) MD -1.30 (-3.67, 1.07)	No significant difference p = 0.16 No significant heterogeneity $l^2 = 0\% (p = 0.60)$
Cannon 2017 SR <i>Moderate</i> <i>quality</i>	N = 1610 (1 RCT, 4 Coh studies) Holcomb 2015 Kutcher 2014 Sperry2008 Guidry 2013 Kim 2014	Adult patients with severe trauma	Various locations (North America, South Korea, Iran)	High vs Iow ratios (plasma:RBC)	RBC in 24 hours, units	NR 9 (7.4) (n=338) 7 (1.7) (n=91) 16 (9) (n=102) 19.3 (14.8) (n=194) 26 (19.8) (n=66)	NR 9 (7.4) (n=341) 10 (3.75) (n=52) 22 (17) (n=313) 13.9 (11) (n=81) 31 (17.8) (n=32)	MD -1.42 (-4.39, 1.54) MD 0.00 (-1.11, 1.11) MD -3.0 (-4.08, -1.92) MD 6.00 (-8.57, -3.43) MD 5.40 (2.23, 8.57) MD -5.00 (-12.80, 2.80)	No significant difference p = 0.35 Substantial heterogeneity $l^2=91\%$ ($p < 0.0001$)
Paediatrics, tra	uma setting – no ide	entified studies repor	ted on outcome of in	terest					

Surgical setting - no identified studies reported on outcome of interest

Cl, confidence interval; Coh, cohort; IQR, interquartile range; ISS, injury severity score; MD, mean difference; NR, not reported; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; SD, standard deviation; SR, systematic review; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

Table 4.25 Results for high ratio of blood components versus low ratio of blood components: Patients *with* critical bleeding – Transfusion volumes, other blood components*

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			High ratio Mean (SD)	Low ratio Mean (SD)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Trauma setting									
Rijnhout 2021 SR <i>Moderate quality</i>	N = 749 (RCTs) Holcomb 2015 Nascimento 2013	Trauma patients with an ISS ranging between 26 and 37	Trauma, military and civilian (North America)	PLT:RBC ≥1 versus PLT:RBC 0.6 or <1	FFP transfusion	NR 7.7 (7.4) (n=338) 6 (3.1) (n=37)	NR 5.7 (6) (n=342) 5 (3.9) (n=32)	MD 1.73 (0.87, 2.60) MD 2.00 (0.99, 3.01) MD 1.00 (-0.68, 2.68)	Favours low ratio p < 0.0001 No significant heterogeneity $l^2 = 0\% (p = 0.32)$
McQuilten 2018 SR	N = 680 (1 RCT)	Trauma patients (paediatric and/or	Trauma centres (US, Canada)	High (1:1:1) vs lower (1:1:2)	PLT ^c in 24 hours	Median (IQR) [n] 12 (6 to 18) [338]	Median (IQR) [n] 6 (0 to 12) [342]	NE	Favours low ratio p < 0.001
Moderate quality	Holcomb 2015	adult)		ratios	CRYO in 24 hours	Median (IQR) [n] 0 (0 to 0) [338]	Median (IQR) [n] 0 (0 to 9) [342]	NE	<i>Favours low ratio</i> <i>p</i> = 0.01
					Total blood products transfused to 24 hrs	Median (IQR) [n] 25.5 (NR) [338]	Median (IQR) [n] 19 (NR) [342]	NE	<i>Favours low ratio</i> p = NR
Paediatrics, traun	na setting – no ide	entified studies rep	orted on outcome	of interest					
Surgical setting -	no identified stud	lies reported on ou	tcome of interest						

* lower is better

CI, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; IQR, interquartile range; ISS, injury severity score; NE, not estimable; NR, not reported; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; SD, standard deviation; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Five- or six-unit pools of whole blood-derived platelets were considered equivalent to a unit of apheresis platelets (e.g. an adult dose of platelets).

4.5 Volume of RBC transfused (Question 4)

Question 4 - (Prognostic)

In patients at risk of critical bleeding, is the transfusion of increased volumes of RBC associated with an increased risk of mortality or adverse effects?

4.5.1 Methods

This review examined the effect of transfusion of increased volumes of RBC in patients at risk of critical bleeding (see Figure 4.17 for PICO criteria).

Figure 4.17 PPO criteria: Question 4 – effect of transfusion of increased volumes of RBC



ARDS, acute respiratory distress syndrome; RBC, red blood cells; TE, thromboembolism Notes:

a. Patients at risk of critical bleeding includes patients with penetration injuries who may not otherwise develop critical bleeding but if over-transfused before haemorrhage control may go on to do so.

b. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months).

c. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. d. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

This systematic review only considered studies published after 2009. Articles published prior to 2009 that had been included within a systematic review were also eligible for inclusion. As outlined in the protocol, this review only considered individual studies published after 2012.

An updated literature search was conducted in August 2019 and again in September 2021 to identify any new SRs or RCTs meeting the eligibility criteria.

4.5.2 Summary of evidence

4.5.2.1 Systematic Reviews

Two systematic reviews (77, 78) were included that examined the impact of transfusion of increased volumes of RBC in patients at risk of clinical bleeding. The main characteristics and quality of these systematic reviews and relevant outcomes assessed are summarised in Table 4.26.

One systematic review (Balvers 2015) identified 50 prospective and retrospective cohort studies but did not report any usable data and is therefore not reported further in this review. The authors searched for (but did not find) any studies that assessed the association between transfusion of increased volumes of RBC in patients *at risk of* critical bleeding in another setting (i.e. perioperative, obstetric, paediatric).

One systematic review (Patel 2014) was carried out in the trauma setting with no limits to trauma severity, mechanism of injury or pattern of injury. The review authors identified 40 prospective and retrospective cohort studies that assessed the association between RBC transfusion and mortality and morbidity in trauma patients. Of these, 21 studies were considered relevant to this review and were included in the meta-analysis.

A matrix illustrating the overlap of prospective and retrospective cohort studies identified in the included systematic reviews is provided in Table 4.27

Review ID <i>Review quality</i>	Study design	Population	Prognostic factor	Outcomes
Trauma setting				
Balvers 2015 (77) <i>Critically low</i>	SR / MA of RCTs, prospective and retrospective cohort studies (50 studies)	Patients aged ≥16 years with blunt or penetrating trauma injuries *mean ISS ≥ 16	Volume of RBC transfused	Morbidity
Patel 2014 (78) Low	SR / MA of prospective and retrospective cohort studies (40 studies)	Trauma patients without restrictions on trauma severity, mechanism of injury or pattern of injury	Volume of RBC transfused	Mortality Morbidity

 Table 4.26
 Characteristics and quality of systematic review evidence: increased RBC transfusion

ALI; acute lung injury; ARDS, acute respiratory distress syndrome; ICU, intensive care unit; ISS, injury severity score; MOF, multiple organ failure; RBC, red blood cells; RCTs, randomised controlled trials

		Tra	uma:	pros	spect	ive o	ohoi	rt stu	dies								Tra	uma:	retr	ospe	ctive	coh	ort s	tudie	es								
	Study ID	Bochicchio 2008	Ciesla 2005	Dunne 2004	Edens 2010	Johnson 2010	Malone 2003	Moore 1997	Sauaia 1994	Silverboard 2005	Brakenridge 2011	Brattstrom 2010	Brown 2012	Cryer 1999	Sperry 2008	Zallen 1999	Barbosa 2011	Charles 2007	Chaiwat 2009	Cotton 2009	Croce 2005	George 2008	Mahambrey	Mostafa 2004	Murrell 2005	Phelan 2010	Plurad 2007	Robinson 2005	Spinella 2008	Texeira 2008	Weinberg 2008	Dewar 2009	Lehmann 1995
iew	Patel 2014	~	~	~	~	~	~	~	~	~				х			~	х	~	~	~	x	~	x	~	~	~	~	~	~	~		
Rev	Balvers 2015																																

Table 4.27 Overlap table showing included systematic reviews and identified cohort studies: increased RBC transfusion

✓ = study included in this review

X = study did not meet the inclusion criteria for this review

--- = study identified by the systematic review authors but not included (no usable data)

4.5.2.2 Prospective cohort studies

Patel 2014 included data from 9 prospective cohort studies that investigated the association between the transfusion of increased volumes of RBC and patient outcomes in patients at risk of critical bleeding. One additional prospective cohort study (Liu 2018) was identified in this systematic review that met our inclusion criteria.

The main characteristics and quality of the prospective cohort studies included in this review is provided in Table 4.28.

Review ID Risk of bias	Study design	Population	Prognostic factor	Outcomes
Trauma settin	g	1	1	
Liu 2018 (79) <i>Serious</i>	Prospective cohort, SC	Adult trauma patients who received between 0 and 87 units of RBC within 24 hours N=131	Volume of RBC transfused	Mortality Hospital LOS
Edens 2010 <i>Moderate</i>	Prospective cohort	Military trauma patients N=66	Volume of RBC transfused	Morbidity (ALI)
Johnson 2010 <i>High</i>	Prospective cohort	Trauma patients with ISS >16 N=1440	Volume of RBC transfused	Morbidity (MOF)
Bochicchio 2008 <i>Moderate</i>	Prospective cohort	Trauma patients admitted to the ICU N=1172	Volume of RBC transfused	Mortality
Ciesla 2005 <i>Low</i>	Prospective cohort	Trauma patients with ISS >15 N=1344	Volume of RBC transfused	Morbidity (MOF)
Silverboard 2005 <i>Low</i>	Prospective cohort	Trauma patients with ISS >16 and intubation N=102	Volume of RBC transfused	Mortality
Dunne 2004 <i>Moderate</i>	Prospective cohort	Trauma patients N=9539	Volume of RBC transfused	Mortality
Malone 2003 <i>Low</i>	Prospective cohort	Any trauma patients over 18 years N=15 534	Volume of RBC transfused	Mortality
Moore 1997 <i>Low</i>	Prospective cohort	Adult trauma patients with ISS >15 N=513	Volume of RBC transfused	Morbidity (MOF)
Sauaia 1994 <i>Moderate</i>	Observational cohort (first year: retrospective; last 2 years: prospective).	Trauma patients with ISS >15 N=394	Volume of RBC transfused	Morbidity (MOF)

 Table 4.28
 Characteristics and quality of prospective cohort evidence: increased RBC transfusion

ALI; acute lung injury; ICU, intensive care unit; ISS, injury severity score; LOS, length of stay; MOF, multiple organ failure; RBC, red blood cells; SC, single centre

Trauma setting

Among the 9 prospective cohort studies identified by Patel 2014, there were 4 studies (Bochicchio 2008, Silverboard 2005, Dunne 2004, Malone 2003) that assessed the effect of RBC on mortality, 4 studies (Ciesla 2005, Johnson 2010, Moore 1997, Sauaia 1994) that assessed the effect of RBC on MOF and one study (Edens 2010) that assessed the effect of RBC on acute lung injury. Meta-analyses were conducted to determine the effect of increased volume of RBC transfusions on each of the outcome measures.

The studies were conducted in the trauma settings and commonly queried trauma databases or registries, resulting in most studies having good representativeness. Overall, Patel 2014 considered there to be no serious concerns of bias among the included prospective cohort studies.

Liu 2018 was a single centre prospective cohort study conducted in the US that investigated the association between RBC transfusion and mortality and hospital LOS in the trauma setting. Included trauma patients (predominantly due to assault and motor vehicle accidents) were over 18 years and had received between 0 and 87 units of RBC within 24 hours of injury. The study was had serious concerns of bias raised, relating to inadequate adjustment for confounders, and lack of details regarding study design.

4.5.2.3 Retrospective cohort studies

Patel 2014 included data from 12 retrospective cohort studies that investigated the association between the transfusion of increased volumes of RBC and patient outcomes in patients at risk of clinical bleeding. One additional retrospective cohort study (Hassanien 2015) (80) was identified in this systematic review that met our inclusion criteria.

The main characteristics and quality of the retrospective cohort studies included in this review is provided in Table 4.29.

Trauma setting

Among the 12 retrospective cohort studies identified by Patel 2014, there were 10 studies (Barbosa 2011, Chaiwat 2009, Mahambrey 2009, Murrell 2005, Phelan 2010, Robinson 2005, Spinella 2008, Croce 2005, Teixeira 2008, Weinberg 2008) that assessed the effect of RBC on mortality, one study (Cotton 2009) that assessed the effect of RBC on MOF and 3 studies (Plurad 2007, Weinberg 2008, Croce 2005) that assessed the effect of RBC on acute respiratory distress syndrome (ARDS). Meta-analyses were conducted to determine the effect of increased volume of RBC transfusions on each of the outcome measures.

The studies were conducted in the trauma settings and commonly queried trauma databases or registries, resulting in most studies having good representativeness. Overall, Patel 2014 considered there to be no serious risk of bias of included studies but noted that the study design is prone to confounding bias (particularly in relation to adjusting for the injury severity scores). Review authors attempted to mitigate confounding by only including studies that attempted to adjust for injury severity in the pooled analysis.

Medical setting

Hassanien 2015 was a retrospective hospital-based study conducted in Egypt. The study included 70 patients with liver cirrhosis and hepatocellular carcinoma presenting with

acute upper gastrointestinal bleeding. Patients must meet criteria of either hematemesis or melena with a diagnostic esophagogastroduodenoscopy, or both. The study was at moderate risk of bias due to a lack of details regarding blinding and study design.

Table 4.29	Characteristics and quality of retrospective cohort evidence: increased RBC
	transfusion

Review ID Risk of bias	Study design	Population N	Prognostic factor	Outcomes	
Trauma setting		1			
Barbosa 2011 <i>Low</i>	Retrospective I cohort	Massively transfused patients N=704	Volume of RBC transfused	Mortality	
Phelan 2010 <i>Low</i>	Retrospective cohort	Adult trauma patients N=399	Volume of RBC transfused	Mortality	
Chaiwat 2009 <i>Moderate</i>	Retrospective cohort	Adults with at least one injury >AIS 3 N=14 070	Volume of RBC transfused	Mortality	
Cotton 2009 <i>Moderate</i>	Retrospective cohort	Surgical patients requiring massive transfusion N=266	Volume of RBC transfused	Morbidity (MOF)	
Mahambrey 2009 <i>Low</i>	Retrospective cohort	Trauma patients requiring massive transfusion N=260	Volume of RBC transfused	Mortality	
Spinella 2008 <i>Serious</i>	Retrospective cohort	Military trauma patients N=708	Volume of RBC transfused	Mortality	
Texeira 2008 <i>Serious</i>	Retrospective cohort	Trauma patients N=25 599	Volume of RBC transfused	Mortality	
Weinburg 2008 <i>Moderate</i>	Retrospective cohort	Patients with blunt trauma and ISS <25 N=1624	Volume of RBC transfused	Mortality Morbidity (ARDS)	
Plurad 2007 <i>Moderate</i>	Retrospective cohort	ICU trauma patients with intubation N=2346	Volume of RBC transfused	Morbidity (ARDS)	
Croce 2005 Low	Retrospective cohort	Patients with blunt trauma requiring ICU admission N=5260	Volume of RBC transfused	Mortality Morbidity (ARDS)	
Murrell 2005 <i>Low</i>	Retrospective cohort	Trauma patients transfused with at least one unit N=275	Volume of RBC transfused	Mortality	
Robinson 2005 <i>Low</i>	Retrospective cohort	Patients with blunt hepatic/splenic injury N=319	Volume of RBC transfused	Mortality	
Medical setting					
Hassanien 2015 (80) <i>Moderate</i>	Retrospective cohort, SC	Adult patients with liver cirrhosis and HCC presenting with acute upper gastrointestinal bleeding N=70	Volume of RBC transfused	Mortality	

AIS, abbreviated injury scale; ARDS, acute respiratory distress syndrome; HCC, hepatocellular carcinoma; ICU, intensive care unit; ISS, injury severity score; MOF, multiple organ failure; RBC, red blood cells; SC, single centre

4.5.3 Results

4.5.3.1 Mortality

A summary of the evidence reported in the identified literature relating to mortality in patients at risk of critical bleeding is presented in Table 4.30.

The identified literature suggests transfusion of increased RBC is associated with an increased risk of mortality among patients at risk of critical bleeding in the trauma setting.

Nine studies identified by Patel 2014 assessed RBC transfusion as a continuous variable. Pooled analysis showed an increased in the odds of mortality with each additional RBC unit transfused (OR 1.07; 95% CI 1.04, 1.10; p < 0.001; random effects; $I^2 = 82.9\%$).

Similarly, Liu 2018 showed increasing odds of mortality with increasing units of RBC transfused.

Due to the limited evidence and significant heterogeneity, no additional meta-analysis was performed.

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results							
Study design ^a	(no. of trials) included in analysis	population	(Location)			High volume RBC n/N (%)	Low volume RBC n/N (%)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b				
Trauma setting													
Patel 2014 SR <i>Low quality</i>	N = 18 009 (9 Coh) Barbosa 2011 Bochicchio 2008 Chaiwat 2009 Mahambrey 2009 Murrell 2005 Phelan 2010 Robinson 2005 Spinella 2008 Silverboard 2005	Trauma patients, not limited by trauma severity, mechanism or pattern of injury	Trauma and non-trauma centres, (US, Canada)	RBC transfusion as continuous variable vs no RBC transfusion	Mortality	NR	NR	OR 1.07 (1.04-1.10) 1.05 (1.03, 1.07) 1.05 (1.03, 1.07) 1.05 (1.00, 1.10) 1.01 (0.97, 1.05) 0.83 (0.69, 0.99) 1.13 (1.10, 1.16) 1.16 (1.01, 1.24) 1.16 (1.09, 1.25) 1.08 (1.04, 1.15)	Favours no RBC transfusion p < 0.001 Substantial heterogeneity I ² = 82.9%; p < 0.0001				
						The odds of morta transfused	ality increased with e	ach additional unit					
	N = 57 875 (6 Coh) Croce 2005 Dunne 2004 Malone 2003 Robinson 2005 Texeira 2008 Weinberg 2008	-		RBC transfusion as dichotomous variable vs no RBC transfusion	Mortality	NR	NR	OR 3.15 (1.82–5.46) 2.46 (2.00, 3.20) 4.23 (3.07, 5.84) 2.83 (1.82, 4.40) 4.75 (1.37, 16.40) 6.70 (6.10, 7.50) 0.96 (0.48, 1.94)	Favours no RBC transfusion p < 0.001 Substantial heterogeneity I² = 94.6%; p < 0.0001				
Liu 2018 Prospective Serious risk of bias	N = 131	Adult trauma patients	Trauma, single centre (US)	High vs low units of RBC	Mortality 0-9 units (n=95) 10-19 units (n=19) 20-29 units (n=8) 30-39 units (n=4) 40+ units (n=5)	4/19 (21) 3/8 (38) 2/2 (50) 4/5 (80)	23/95 (24)	OR 0.83 (0.25, 2.77) OR 1.88 (0.42, 8.47) OR 3.13 (0.41, 23.49) OR 12.52 (1.33, 117.7)	OR for 40+ units was 12.52 and did not contain the null, indicating a statistically significant difference from control (0-9 units)				
Medical setting													
Hassanien 2015 Retrospective Moderate risk of bias	N = 70	Patients with liver cirrhosis and HCC presenting with AUGI bleeding	Single medical centre (Egypt)	Survivor vs non- survivor	Units of RBC transferred	Survivor (n=32) 1.9 ± 0.23	Non-survivor (n=38) 2.6 ± 0.74	NR	<i>p</i> < 0.01				

Table 4.30 Results for increased volume of RBC transfused versus decreased volume of RBC transfused: Patients at risk of critical bleeding - Mortality

AUGI; acute upper gastrointestinal; CI, confidence interval; Coh, cohort; HCC, hepatocellular carcinoma; NR, not reported; OR, odds ratio; RBC, red blood cells; SR, systematic review; US, United States a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

4.5.3.2 Morbidity

A summary of the evidence reported in the identified literature relating to mortality in patients at risk of critical bleeding is presented in Table 4.31.

Identified literature suggests transfusion of increased RBC is associated with an increased risk of morbidity among patients at risk of critical bleeding in the trauma setting.

Multiple organ failure

Three studies identified by Patel 2014 assessed RBC transfusion as a continuous variable. Pooled analysis showed a significant increase in the odds of MOF with each additional RBC unit transfused (OR 1.08; 95% Cl 1.02, 1.14; p = 0.012; random effects; $l^2 = 95.9\%$).

Due to the limited evidence and significant heterogeneity, no additional meta-analysis was performed.

Acute respiratory distress syndrome / acute lung injury

Two studies identified by Patel 2014 assessed RBC transfusion as a continuous variable. Pooled analysis showed a significant increase in the odds of ARDS or ALI with each additional RBC unit transfused (OR 1.06; 95% CI 1.03, 1.10; p < 0.001; random effects; $I^2 = 0$ %).

Due to the limited evidence and significant heterogeneity, no additional meta-analysis was performed.

Length of stay

One cohort study by Liu (2018) found no association between the transfusion of increased RBC and length of stay in trauma patients.

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			High volume RBC mean ± SD	Low volume RBC mean ± SD	Risk estimate (95% Cl)	Statistical significance <i>p</i> -value Heterogeneity ^b
Trauma setting	g								
Patel 2014 SR <i>Low quality</i>	N = 3050 (3 Coh) Ciesla 2005 Cotton 2009 Johnson 2010	Trauma patients, not limited by trauma severity, mechanism or	Trauma (US)	RBC transfusion as continuous variable vs no RBC transfusion	MOF	NR	NR	OR 1.08 (1.02, 1.14) 3.40 (2.53, 4.58) 2.90 (1.20, 6.70) 8.60 (4.20, 17.70)	Favours no RBC transfusion p = 0.012 Substantial heterogeneity $l^2 = 95.9\%; p < 0.0001$
	N = 2251 (3 Coh) Ciesla 2005 Moore 1997 Sauaia 1994	pattern of injury		RBC transfusion of > 6 units as a dichotomous variable vs no transfusion	-	NR	NR	OR 4.30 (2.36, 7.85) 3.40 (2.53, 4.58) 2.90 (1.20, 6.70) 8.60 (4.20, 17.70)	Favours RBC transfusion ≤ 6 units p < 0.0001 Moderate heterogeneity $l^2 = 65.9\%; p = 0.053$
	N = 14 136 (2 Coh) Chaiwat 2009 Edens 2010	-		RBC transfusion as continuous variable vs no RBC transfusion	ARDS/ALI	NR	NR	OR 1.06 (1.03–1.10) 1.06 (1.03, 1.10) 1.09 (0.74, 1.58)	Favours no RBC transfusion p < 0.001 No heterogeneity $l^2 = 0.0\%$, $p = 0.886$
	N = 9230 (3 Coh) Plurad 2007 Weinburg 2008 Croce 2005			RBC transfusion as a dichotomous variable vs no transfusion	_	NR	NR	OR 2.04 (1.47, 2.83) 1.98 (1.38, 2.83) 1.96 (0.73, 5.26) 3.42 (2.02, 34.20)	Favours no RBC transfusion p < 0.001 No heterogeneity l² = 0.0%, p = 0.761
Liu 2018 Prospective Serious risk of bias	N = 131	Adult trauma patients	Trauma, single centre (US)	High vs low units of RBC	LOS, mean ±SD 0-9 units (n=95) 10-19 units (n=19) 20-29 units (n=8) 30-39 units (n=4) 40+ units (n=5)	9.3 ± 5.5 9.0 ± 8.0 6.8 ± 6.0 4.6 ± 6.2	10.1 ± 12.1	NR	No significant difference p = 0.793 p = 0.806 p = 0.588 p = 0.321

Table 4.31 Results for increased volume of RBC transfused versus decreased volume of RBC transfused: Patients at risk of critical bleeding - Morbidity

Medical setting - no identified studies reported outcome of interest

ALI, acute lung injury; ARDS, acute respiratory distress syndrome; CI, confidence interval; Coh, cohort; LOS, length of stay; MOF, multiorgan failure; NR, not reported; OR, odds ratio; RBC, red blood cells; SD, standard deviation; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

4.5.3.3 Time on mechanical ventilator

No studies identified.

4.6 Recombinant factor VIIa (Question 5)

Question 5 - (interventional)

In patients *with* critical bleeding, what is the effect of rFVIIa treatment on morbidity, mortality and transfusion rate?

4.6.1 Methods

This review examined the effect of rFVIIa treatment on outcomes in patients with critical bleeding (i.e. major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion) as outlined in Figure 4.18.

For this question there was particular focus on *patients who failed to achieve adequate haemostasis despite surgical management and appropriate blood component therapy.* Studies in patients with haemophilia and studies that examined the prophylactic use of rFVIIa were not eligible for inclusion.





Abbreviations: ARDS, acute respiratory distress syndrome; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; PLT, platelets; RBC, red blood cells; rFVIIa, recombinant activated factor seven; TE, thromboembolic event

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other.

c. Newborns up to 28 days following birth.

d. rFVIIa is approved in Australia and NZ for the control of bleeding and prophylaxis for surgery in patients with specific clotting disorders. Use outside these indications (including critical bleeding following trauma) is considered 'off-label'.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2009, noting primary studies published prior to 2009 that had been included in a systematic review were also eligible for inclusion. Nonrandomised studies (with concurrent or noncurrent controls) and observational cohort studies were excluded. There were no restrictions applied to study sample size.

Assuming all relevant primary studies had been identified in the included systematic review studies; the systematic screen for RCTs was limited to studies published from January 2015. This was based on the literature search date of the most recent identified systematic review (Canon 2017), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting. A targeted search²⁹ for studies that focused on critical bleeding in the obstetrics and surgical setting and had been published between 2010 and 2015 was also conducted.

The literature search was updated in August 2019 with no new systematic review found.

In March 2021 it was agreed that this question would be retired, as research in this field is not evolving and updates to the literature search would likely find no new evidence (i.e. the citations in the September 2021 literature search update relating to rFVIIa were not screened).

4.6.2 Summary of evidence

4.6.2.1 Systematic reviews

Eight systematic reviews (61, 81-87) were included that evaluated the effects of rFVIIa treatment in patients with critical bleeding. The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.32.

Five systematic reviews were focused on critical bleeding in the trauma and non-trauma setting (Cannon 2017, Curry 2011, McQuilten 2015, Simpson 2012; Yank 2011), one in paediatric surgical trauma (Okonta 2012), and 2 in the obstetric setting (Franchini 2010, Magon 2012).

A matrix illustrating the overlap of studies included in each review is provided in Table 4.33 and Table 4.34. Among the 48 publications identified by the included systematic reviews, there were 3 RCTs and 34 nonrandomised cohort studies that were not included in the evidence evaluation as they did not meet the review criteria for this question (see Section 3.1.4).

4.6.2.2 Randomised controlled trials

There were 11 citations related to 9 RCTs (88-98) that were considered relevant to this review. The primary studies varied in the clinical setting, and included trauma (blunt or penetrating), surgical (cardiac), medical emergency and obstetrics. All identified studies were supported by the manufacturer.

Two additional RCTs were identified through the handsearching process. One RCT sponsored by the manufacturer (Novo Nordisk A/S; NCT00323570) was withdrawn (and merged with the RCT reported by Hauser 2010). The other evaluated the safety and effectiveness of rFVIIa in women with severe primary postpartum haemorrhage (99).

A summary of the characteristics and quality of all identified RCTs is provided in Table 4.35.

²⁹ Keyword search for "obstetrics" and "maternity" or "cardiac" in identified studies.

Review ID	Study design	Population	Intervention	Comparator	Outcomes
Review quality					
Trauma and no	n-trauma settir	ng			
Cannon 2017 (61) <i>Moderate</i>	SR of RCTs and cohort studies	Patients at risk of death from haemorrhage *defined as patients requiring blood transfusions and/or injury severity score greater than 25	rFVIIa	NorFVIIa	Mortality Morbidity (TE) Blood components used Massive transfusion
McQuilten 2015 (81) <i>Moderate</i>	SR review of SRs and RCTs	Patients with critical bleeding who had received or were anticipated to receive a massive transfusion in any clinical setting	rFVIIa	No rFVIIa	Mortality Morbidity (hospital LOS, serious AE, transfusion- related AE) Transfusion volume
Simpson 2012 <i>High</i> (84)	SR of RCTs	Patients who had received treatment to manage bleeding (medical, surgical, or obstetric)	rFVIIa	Placebo or different dose	Mortality Morbidity (TE) Transfusion volume
Curry 2011 (85) <i>Moderate</i>	SR of RCTs	Trauma patients with haemorrhagic shock within the first 24 hours of injury	rFVIIa	NorFVIIa	Mortality Morbidity (MOF, ARDS) Transfusion volume
Yank 2011ª (86, 100) <i>High</i>	SR of RCTs, cohort studies, case series and case reports	Hospitalised patients with off-label use in cardiac surgery, body trauma, ICH, TBI, liver transplantation and prostatectomy ^b	rFVIIa	Placebo, alternative therapies or usual care	Mortality Morbidity (ARDS, TE) Transfusion volume
Surgical setting	1				
Okonta 2012 (83) <i>Critically low</i>	SR of best available evidence	Paediatric patients with excessive bleeding after cardiac surgery	rFVIIa	Placebo or other dose	Mortality Morbidity (TE) Transfusion volume
Obstetrics and	maternity settii	ng			
Magon 2012 (82) Critically low	SR of best available evidence	Women with major postpartum haemorrhage	rFVIIa	Not specified	Mortality Transfusion volume
Franchini 2010 (87) <i>Critically low</i>	SR of best available evidence	Women with major postpartum haemorrhage	rFVIIa	Not specified	Mortality Transfusion volume

Table 4.32 Characteristics and quality of systematic review evidence: rFVIIa

Abbreviations: AE, adverse events; ARDS, acute respiratory distress syndrome; ICH, intracranial haemorrhage; LOS, length of stay; MOF, multiorgan failure; RCT, randomised controlled trial; SR, systematic review; TBI, traumatic brain injury, TE, thromboembolic events

a. An updated 2016 report (100) of the published systematic review by Yank 2011 also reviewed.

b. Data for ICH, TBI, liver transplantation and prostatectomy not relevant for this review (not critical bleeding or prophylactic use).

		Traun	าล					Medi	cal er	nerge	Surgical (cardiac)					
	Study ID	Dutton 2011ª	Hauser 2010	Boffard 2005a&b	McMullin 2010 ^b	Boffard 2009 ^b	Rizoli 2006(a) ^b	Narayan 2008 °	Bosch 2008	Chuansumrit 2005	Pihusch 2005	sosch 2004	Gill 2009	Ma 2006	Diprose 2005	
	Cannon 2017		~	~												
	McQuilten 2015	✓	✓	✓												
₽	Simpson 2012		~			~		Х	~	×	×	~				
3	Okonta 2012 ^d															
Revie	Magon 2012 d															
	Curry 2011		~	~	~	~	~									
	Yank 2011		~	~									~	Х	Х	
	Franchini 2010 d															

 \checkmark = study included in this review X = study did not meet the inclusion criteria for this review.

a. Dutton 2011 and Hauser 2010 report data from the same randomised trial (CONTROL) b. Post-hoc analysis of Boffard 2005a&b.

c. Population out of scope. Adult patients with traumatic intracranial haemorrhage with contusion.

d. Authors did not identify any RCTs in the population of interest.

		Trau	ıma					Surgical (cardiac, adult)							Surgical (cardiac, paediatric)												Obstetric and maternity									
	Study ID	Wade 2010	Woodruff 2010	Fox 2009	Spinella 2008	Rizoli 2006(b)	Harrison 2005	Mitra 2010	Gelsomino 2008	Trowbridge 2008	Tritapepe 2007	Karkouti 2005	Von Heymann 2005	Ekert 2006 ^a	Warren 2009 (SR)	Singh 2012	Karsies 2010	Guzzetta 2009	Niles 2008	Agarwal 2007	Veldman 2007	Kylasam 2006	Razon 2005	Egan 2004	Pynchynka-Pokorska	Tobias 2004	Isbister 2008	Bouma 2008	Ahonen 2007	Alfirevic 2007	Barillari 2007	Hossain 2007	Ahonen 2005	Sobieszczyk 2006	Segal 2004	
	Cannon 2017				Х	Х	X																													
	McQuilten 2015 ^b																																			
٥	Magon 2012																										No	stud	ies ic	lenti	fied					
Ň	Okonta 2012													x	Х	x	Х	х	х	X	х	х	Х	Х	X	Х										
evie	Simpson 2012 ^b																																			
Ř	Curry 2011 ^b																																			
	Yank 2011	Х	Х	Х	X	Х		Х	Х	Х	Х	Х	Х																							
	Franchini 2010																										Х	Х	Х	Х	Х	Х	Х	x	Х	

Table 4.34 Overlap table of nonrandomised cohort studies identified by included systematic reviews: rFVIIa

✓ = study included in this review X = study did not meet the inclusion criteria for this review (nonrandomised studies not eligible for this question).

a. Intervention out of scope. Prophylactic use.

b. Authors did not search for nonrandomised studies.
Study ID Risk of bias	Study design	Population	Intervention	Comparator	Outcomes
Trauma settin	g				
Hauser 2010 (CONTROL) (89) <i>Unclear</i>	RCT	Adult patients (aged 18 to 70 years) with major haemorrhage from blunt or penetrating trauma who received 4 to 8 units of RBC within 12 hours of injury	rFVIIa at 0, 1, 3 hrs (total 400 µg/kg)	Placebo	Mortality Morbidity (MOF, SOF, ARDS, TE, DIC, SAEs) Blood component use
Boffard 2005a (90, 91, 94, 95) <i>High</i>	RCT	Adult patients with haemorrhage from a severe blunt traumatic injury requiring \geq 6 units of RBC within 4 hours of hospitalisation	rFVIIa at 0, 1, 3 hrs (total 400 µg/kg)	Placebo	Mortality Morbidity (MOF, ARDS) Number of RBC transfused
Boffard 2005b (90, 91, 94, 95) <i>High</i>	RCT	Adult patients with haemorrhage from a severe penetrating traumatic injury requiring ≥ 6 units of RBC within 4 hours of hospitalisation	rFVIIa at 0, 1, 3 hrs (total 400 µg/kg)	Placebo	Mortality Morbidity (MOF, ARDS) Number of RBC transfused
Medical					
Bosch 2008 <i>Unclear</i> (93)	RCT	Adult patients with cirrhosis and upper gastrointestinal haemorrhage	rFVIIa at 0, 2, 8, 14 & 20 hrs (total 1000 µg/kg)	Placebo	Mortality Morbidity (AE) RBC transfusion volume
Bosch 2004 (98) <i>Unclear</i>	RCT	Adult patients with cirrhosis and upper gastrointestinal haemorrhage	rFVIIa at 0, 2, 4, 6, 12, 18, 24 & 30 hrs (total 800 µg/kg)	Placebo	Mortality Morbidity (AE) RBC transfusion volume
Haematology/	oncology				
Chuansumrit 2005 (96) <i>High</i>	RCT	Children with dengue haemorrhagic fever	rFVIIa 100 µg/kg additional dose allowed after 30 minutes	Placebo	Transfusion volume (RBC, PLT, FFP)
Pihusch 2005 (97) <i>High</i>	RCT	Patients (aged >12 yrs.) with bleeding occurring 2 to 180 days after HSCT	rFVIIa 40, 80 or 160 µg/kg every 6 hrs for 36 hrs	Placebo	Morbidity (AE) Transfusion volume (RBC, PLT, FFP)
Surgical settin	g				
Gill 2009 (92) <i>Unclear</i>	RCT	Patients with bleeding after cardiac surgery ^a	rFVIIa 40, 80 or 160 µg/kg	Placebo	Morbidity (SAEs) Transfusion volume (RBC, PLT, FFP)
Obstetrics and	l maternit	y			
Lavigne- Lissalde 2015 (99) <i>High</i>	RCT	Women with severe PPH ^b after vaginal or caesarean birth that persisted after sulprostone treatment	rFVIIa 60 μg/kg	SoC ^c	Mortality Morbidity (TE, reduction in need for specific second-line therapies)

Table 4.35 Characteristics and	d quality of RCT evidence: rFVIIa
--------------------------------	-----------------------------------

Abbreviations: AE, adverse events; ARDS, acute respiratory distress syndrome; DIC, disseminated intravascular coagulation; FFP, fresh frozen plasma; HSCT, haematopoietic stem cell transplant; MOF, multiorgan failure;; RCT, randomised controlled trial; rFVIIa, recombinant activated factor seven; PLT, platelets; PPH, primary postpartum haemorrhage; SAEs, serious adverse events; SoC, standard of care; SOF, single organ failure; TE, thromboembolic events

a. defined as post-operative bleeding for a minimum of 30 minutes following completion of a stabilisation period, and meeting at least one of the following criteria: \geq 200 ml/hr in any one hour or part thereof; or \geq 2 ml/kg/hr for 2 consecutive hours.

b. defined as the loss of more than 1500 mL of blood within 24 hours.

c. compassionate rFVIIa given late to avoid emergency peripartum hysterectomy.

Trauma

Three RCTs (Boffard 2005a & b, Hauser 2010) were identified that examined the effect of rFVIIa in patients with critical bleeding after blunt or penetrating trauma (89, 95). All 3 RCTs were judged to be of an overall unclear to high risk of bias (84), with high threats to validity due to lack of details provided by Boffard 2005a and 2005b (selective reporting) or unclear blinding of outcome assessment in Hauser 2010, which may have favoured the intervention.

Two parallel, double-blind RCTs were run simultaneously and published in the one article (95). The studies enrolled patients with haemorrhage from a blunt (Boffard 2005a) or penetrating (Boffard 2005b) traumatic injury requiring a least 6 unit of RBC within 4 hours of hospitalisation. The studies were sponsored by the manufacturer and enrolled 301 patients (143 blunt and 134 penetrating) from 32 centres across 8 countries (including Australia, Canada, France, Germany, Israel, Singapore, South Africa and the UK). Both RCTs censored deaths that occurred within 48 hours (comprising nearly 20% of patients) as the primary outcomes were RBC transfusion needs during the 48-hour observation period, which indicates that some end-stage use of rFVIIa may have occurred. Mortality and morbidity (ARDS, TE) were also reported, noting the studies were not powered to detect a difference in these outcomes. Post-hoc analyses on the effect of rFVIIa on coagulopathic patients (94), on trauma patients who survived the first 48 hours after randomisation (91), and exploring the association between poorer outcomes and baseline haematologic and coagulation parameters (90) were also identified.

The double-blind RCT published by Hauser 2010 (CONTROL) enrolled patients with blunt or penetrating trauma who, despite strict damage control resuscitation and operative management had continued bleeding after receiving 4 units of RBC within 12 hours of injury. The study was sponsored by the manufacturer and enrolled 573 patients (481 blunt and 92 penetrating) from 150 hospitals in 26 countries. Subgroup analyses on patients with blunt (Hauser 2010a) and penetrating (Hauser 2010b) trauma were also conducted. The aim of the study was to detect a 16.7% mortality reduction with rFVIIa, assuming a 30% mortality in placebo patients, however, the study was terminated early due to unexpectedly low mortality in the placebo group detected during planned interim futility analysis. Extended safety data on patients enrolled in CONTROL are also available (88).

The 3 RCTs evaluated a total dose of 400 μ g/kg intravenous rFVIIa administered in 3 doses (200 μ g/kg at 0 hour, 100 μ g/kg at one and 3 hours); which is higher than that reported among trauma patients in the Australian and New Zealand Haemostasis Registry, with 76% of patients (352/461) receiving only a single dose (median first dose of 95 μ g/kg; IQR 80 to 108) (101). Patients enrolled in Hauser 2010 received the first dose earlier during the resuscitation period (after the fourth unit of RBC) and required participating hospitals to use a prespecified resuscitation protocol.

Medical emergency

Two RCTs (Bosch 2004, Bosch 2008) were identified in the medical emergency setting that evaluated the therapeutic use of rFVIIa in patients with cirrhosis presenting with upper gastrointestinal haemorrhage (93, 98). Both RCTs were assessed to be at low to

unclear risk of bias, predominantly due to lack of clear detail and poor reporting in the published reports (84).

The RCT reported by Bosch 2004 was conducted in 245 cirrhotic patients with upper gastrointestinal bleeding (UGIB) enrolled from 26 centres in Europe. Subject were administered 100 μ g/kg rFVIIa 8 times *before* first endoscopy (t0), then at 2, 4, 6, 12, 18, 24, and 30 hours *after* endoscopy (total dose: 800 μ g/kg total), with follow-up of patients occurring through to 42 days.

In the second RCT reported by Bosch 2008, 256 patients with advanced cirrhosis and active variceal bleeding were enrolled from 31 hospitals across Europe and Asia. Patients were randomised to receive 200 μ g/kg rFVIIa initially as soon as possible *after* endoscopy, then either 4 x 100 μ g/kg (total dose: 600 μ g/kg) or a single 100 μ g/kg (total dose: 300 μ g/kg), or placebo; with the subsequent doses given at 2, 8, 14 and 20 hours after the first dose.

In both trials, the total dose of rFVIIa is again notably higher than that reported among patients with UGIB in the Australian and New Zealand Haemostasis Registry, with 74% of patients (140/189) receiving only a single dose (median first dose of 89 μ g/kg; IQR 67 to 104) (101). The primary outcome measures in both trials were a composite of failure to control UGIB within 24 hours after first dose, failure to prevent rebleeding between 24 hours and day five, or death within 5 days. Outcomes of relevance for this review were transfusion requirements within 5 days (at discharge), and mortality and thromboembolic events recorded at latest follow-up.

Haematology/oncology setting

One multicentre RCT (Pihusch 2005) was identified that evaluated the use of rFVIIa in 100 patients with moderate or severe bleeding complications following haematopoietic stem cell transplantation (+2 to +180 weeks post-transplant) (97). Patients with bleeding (52 gastrointestinal; 26 haemorrhagic cystitis; 7 pulmonary; one cerebral; 14 other) were randomised to receive 7 doses of rFVIIa at 40, 80 or 160 μ g/kg (total dose: 280, 560, or 1120 μ g/kg) or placebo every 6 hours. The primary efficacy endpoint was the change in bleeding score between the first administration and 38 hours. The study was at high risk of bias due to baseline difference observed between treatment groups, suggesting randomisation or allocation concealment was compromised (84).

One RCT (Chuansumrit 2005) conducted in 25 paediatric patients with active bleeding due to dengue fever was identified in the literature (96). Patients were administered 100 μ g/kg rFVIIa with repeat dose at 30 minutes if ongoing bleeding was observed. The study was small and not powered to detect differences in any outcomes and was therefore considered to be at high risk of bias for all outcomes (84).

Surgical setting

One Phase II dose-escalation study (Gill 2009) conducted in 13 countries in Africa, Asia, Europe, South America and US was identified that evaluated the therapeutic use of rFVIIa in patients with intractable bleeding after cardiac surgery (92). Patients were randomised to receive either 40 or 80 μ g/kg rFVIIa (n= 35 and n=69, respectively) or placebo (n=68) after cardiopulmonary bypass (CPB) as treatment for excessive post-operative bleeding in the ICU.

The trial was terminated in November 2007 without proceeding to the highest dosing cohort (160 μ g/kg) as it was determined to no longer reflect common clinical practice. The primary outcome was the incidence of critical serious adverse events at 30 days. The RCT was had overall unclear risk of bias (84).

Obstetrics and maternity

One multicentre RCT (Lavigne-Lissalde 2015) was identified that assessed the safety and effectiveness of rFVIIa given to women with severe primary postpartum haemorrhage (PPH), defined as loss of more than 1500 mL of blood within 24 hours after birth, after sulprostone failure (99). The women were aged over 18 years and had delivered after the end of 27 weeks of gestation by either vaginal or Caesarean section.

Subjects were randomly assigned to receive a single dose of 60 µg/kg rFVIIa or not, with the primary outcome being a reduction in the need for specific second-line therapies (inclusive of arterial embolisation, hysterectomy). Safety outcomes were also recorded up to 5 days post infusion.

The study was assessed to be at high risk of bias due to non-blinding that seriously weakens confidence in the results. The study allowed for compassionate use of rFVIIa in the comparator arm (8 out of 42 women in the standard care group received late rFVIIa) so it is possible that this introduced bias into the subsequent management of patients.

4.6.3 Results

4.6.3.1 Mortality

A summary of the evidence relating to in-hospital mortality in patients with critical bleeding treated with rFVIIa is presented in Table 4.36.

For most bleeding patients there does not seem to be clear significant survival benefits associated with rFVIIa (*GRADE: low or very low*).

A meta-analysis of data from RCTs included in this review (see Figure 4.19) showed the mortality rate (latest timepoint) in patients with critical bleeding to be comparable among those who received rFVIIa (157/934, 16.8%) and those who did not (120/776, 15.5%) with a relative risk (RR) of 0.99 observed (95% CI 0.80, 1.23; p = 0.84; fixed effect, I² = 0%). For most bleeding patients, there is little or no difference in mortality compared with placebo or no rFVIIa.

Among patients with blunt and penetrating trauma, a total of 409 patients received rFVIIa compared with 428 patients who did not, with no difference in mortality observed (16.6% vs 17.1%, RR 0.96; 95% CI 0.71, 1.29; p = 0.71; fixed effect, $I^2 = 0\%$) (*GRADE: low*).

Among patients with UGIB who received rFVIIa, the mortality rate of 19.2% (55/286) was not significantly different from the mortality rate of 17.5% (36/206) observed among those who did not receive rFVIIa. This corresponded to a RR of 1.02 (95% CI 0.55, 1.90; p = 0.95; random effects, $l^2 = 56\%$)(*GRADE: very low*).

Among patients with uncontrolled bleeding due to other medical conditions (after HSCT, Dengue fever), the mortality rate was 25.8% (24/93) among those who received rFVIIa, compared with 21.9% (7/32) in those who did not, corresponding to a RR of 1.02 (95% CI 0.51, 2.07; p = 0.95; fixed effects, I² = not applicable [one study]) (*GRADE: very low*).

Among patients with intractable bleeding after cardiac surgery, the mortality rate among those who received rFVIIa (9.6%)³⁰ was higher than that observed among those who did not receive rFVIIa (5.9%); however, this difference was not significant (RR 1.63; 95% CI 0.53, 5.00; p = 0.95; fixed effects, I² = not applicable [one study]) (*GRADE: very low*).

No deaths were observed in the RCT that assessed the effects of rFVIIa among women with severe PPH with persistent bleeding after sulprostone treatment and the included RCT was not large enough to detect differences in mortality (*GRADE: very low*).

In agreeance with our findings, all identified systematic reviews reported no significant difference in mortality between patients who received in rFVIIa compared with those who did not, regardless of clinical setting (61, 81-87). There was also no significant effect on mortality shown in any of the subgroup analyses.

³⁰ noting the mortality rate among patients administered 40 and 80 µg/kg rFVIIa was 11.4% (4/35) and 8.7% (6/69), respectively.

Figure 4.19 Forest plot of comparison: rFVIIa vs placebo, outcome: Mortality, latest timepoint

	Experim	ental	Contr	ol		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	Year	M-H, Fixed, 95% CI
1.1.1 Trauma (RCTs)								
Boffard 2005a (blunt)	17	69	22	74	16.1%	0.83 [0.48, 1.42]	2005	
Boffard 2005b (penetrating)	17	70	18	64	14.3%	0.86 [0.49, 1.53]	2005	
Hauser 2010a (blunt)	26	224	28	250	20.1%	1.04 [0.63, 1.71]	2010	
Hauser 2010b (penetrating)	8	46	5	40	4.1%	1.39 [0.49, 3.91]	2010	
Subtotal (95% CI)		409		428	54.6%	0.96 [0.71, 1.29]		+
Total events	68		73					
Heterogeneity: Chi# = 1.00, df = 3 (P = 0.80); I# =	0%							
Test for overall effect Z = 0.30 (P = 0.77)								
1.1.3 Medical emergency								
Bosch 2004 (GI haemorrhage in cirrhosis)	18	116	11	120	8.2%	1.50 [0.73, 3.10]	2004	
Bosch 2008 (GI haemorrhage in cirrhosis) (1)	39	170	25	86	25.3%	0.79 [0.51, 1.21]	2008	
Subtotal (95% CI)		286		206	33.5%	0.96 [0.67, 1.39]		+
Total events	55		36					
Heterogeneity: Chi2 = 2.29, df = 1 (P = 0.13); P =	56%							
Test for overall effect: Z = 0.19 (P = 0.05)								
1.1.4 Haemotology/Oncology								
Pihusch 2005 (haemonhage after HSCT)	24	77	7	23	8.2%	1.02 [0.51, 2.07]	2005	
Chuansumrit 2005 (dengue)	0	16	0	9		Not estimable	2005	
Subtotal (95% CI)		93		32	8.2%	1.02 [0.51, 2.07]		-
Total events	24		7					
Heterogeneity: Not applicable								
Test for overall effect $Z = 0.07 \ (P = 0.95)$								
1.1.5 Cardiac, adult								
Gill 2009 (treatment)	10	104	4	68	3.7%	1.63 [0.53, 5.00]	2009	
Subtotal (95% CI)		104		68	3.7%	1.63 [0.53, 5.00]		
Total events	10		4					
Heterogeneity: Not applicable								
Test for overall effect: Z = 0.06 (P = 0.39)								
1.1.6 Obstetrics								
Lavigne-Lissalde 2015 (PPH) Subtotal (95% CD	0	42	0	42		Not estimable		
Total quarte	0	42	0	16		not countable		
Heterneonethe Nationalicable	0		0					
Test for overall effect. Not applicable								
Total (95% CI)		934		776	100.0%	0.99 [0.80, 1.23]		•
Total events	157		120					S 8 9 9 9
Heterogeneity: Chi2 = 4.22, df = 7 (P = 0.75); P =	0%							
Test for overall effect: Z = 0.09 (P = 0.92)								0.05 0.2 1 5 20
Test for subgroup differences: Chi ² = 0.85, df = 3	P = 0.84	, P= 09	5					Favours (Fylia Favours standard Care
Footnates								

(1) Data for 600 and 300 ug/kg rFIIVa groups combined.

Table 4.36 Results for rFVIIa versus no rFVIIa: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
Any setting									
Simpson 2012 SR High quality	N = 2856 (13 RCTs)	Patients with critical bleeding due to trauma, or who had received treatment to manage bleeding	trauma, medical, oncology (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality, in- hospital, 28 or 30 day (including patients with spontaneous ICH)	332/1777 (18.7)	202/1079 (18.7)	RR 0.91 (0.78, 1.06)	No significant difference p = 0.2 No significant heterogeneity $l^2 = 0\% (p = 0.66)$
	N = 1569 (8 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Bosch 2004 Bosch 2008 Pihusch 2005 Chuansumrit 2005 Narayan 2008ª				Mortality, In- hospital, 28 or 30 day (excluding patients with spontaneous ICH)	154/849 (18.1) 17/6 17/7 26/22 8/4 16/1 39/17 24/7 0/ 7/	120/720 (16.7) 9 22/74 0 18/64 4 28/250 6 5/40 6 11/120 0 25/86 7 7/23 6 0/9 51 4/36	RR 0.95 (0.77, 1.18) RR 0.83 (0.48, 1.42) RR 0.86 (0.49, 1.53) RR 1.04 (0.63, 1.71) RR 1.39 (0.49, 3.91) RR 1.50 (0.73, 3.10) RR 0.79 (0.51, 1.21) RR 1.02 (0.51, 2.07) Not estimable RR 1.03 (0.32, 3.29)	No significant difference $p = 0.65^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.85)$
Trauma setting									
Cannon 2017 SR <i>Moderate quality</i>	N = 1292 (3 RCTs, 3 NRSIs) ^e	Patients with severe trauma at risk of death from haemorrhage	Trauma	rFVIIa vs no rFVIIa	Mortality, in- hospital, 28 or 30 day	112/517 (21.7)	237/775 (30.6)	OR 0.88 (0.64, 1.20)	No significant difference p = 0.42 No significant heterogeneity $l^2 = 15\% (p = 0.32)$
	N = 825 (3 RCTs) Boffard 2005a&b Hauser 2010					66/401 (16.5) 34/139 (24. 32/262 (12.	71/424 (16.7) 5) 40/144 (27.8) 2) 31/280 (11.1)	OR 0.97 (0.67, 1.41) OR 0.84 (0.49, 1.43) OR 1.12 (0.66, 1.89)	No significant difference p = 0.88 No heterogeneity $l^2 = 0\% (p = 0.46)$
Simpson 2012 SR High quality	N = 934 (4 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Hauser 2010 Narayan 2008 d	Adult patients with severe bleeding due blunt or penetrating trauma, or traumatic ICH	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality blunt penetrating blunt penetrating ICH	75/470 (16.0) 17/6 17/7 26/22 8/4 7/	77/464 (16.6) 9 22/74 0 18/64 4 28/250 6 5/40 11 4/36	RR 0.95 (0.71, 1.27) RR 0.83 (0.48, 1.42) RR 0.86 (0.49, 1.53) RR 1.04 (0.63, 1.71) RR 1.39 (0.49, 3.91) RR 1.03 (0.32, 3.29)	No significant difference $p = 0.73^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.91)$
Curry 2011 SR <i>Moderate quality</i>	N = 850 (3 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Hauser 2010	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality blunt penetrating blunt penetrating	NR/412 NR/6 NR/7 NR/22 NR/4	NR/438 9 NR/74 0 NR/64 6 NR/255 7 NR/45	Not calculated	No significant difference p = 0.58 p = 0.69 p = NR p = NR

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
Medical emergen	cy			- -					
Simpson 2012 SR High quality	N = 934 (3 RCTs) Bosch 2004 Bosch 2008 Chuansumrit 2005	Adult patients with cirrhosis and GI haemorrhage or children with dengue haemorrhagic fever	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality, not specified	55/302 16/116 39/170 0/16	36/215 11/120 25/86 0/9	RR 1.02 (0.55, 1.90) RR 1.50 (0.73, 3.10) RR 0.79 (0.51, 1.21) Not estimable	No significant difference $p = 0.95^{\circ}$ Substantial heterogeneity $l^2 = 56\% (p = 0.13)$
McQuilten 2015 SR	N = 510 (1 SR, κ=2 RCTs)	Adult patients with cirrhosis and Gl	Medical (multicentre,	rFVIIa (therapeutic)	Mortality, 5 days	NR	NR	RR 0.95 (0.36, 2.50)	No significant difference p = 0.16
Moderate quality	Marti-Caravajal 2012	haemorrhage	multicountry)	vs placebo	Mortality, 42 days	NR	NR	RR 1.01 (0.55, 1.87)	No significant difference p = 0.14
Oncology setting								1	
Simpson 2012 RCT High risk of bias	N = 934 (1 RCT) Pihusch 2005	Patients with moderate to severe bleeding after HSCT	Oncology (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality	24/77	7/23	1.02 (0.51, 2.07)	No significant difference p = 0.95° Heterogeneity NA
Surgical setting	1				1	1	1	1	
Yank 2011 RCT Unclear risk of bias	N = 172 (1 RCT) Gill 2009	Adult patients who had undergone cardiac surgery and were bleeding	Cardiology (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Mortality 40 ug/kg rFVIIa 80ug/kg rFVIIa	10/104 4/35 (11.4) 6/69 (8.7)	4/68 (5.8)	RD 0.04 (-0.04, 0.12)	No significant difference Heterogeneity NA p = NR
Obstetrics and ma	aternity setting								
Lavigne-Lissalde 2015 RCT High risk of blas	N = 84	Women (aged 18 years or older) with severe persistent primary PPH 'after sulprostone treatment	Obstetrics (multicentre, France, Switzerland)	rFVIIa (therapeutic) vs standard care	Mortality	0/42 (0)	0/42 (0)	Not estimable	Not estimable
Franchini 2010 SR (case series)	N = 272 (9 case series)	Women with severe PPH (≥ 500 mL after vaginal delivery and ≥ 1000 mL after caesarean delivery)	Obstetrics and gynaecology (multicountry including Europe and Australia)	rFVIIa	Mortality	The authors identified no studies, therefore attemp published case reports (N management of severe F Two retrospective Coh st discussed in PBM Module	RCTs, case-control o bted to extract useful J>10) to provide recon OPH. udies (Kalina 2011, Ho e 5 (technical report v	interventional cohort information from nmendations for the ssain 2007) identified and rol. 1 Section 3.4.4).	One study reported an association favouring rFVIIa after adjustment for confounders.

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Deadiatrias no a	omporativa avidan	a found							

Paediatrics - no comparative evidence found

Abbreviations: CI, confidence interval; GI, gastrointestinal; HSCT, haemopoietic stem cell transplant; ICH, intracerebral haemorrhage; NA, not applicable; NR, not reported; NRSIs, non-randomised study of intervention; OR, odds ratio; PPH, postpartum haemorrhage; RCTs, randomised controlled, trials; rFVIIa, recombinant activated factor seven; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational or cohort studies, the evidence has been considered as According to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H, random effects.

d. Narayan 2008 does not meet the PICO criteria for this review as it in a population with ICH (not critical bleeding with haemodynamic compromise).

e. NRSIs not included in the review for this question.

f. Defined as the loss of more than 1500 mL of blood within 24 hr after vaginal or caesarean delivery.

4.6.3.2 Morbidity

Thromboembolic events

A summary of the evidence relating to thromboembolic events in patients with critical bleeding treated with rFVIIa is presented in Table 4.37.

Overall, the evidence for harms (thromboembolic events) is limited. The studies were not large enough to detect important differences with variance for methods for detection of thromboembolic events also noted (*GRADE: very low*).

In a meta-analysis of data from included RCTs (Figure 4.20), there was slight increased risk of total thromboembolic events among patients administered rFVIIa (77/945; 8.1%) compared with placebo or no rFVIIa (58/780; 7.4%), however the difference was not statistically significant (RR 1.17, 95%CI 0.85, 1.63, p = 0.52, fixed effect, $I^2 = 0$ %).

Among patients with blunt and penetrating trauma who received rFVIIa, 10.8% (44/409) had a thromboembolic event compared with 10.0% (43/428) in the placebo group, corresponding to a nonsignificant difference between treatment groups (RR 1.10; 95% CI 0.74, 1.63; p = 0.63, fixed effect, $I^2 = 0$ %).

Among patients with UGIB, the rate of thromboembolic events in patients who received rFVIIa was also not significantly different from those who did not receive rFVIIa (5.4% vs 6.6%, RR 0.80; 95% CI 0.40, 1.60, p = 0.54, fixed effect, I² = 0%).

Among patients with uncontrolled bleeding after HSCT, the risk of thromboembolic events was higher in the group who received rFVIIa (8/93, 10.4%) compared with those who did not (0/23, 0%) (RR 5.23; 95% CI 0.31, 87.34; p = 0.25).

Among patients with uncontrolled bleeding due after cardiac surgery, the risk of thromboembolic events was higher in the group who received rFVIIa (7/104, 6.7%) compared with those who did not (1/68, 1.5%) (RR 4.58; 95% CI 0.58, 36.38; p = 0.15).

Among patients with PPH, the risk of thromboembolic events was higher in the group who received rFVIIa (2/42, 4.8%) compared with those who did not (0/42, 0%) (RR 5.00; 95% CI 0.25, 101.11; p = 0.29).

All identified systematic review suggested no increased risk of thromboembolic events among patients treated with rFVIIa, except one (Yank 2011), who suggested an increased (borderline) risk among patients with intractable bleeding after cardiac surgery. This is consistent with the review by Simpson 2012 and McQuilten 2015, who noted an increased risk of arterial thromboembolic events when both prophylactic and therapeutic studies were considered.

Figure 4.20 Forest plot of comparison: rFVIIa vs placebo, outcome: total thromboembolic events

	rFVII	a	standard	care		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	Year	M-H, Fixed, 95% CI
1.4.1 Trauma								
Boffard 2005a (blunt)	2	69	3	74	4.8%	0.71 [0.12, 4.15]	2005	
Boffard 2005b (penetrating)	4	70	3	64	5.2%	1.22 [0.28, 5.24]	2005	-
Hauser 2010a (blunt)	36	224	33	250	51.7%	1.22 [0.79, 1.88]	2010	
Hauser 2010b (penetrating)	2	46	4	40	7.1%	0.43 [0.08, 2.25]	2010	
Subtotal (95% CI)		409		428	68.9%	1.10 [0.74, 1.63]		•
Total events	44		43					
Heterogeneity: Chi ² = 1.68, df = 3 (P = 0.64); l ² = 0)%							
Test for overall effect: Z = 0.49 (P = 0.63)								
1.4.2 Medical emergency								
Bosch 2004 (GL baemorrhage in cirrhosis)	7	121	7	121	11.6%	1.00.00.36.2.761	2004	
Bosch 2008 (Gl haemorrhage in cirrhosis) (1)	9	176	. 7	89	15.4%	0.65 (0.25, 1.69)	2008	
Subtotal (95% CI)		297		210	27.0%	0.80 [0.40, 1.60]	2000	-
Total events	16		14					-
Heterogeneity: $Chi^2 = 0.37$ df = 1 (P = 0.54); $I^2 = 0$	196							
Test for overall effect: $Z = 0.63$ (P = 0.53)								
1.4.3 Oncology								
Chuansumrit 2005 (dengue)	0	16	0	9		Not estimable	2005	
Pihusch 2005 (haemorrhage after HSCT)	8	77	0	23	1.3%	5.23 [0.31, 87.34]	2005	
Subtotal (95% CI)		93		32	1.3%	5.23 [0.31, 87.34]		
Total events	8		0					
Heterogeneity: Not applicable								
Test for overall effect: Z = 1.15 (P = 0.25)								
1.4.5 Cardiac, adult	_							
Gill 2009 (treatment)	1	104	1	68	2.0%	4.58 [0.58, 36.38]	2009	
Subtotal (95% CI)	_	104		00	2.0%	4.56 [0.56, 50.56]		
i otal events			1					
Heterogeneity: Not applicable								
Test for overall effect: $Z = 1.44$ (P = 0.15)								
1.4.6 Obstetrics								
Lavigne-Lissalde 2015 (PPH)	2	42	0	42	0.8%	5.00 [0.25, 101.11]		
Subtotal (95% CI)		42		42	0.8%	5.00 [0.25, 101.11]		
Total events	2		0					
Heterogeneity: Not applicable								
Test for overall effect: Z = 1.05 (P = 0.29)								
Total (95% CI)		945		780	100.0%	1.17 [0.85, 1.63]		•
Total events	77		58					·
Heterogeneity: $Chi^2 = 6.94$ df = 8 (P = 0.54): $I^2 = 0$	196							++
Test for overall effect: $Z = 0.96$ (P = 0.33)								0.02 0.1 1 10 50
Test for subgroup differences: $Chi^2 = 4.78 df = 4$	(P = 0.31)) ²=1	6.3%					⊢avours r⊢VIIa ⊢avours standard care
Ecotoptos	v = 0.01		0.070					

Footnotes (1) Data for 600 and 300 ug/kg rFIIVa groups combined.

Table 4.37 Results for rFVIIa versus no rFVIIa: Patients with critical bleeding – Me	lorbidity: thromboembolic events
--	----------------------------------

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results						
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%) mean ± SD (n)	No rFVIIa n/N (%) mean ± SD (n)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b			
Any setting												
Simpson 2012 SR High quality	N = 2873 (13 RCTs)	Patients with critical bleeding due to trauma, or who had received treatment to manage bleeding	Trauma, medical, oncology (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events (including patients with spontaneous ICH)	169/1789	89/1084	RR 1.14 (0.89, 1.47)	No significant difference p = 0.30 No significant heterogeneity $l^2 = 0\% (p = 0.67)$			
	N = 1566 (9 RCTs) Boffard 2005a Boffard 2005b Hauser 2010a Hauser 2010b Bosch 2004 Bosch 2008 Pihusch 2005 Chuansumrit 2005 Narayan 2008 ^a	-			Total thromboembolic events (excluding patients with spontaneous ICH)	81/860 2/69 4/70 36/224 2/46 7/121 9/176 8/77 0/16 13/61	62/706 3/74 3/64 33/250 4/40 7/121 7/89 0/23 0/9 5/36	RR 1.10 (0.80, 1.52) 0.71 (0.12, 4.15) 1.22 (0.28, 5.24) 1.22 (0.79, 1.88) 0.43 (0.08, 2.25) 1.00 (0.36, 2.76) 0.65 (0.25, 1.69) 5.23 (0.31, 87.34) Not estimable 1.53 (0.60, 3.95)	No significant difference $p = 0.56^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.72)$			
McQuilten 2015 SR <i>Moderate quality</i>	N = 4119 (1 SR, κ=35 studies) Levi 2010	Off-label use in bleeding patients and healthy volunteers	Any	rFVIIa versus placebo	Total thromboembolic events Venous Cerebrovascular Arterial Coronary	NR	NR	OR 1.17 (0.94, 1.47) OR 0.93 (0.70, 1.23) OR 1.27 (0.74, 2.17) OR 1.68 (1.2, 2.36) OR 2.39 (1.39, 4.09)	No significant difference p = 0.16 p = 0.61 Not significant p = 0.39 Not significant p = 0.003 Favours rFVIIa p = 0.002 Favours rFVIIa			
Trauma setting												
Cannon 2017 SR High quality	N = 1061 (2 RCTs, 2 Coh) ^e	Patients with severe trauma at risk of death from haemorrhage	Trauma	rFVIIa vs no rFVIIa	Venous thromboembolic events	48/487 (9.9%)	57/574 (9.9%)	OR 0.97 (0.49, 1.92)	No significant difference p = 0.94 Mild heterogeneity $l^2 = 29\% (p = 0.24)$			
	N = 837 (2 RCTs) Boffard 2005 Hauser 2010					44/409 (10.8%) 6/139 (4.3%) 38/270 (14.1%)	43/428 (10.0%) 6/138 (4.3%) 37/290 (12.8%)	OR 1.10 (0.70, 1.72) OR 0.99 (0.31, 3.16) OR 1.12 (0.69, 1.82)	No significant difference p = 0.68 No heterogeneity $l^2 = 0\% (p = 0.85)$			

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%) mean ± SD (n)	No rFVIIa n/N (%) mean ± SD (n)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
McQuilten 2015 SR <i>Moderate</i>	N = 560 (1 RCT) Dutton 2011	Adult patients with blunt and/or penetrating trauma with continuing bleeding after receiving 4 units RBC despite standard haemostatic interventions	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Thromboembolic events Venous Arterial	25/270 (9) 16/270 (6)	26/287 (9) 12/290 (4)	NR NR	No significant difference p = 0.90 p = 0.33
Simpson 2012 SR High quality	N = 934 (4 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Hauser 2010 Nereysan 2000 r	Adult patients with severe bleeding due blunt or penetrating trauma, or traumatic ICH	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events	57/470 2/69 4/70 36/224 2/46 13/61	46/464 3/74 3/64 33/250 4/40 5/36	RR 1.17 (0.81, 1.69) RR 0.71 (0.12, 4.15) RR 1.22 (0.28, 5.24) RR 1.22 (0.79, 1.88) RR 0.43 (0.08, 2.25) RR 1.53 (0.60, 3.95)	No significant difference $p = 0.40^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.73)$
Medical emergend	cy								
McQuilten 2015 SR <i>Moderate</i>	N = 510 (1 SR, κ=2 RCTs) Marti-Caravajal 2012	Adult patients with cirrhosis and GI haemorrhage	Medical (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events	NR	NR	RR 0.80 (0.40, 1.60)	No significant difference p = 0.20
Simpson 2012 SR High quality	N = 532 (3 RCTs) Bosch 2004 Bosch 2008 Chuansumrit 2005	Adult patients with cirrhosis and GI haemorrhage or children with dengue haemorrhagic fever	Medical (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events	16/313 7/121 9/176 0/16	14/219 7/121 7/89 0/9	RR 0.80 (0.40, 1.60) RR 1.00 (0.36, 2.76) RR 0.65 (0.25, 1.69) Not estimable	No significant difference $p = 0.52^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.54)$
Oncology setting							·		
Simpson 2012 RCT High risk of bias	N = 100 (1 RCT) Pihusch 2005	Patients with moderate to severe bleeding after HSCT	Oncology trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events	8/77	0/23	RR 5.23 (0.31, 87.34)	No significant difference p = 0.25 Heterogeneity NA
Surgical setting									
Yank 2011 RCT High risk of bias	N = 172 (1 RCT) Gill 2009	Adult patients who had undergone cardiac surgery and were bleeding	Surgical (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Total thromboembolic events 40 ug/kg rFVIIa 80ug/kg rFVIIa	7/104 (6.7) 3/35 (8.6) 4/69 (5.8)	1/68 (1.5)	RD 0.05 (0.00, 0.11)	Favours rFVIIa p = NR (borderline)

Study ID	Sample size	Patient	Setting	Setting Comparison C		Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%) mean ± SD (n)	No rFVIIa n/N (%) mean ± SD (n)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Obstetrics and ma	nternity setting									
Lavigne-Lissalde 2015 RCT High risk of bias	N = 84	Women (aged 18 years or older) with severe persistent primary PPH ^r after sulprostone treatment	Obstetrics (multicentre, France, Switzerland)	rFVIIa (therapeutic) vs standard care	Thromboembolic events	2/42 (5)	0/42 (0)	NR	<i>No significant difference</i> <i>p</i> = 0.25	
Franchini 2010 SR (case series) <i>High risk of bias</i>	N = 272 (9 case series)	Women with severe PPH (≥ 500 mL after vaginal delivery and ≥ 1000 mL after caesarean delivery)	Obstetrics and gynaecology (various countries including Europe and Australia)	rFVIIa vs no rFVIIa	Thromboembolic events	The authors identifi studies, therefore at published case report the management o Three retrospective Hossain 2007) ident Section 3.4.4). One T	ed no RCTs, case-cont tempted to extract us rts (total N>10) to pro f severe PPH. cohort studies (Kalina ified and discussed in 'E event reported.	rol or interventional cohort seful information from vide recommendations for a 2011, Ahonen 2007, PBM Module 5 (TR vol. 1	No significant association between treatment with rFVIIa and TE observed.	

Paediatrics - no comparative evidence found

Abbreviations: CI, confidence interval; GI, gastrointestinal; HSCT, haemopoietic stem cell transplant; NR, not reported; OR, odds ratio; PPH, postpartum haemorrhage; RCTs, randomised controlled, trials; rFVIIa, recombinant activated factor seven; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational or cohort studies, the evidence has been considered according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

d. Narayan 2008 does not meet the PICO criteria for this review as it in a population with ICH (not critical bleeding with haemodynamic compromise).

e. NRSIs not included in the review for this question.

f. Defined as the loss of more than 1500 mL of blood within 24 hr after vaginal or caesarean delivery.

Other adverse events

A summary of the evidence relating to other adverse events in patients with critical bleeding treated with rFVIIa is presented in Table 4.38.

The available evidence suggested a slight increased benefit associated with a reduced incidence of acute respiratory distress syndrome (ARDS) and multiorgan failure (MOF) among patients with blunt or penetrating trauma, however it is noted that the evidence is weak and limited to post-hoc analyses (*GRADE: low*).

In a meta-analysis of data from RCTs included in this review (Figure 4.21), fewer patients with blunt and penetrating trauma who received rFVIIa were reported to have ARDS compared with those who received placebo (3.4% vs 8.9%); an effect that was statistically significant (RR 0.39, 95%CI 0.22, 0.71, p = 0.002, fixed effect, $I^2 = 0$ %). Similarly, significantly fewer patients who received rFVIIa were reported to have MOF compared with the placebo group (4.4% vs 7.9%; RR 0.56; 95% CI 0.32, 0.97; p = 0.004, fixed effect, $I^2 = 0$ %).

Evidence for ARDS or MOF was not reported in the RCTs evaluating the effects of rFVIIa in patients with UGIB or those with uncontrolled bleeding due to other medical conditions (after HSCT, dengue haemorrhagic fever, cardiac surgery, primary PPH).

One RCT (Hauser 2010) examining the effects of rFVIIa among patients with blunt or penetrating trauma also reported on the incidence of sepsis and disseminated intravascular coagulation among treated patients. For both outcomes, a non-significant difference between the treatment groups was observed (RR 0.86; 95% CI 0.58, 1.28; p = 0.47, fixed effect, I² = 0% and RR 0.69; 95% Ci 0.27, 1.76; p = 0.44, fixed effect, I² = 0%, respectively).

Evidence from the included systematic reviews suggested a slight increased benefit among patients treated with rFVIIa associated with a reduced incidence of acute respiratory distress syndrome (ARDS) and multiorgan failure (MOF) among patients with blunt or penetrating trauma, however it was noted that the evidence is weak and limited to post-hoc analyses.

Figure 4.21 Forest plot of comparison: rFVIIa vs placebo, outcome: Other adverse events (trauma setting)

	rFVIIa	1	Standard	care		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% Cl	M-H, Fixed, 95% CI
1.5.1 Acute respiratory distre	ss syndro	ome					
Boffard 2005a (blunt)	2	69	12	74	30.8%	0.18 [0.04, 0.77]	_
Boffard 2005b (penetrating)	4	70	5	64	13.9%	0.73 [0.21, 2.61]	
Hauser 2010a (blunt)	8	224	18	250	45.3%	0.50 [0.22, 1.12]	
Hauser 2010b (penetrating)	0	46	3	40	10.0%	0.12 [0.01, 2.34]	← ■
Subtotal (95% CI)		409		428	100.0%	0.39 [0.22, 0.71]	◆
Total events	14		38				
Heterogeneity: Chi ² = 2.94, df =	= 3 (P = 0.4	40); l² :	= 0%				
Test for overall effect: Z = 3.10	(P = 0.002)	2)					
1.5.2 Multiorgan failure							
Boffard 2005a (blunt)	5	69	9	74	26.2%	0.60 [0.21, 1.69]	
Boffard 2005b (penetrating)	2	70	7	64	22.1%	0.26 [0.06, 1.21]	
Hauser 2010a (blunt)	10	224	17	250	48.5%	0.66 [0.31, 1.40]	
Hauser 2010b (penetrating)	1	46	1	40	3.2%	0.87 [0.06, 13.46]	
Subtotal (95% CI)		409		428	100.0%	0.56 [0.32, 0.97]	\bullet
Total events	18		34				
Heterogeneity: Chi ² = 1.23, df =	= 3 (P = 0.1	75); I ² :	= 0%				
Test for overall effect: Z = 2.06	(P = 0.04)						
4.E.2. Consis							
1.5.5 Sepsis							
Hauser 2010a (blunt)	33	224	45	250	95.2%	0.82 [0.54, 1.23]	
Hauser 2010b (penetrating)	4	270	2	40	4.8%	1.74 [0.34, 9.00]	
Subtotal (95% CI)	07	210	47	290	100.0%	0.00 [0.30, 1.20]	
Lotaregeneity ObiZ = 0.76 df-	- 1 (D - 0 '	- 51 - 40 -	- 00/				
Teet for everall effect: 7 = 0.76	= 1 (P = 0 7D = 0.47)	38), 179	= 0%				
Test for overall effect. $\Sigma = 0.73$	(P = 0.47)						
1.5.4 DIC							
Hauser 2010a (blunt)	6	224	10	250	89.8%	0.67 [0.25, 1.81]	
Hauser 2010b (penetrating)	1	46	1	40	10.2%	0.87 [0.06, 13.46]	
Subtotal (95% CI)		270		290	100.0%	0.69 [0.27, 1.76]	
Total events	7		11				
Heterogeneity: Chi ² = 0.03, df =	= 1 (P = 0.0	86); I ž =	= 0%				
Test for overall effect: Z = 0.78	(P = 0.44)						
							Eavours rEVIIa Eavours standard care
Toot for outgroup differences:	ONIZ - E O	11 df -	2/D = 0.4	73 12 - 4	0.40/		. arouro n' ma l'arouro otanidata care

Test for subgroup differences: Chi² = 5.01, df = 3 (P = 0.17), l² = 40.1 \%

Table 4.38	Results for rFVIIa versus no rFVIIa: Patients with critical bleeding	g – Morbidity: other adverse events
------------	--	-------------------------------------

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting				·						
Updated in this review *data retrieved from primary studies	N = 961 (3 RCTs) Boffard 2005a Boffard 2005b Hauser 2010	Adult patients with blunt and/or penetrating trauma with critical bleeding	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	ARDS, 30 or 90 days blunt penetrating blunt penetrating MOF, 30 or 90 days blunt penetrating blunt	14/409 (3.4) 3/69 (4) 4/70 (5) 8/224 (3.6) 0/46 (0.0) 18/409 (4.4) 5/69 (7) 2/70 (3) 10/224 (4.5)	38/428 (8.9) 12/74 (16) 5/64 (8) 18/250 (7.2) 3/40 (7.5) 34/428 (7.9) 9/74 (12) 7/64 (11) 17/250 (6.8)	RR 0.39 (0.22, 0.71) 0.18 (0.04, 0.77) 0.73 (0.21, 2.61) 0.5 (0.22, 1.12) 0.12 (0.01, 2.34) RR 0.56 (0.32, 0.97) RR 0.6 (0.21, 1.69) 0.26 0.06, 1.21) 0.66 (0.31, 1.40)	Favours rFVIIa $p = 0.002^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.4)$ No significant difference $p = 0.04^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.75)$	
	N = 560 (1 RCT) Hauser 2010					Sepsis, 90 days blunt	37/270 33/224 (14.7) 4/46 (8.7)	47/290 45/250 (18) 2/40 (5.0)	RR 0.86 (0.57, 1.28) RR 0.82 (0.54, 1.23) RR 1.74 (0.34, 9.00)	No significant difference $p = 0.44^{\circ}$ No significant subgroup heterogeneity $l^2 = 0\% (p = 0.38)$
					DIC, 90 days blunt penetrating	7/270 6/224 (2.7) 1/46 (2.2)	11/290 10/250 (4.0) 1/40 (2.5)	RR 0.69 (0.27, 1.76) RR 0.67 (0.25, 1.81) RR 0.87 (0.06, 13.46)	No significant difference p = 0.44 ^c No significant subgroup heterogeneity $l^2 = 0\%$ ($p = 0.86$)	
	N = 220 (2 RCTs) Boffard 2009	Patients who did not survive the first 48 hours excluded	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	ARDS, 30 days blunt trauma penetrating trauma	11/NR (5) NR	3/NR (18) NR	NR NR	Favours rFVIIa p = 0.05 (blunt) p = NR	
						Logistic regression demographic and confirmed reduced p = 0.0163 in patien Data on patients w	analysis (See Boffard baseline variables kr d risk favouring rFVII nts with blunt traum ith penetrating trau	d 2009) adjusted for own to predict ARDS a (OR 0.16; 95% CI 0.02, 0.73; a. ma NR.		
					MOF, 30 day blunt trauma penetrating trauma	7/NR (11) NR	3/NR (5) NR	NR NR	No significant difference p = 0.30 (blunt) p = NR	
					Logistic regression baseline variables	analysis adjusted fo known to predict MC	demographic and DS suggested a significant			

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results										
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b							
						difference in favou 0.0406). (See Boffa Data on patients w	r of rFVIIa (OR 0.05: 9 rd 2009) ⁄ith penetrating trau	5% CI 0.00, 0.89; <i>p</i> = ma NR.								
McQuilten 2015 SR	N = 560 (1 RCT) Dutton 2011	Adult patients with blunt and/or	Trauma (multicentre,	rFVIIa (therapeutic) vs	ARDS, 30 days	8/270 (3)	21/290 (7.2)	NR	Favours rFVIIa p = 0.02							
Moderate quality	N = 573 (1 RCT) Hauser 2010	penetrating trauma with continued bleeding after receipt of 4U RBC despite standard haemostatic interventions	multicountry)	placebo	MOF, 30 day blunt and penetrating blunt	NR/267 (23) NR/221 (45)	NR/287 (24) NR/247 (53)	NR NR	No significant difference p = 0.09 p = 0.06							
Curry 2011 SR <i>Moderate quality</i>	N = 850 (3 RCTs) Boffard 2005a Boffard 2005b Hauser 2010	Adults patients with blunt or penetrating injury	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	ARDS, not specified blunt penetrating blunt & penetrating	NR/411 NR/69 NR/70 NR/272	NR/438 NR/74 NR/64 NR/300	NR NR NR NR	p = NR Favours rFVIIa p = NR no difference p = NR							
			-		Post-hoc analys	ses of Boffard 200	5 a&b	trend towards (blunt only)								
						Rizoli 2006 (coagulopathic patients)	NR/60	NR/76	NR	Favours rFVIIa p = NR						
					Boffard 2009 (patients surviving 48 hours or more)	NR/139	NR/138	OR 0.16 (0.02, 0.73)	<i>Favours rFVIIa</i> p = NR							
					MOF, not specified blunt penetrating blunt & penetrating	NR/411 NR/69 NR/70 NR/272	NR/438 NR/74 NR/64 NR/300	NR	p = NR no difference p = NR trend towards p = NR trend towards (blunt only)							
						Post-hoc analys	es of Boffard 200	5 a&b								
					Rizoli 2006 (coagulopathic patients)	NR/60	NR/76	NR	p = NR trend towards							
												Boffard 2009 (patients surviving 48 hours or more)	NR/69 (blunt) NR/70 (penetrating)	NR/74 NR/64	OR 0.05 (0.0, 0.89) NR	Favours rFVIIa (blunt injury) p = NR
					MOF and ARDS Boffard 2009 (patients surviving 48 hours or more)	NR/139	NR/138	OR 0.16 (0.02, 0.81)	Favours rFVIIa (blunt injury) p = NR							
Yank 2011	N = 277 (2 RCTs)				ARDS											

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%)	No rFVIIa n/N (%)	Risk estimate (95% Cl)	Statistical significance		
									<i>p</i> -value		
									Heterogeneity ^b		
SR	Boffard 2005a				blunt	NR/69 (4.3)	NR/74 (16.2)	NR	p = 0.03 Favours rFVIIa		
High quality	Boffard 2005b				penetrating	NR/70 (5.7)	NR/64 (7.8)	NR	p = 0.74 no difference		
Obstetrics and maternity setting - no comparative evidence found											

Paediatrics - no comparative evidence found

Abbreviations: CI, confidence interval; DIC, disseminated intravascular coagulation; GI, gastrointestinal; HSCT, haemopoietic stem cell transplant; MODS, multiorgan dysfunction syndrome; NR, not reported; OR, odds ratio; RCTs, randomised controlled, trials; rFVIIa, recombinant activated factor seven; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H, random effects.

Need for second-line therapies

A summary of the evidence relating to other second-line therapies in the obstetrics and maternity setting in patients with critical bleeding treated with rFVIIa is presented in Table 4.37. This outcome was recognised as a critical patient relevant outcome of interest in this setting only.

Among women with severe PPH with persistent bleeding after sulprostone treatment, the use of rFVIIa was reported to reduce the incidence of second-line therapies compared with standard care (RR 0.56; 95% CI 0.42, 0.76; p = 0.0002); however, the data is limited by low patient numbers. *(GRADE: very low)*

Specifically, there was a reduced need for arterial embolisation (see Figure 4.23).

Figure 4.22 Forest plot of comparison: rFVIIa vs placebo, outcome: Morbidity – need for second-line intervention (obstetrics and maternity)

	rFVI	a	Standard	care		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% Cl		M-H, Fixed, 95% CI	
1.4.5 Need for other second line	e therapie	es							
Lavigne-Lissalde 2015 (PPH) Subtotal (95% CI)	22	42 42	39	42 42	100.0% 100.0%	0.56 [0.42, 0.76] 0.56 [0.42, 0.76]		•	
Total events Heterogeneity: Not applicable	22		39						
Test for overall effect: Z = 3.74 (P	= 0.0002	2)							
Total (95% CI)		42		42	100.0%	0.56 [0.42, 0.76]		◆	
Total events	22		39						
Heterogeneity: Not applicable									100
Test for overall effect: Z = 3.74 (P					0.01	Eavours rEVIIa Eavours standard ca	100		
Test for subgroup differences: N	ot applica	able							10

Figure 4.23 Forest plot of comparison: rFVIIa vs placebo, outcome: Morbidity - other secondline interventions (obstetrics and maternity)

	rFVII	a	Standard	care		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 95% CI
1.3.1 Arterial embolization							_
Lavigne-Lissalde 2015 (PPH) Subtotal (95% CI)	12	42 42	24	42 42	100.0% 100.0%	0.50 [0.29, 0.86] 0.50 [0.29, 0.86]	
Total events	12		24				
Heterogeneity: Not applicable							
Test for overall effect: Z = 2.49 (F	° = 0.01)						
1.3.2 Arterial ligation							_
Lavigne-Lissalde 2015 (PPH) Subtotal (95% CI)	9	42 42	12	42 42	100.0% 100.0%	0.75 [0.35, 1.59] 0.75 [0.35, 1.59]	
Total events	9		12				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.75 (F	° = 0.45)						
1.3.3 Peripartum hysterectomy	1						
Lavigne-Lissalde 2015 (PPH)	3	42	8	42	100.0%	0.38 [0.11, 1.32]	
Subtotal (95% CI)		42		42	100.0%	0.38 [0.11, 1.32]	
l otal events	3		8				
Heterogeneity: Not applicable	0.400						
Test for overall effect. $Z = 1.53$ (F	-= 0.13)						
1.3.4 Other (B-Lynch, Bakri Bal	loon, etc)						_
Lavigne-Lissalde 2015 (PPH)	4	42	6	42	100.0%	0.67 [0.20, 2.19]	
Subtotal (95% CI)		42		42	100.0%	0.67 [0.20, 2.19]	
Total events	4		6				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.67$ (F	P = 0.50)						
							0.2 0.5 1 2 5
Test for subgroup differences: C	>hi² = 1.22	2. df = 3	(P = 0.75)	. I² = 0%			Favours rFVIIa Favours standard care

Table 4.39 Results for rFVIIa versus no rFVIIa: Patients with critical bleeding – Morbidity: other second-line interventions (obstetrics and maternity only)

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ª	(no. of trials) included in analysis	population	(Location)			rFVIIa n/N (%) mean ± SD (n)	No rFVIIa n/N (%) mean ± SD (n)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Obstetrics and maternity setting									
Lavigne-	N = 84	Women (aged 18	Obstetrics	rFVIIa	Reduction in the need for	22/42	39/42 (93)	RR 0.56 (0.42, 0.76)	Favours rFVIIa
Lissalde 2015		years or older) with	(multicentre,	(therapeutic) vs	specific second-line				<i>p</i> < 0.0001
RCT		severe persistent	France,	standard care	therapies (composite)				
High risk of bias		primary PPH ^c after	Switzerland)		Arterial embolisation	12/42 (29)	24/42 (57)	RR 0.50 (0.29, 0.86)	p = 0.0082
_		sulprostone			Arterial ligation	9/42 (21)	12/42 (29)	RR 0.75 (0.35, 1.59)	p = 0.45
		treatment			Peripartum hysterectomy	3/42 (7)	8/42 (19)	RR 0.38 (0.11, 1.32)	p = 0.11
					Other (B-lynch, Bakri Balloon etc.)	4/42 (10)	6/42 (14)	RR 0.67 (0.20, 2.19)	p = 0.50

Abbreviations: CI, confidence interval; GI, gastrointestinal; HSCT, haemopoietic stem cell transplant; NR, not reported; OR, odds ratio; PPH, postpartum haemorrhage; RCTs, randomised controlled, trials; rFVIIa, recombinant activated factor seven; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as According to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Defined as the loss of more than 1500 mL of blood within 24 hr after vaginal or caesarean delivery.

4.6.3.3 Transfusion volumes

Red blood cells

A summary of the evidence relating to RBC transfusion volumes in patients with critical bleeding treated with rFVIIa is presented in Table 4.40.

Only limited conclusions can be drawn from the available evidence, with an overall modest reduction in the volume of RBC transfused (less than one red cell unit saved). The Cochrane review by Simpson 2012 noted that these favourable findings were likely overestimated because data were not available from larger negative studies for inclusion in the meta-analysis.

A meta-analysis of data from RCTs included in this review (see Figure 4.24) revealed a reduction in the volume of RBC transfusion in patients with critical bleeding who received rFVIIa (n=552) compared with those who did not (n=579), with an overall mean difference (MD) of –0.90 units observed (95% CI –1.82, 0.02; p = 0.05; random effect, I² = 58%).

There was a large difference among the subgroups, with the RCTs conducted in bleeding patients with blunt or penetrating trauma suggesting a reduction in the volume of RBC transfusion to be closer 2 units saved (MD –2.35; 95% CI –3.70, –1.00; p = 0.0007) (*GRADE: very low*). It was noted that these data are confounded by the exclusion of trauma patients who died within 48 hours of admission to hospital.

Among patients with UGIB who received rFVIIa there was no difference in RBC transfusion volumes between treatment groups (MD –0.24, 95% CI –1.17, 0.69; p = 0.61, I² = 62%) (*GRADE: very low*). A similar result was observed in paediatric patients with dengue haemorrhagic fever (MD 0.10, 95% CI –1.24, 1.44; p = 0.88) (*GRADE: very low*).

The volume of RBC transfused was not reported in the RCTs conducted in patients with intractable bleeding after cardiac surgery or in women with severe PPH with persistent bleeding after sulprostone treatment.

Figure 4.24 Forest plot of comparison: rFVIIa vs placebo, outcome: RBC transfusion volume, Units

	Expe	erimen	tal	C	ontrol			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.7.1 Trauma (from dosing to 48 hour)									
Boffard 2005a (blunt) (1)	6.9	6.2	52	10.9	9.3	59	7.5%	-4.00 [-6.91, -1.09]	+
Boffard 2005b (penetrating) (2)	4.5	5.3	57	7.7	9.9	52	7.1%	-3.20 [-6.22, -0.18]	← <u></u>
lauser 2010a (blunt)	7.8	10.6	191	9.1	11.3	228	11.7%	-1.30 [-3.40, 0.80]	
Hauser 2010b (penetrating) Subtotal (95% CI)	5	7.4	39 339	6.8	6.9	35 374	6.3% 32.7%	-1.80 [-5.06, 1.46] -2.35 [-3.70, -1.00]	•
leterogeneity: Tau ^a = 0.00; Chi ^a = 2.61, df = 3 (f	P = 0.46	$ ^{2} = 0^{4}$	36						
Fest for overall effect Z = 3.41 (P = 0.0007)									
1.7.2 Medical emergency									
3osch 2004 (GI haemorrhage in cirrhosis)	1.5	3.7	121	1.3	1.9	121	25.3%	0.20 [-0.54, 0.94]	
Bosch 2008 (GI haemorrhage in cirrhosis) (3) Subtotal (95% CI)	2.55	2.4	76	3.3	3.1	75	23.6% 48.9%	-0.75 [-1.64, 0.14] -0.24 [-1.17, 0.69]	
-leterogeneity: Tau ^a = 0.28; Chi ^a = 2.60, df = 1 (f Fest for overall effect: Z = 0.51 (P = 0.61)	P = 0.11)	I ² = 6.	2%						
1.7.3 Haematology/oncology									
Chuansumril 2005 (dengue) Subtotal (95% CI)	0.44	2.7	16	0.34	0.34	9	18.4%	0.10 [-1.24, 1.44]	-
Heterogeneity. Not applicable									
Fest for overall effect: Z = 0.15 (P = 0.88)									
fotal (95% CI)			552			579	100.0%	0.90 [-1.82, 0.02]	-
Heterogeneity: Tau* = 0.73; Chi* = 14.22, df = 6	(P = 0.03)	$0; 1^{2} = 0$	58%						
fest for overall effect: Z = 1.92 (P = 0.05)	2010/01/04	2011/08							-4 -2 U 2 4
Fest for subgroup differences: Chi ² = 8.00, df =	2 (P = 0.)	02), P :	75.09	6					Favours (Fivial Favours pracedo
Footnotes									
1) excludes patients who died within 48 hours.									

(1) excludes patients who died within 48 hours (2) excludes patients who died within 48 hours (3) Data for 600 and 300 ug/kg rFIIVa groups combined (average).

HTANALYSTS | National Blood Authority | Critical bleeding | Technical report Volume 1

Table 4.40 Results for rFVIIa versus no rFVIIa: Patients with critical bleeding – RBC transfusion volume

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results			
Study design ª	(no. of trials) included in analysis		(Location)			rFVIIa mean ± SD (n)	No rFVIIa mean ± SD (n)	Mean Difference (95% CI)	Statistical significance p-value Heterogeneity ^b
Any setting	1		1	1	1		1	1	
Simpson 2012 SR High quality	N = 911 (4 RCTs) Hauser 2010 Bosch 2004 Bosch 2008 Chuansumrit 2005	Patients with critical bleeding due to trauma, or who had received treatment to manage bleeding	Trauma, medical, oncology (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion volume, mL blunt penetrating Gl haemorrhage Gl haemorrhage dengue	(n = 443) 2340 ± 3180 (191) 1500 ± 2220 (39) 450 ± 1110 (121) 764 ± 719 (76) 131 ± 812 (16)	(n = 468) 2730 ± 3390 (228) 2040 ± 2070 (35) 390 ± 570 (121) 990 ± 930 (75) 103 ± 102 (9)	MD -88.60 (-263.88, 86.68) -390.00 (-1020.09, 240.09) -540.00 (-1517.62, 437.62) 60.00 (-162.33, 282.33) -226.00 (-491.39, 39.39) 28.00 (-375.41, 431.41)	No significant difference p = 0.32 Mild heterogeneity $I^2 = 16\% (p = 0.32)$
Trauma settin	g								
Cannon 2017 SR High quality	N = 933 (3 RCTs, 2 NRSIs) ^c	Patients with severe trauma at risk of death from haemorrhage	Trauma	rFVIIa vs placebo or no rFVIIa	RBC transfusion volume, units to 24 hours	(n = 424)	(n = 509)	MD -0 .92 (-2.31, 0.47)	No significant difference p = 0.19 No significant heterogeneity $l^2 = 17\%$ ($p = 0.30$)
	N = 742 (3 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Hauser 2010				blunt penetrating blunt penetrating	(n = 354) 7.8 ± 12 (64) 4 ± 9.25 (69) 6.9 ± 10.4 (184) 4.5 ± 7.3 (37)	(n = 388) 7.2 ± 8.75 (72) 4.8 ± 10.25 (61) 8.1 ± 10.9 (222) 6.2 ± 6.5 (33)	MD -0.94 (-2.36, 0.48) MD 0.60 (-2.97, 4.17) MD -0.80 (-4.17, 2.57) MD -1.20 (-3.28, 0.88) MD -1.70 (-4.93, 1.53)	No significant difference p = 0.20 No heterogeneity $l^2 = 0\% (p = 0.80)$
McQuilten 2015 SR <i>Moderate</i> <i>quality</i>	N = 554 (1 RCT) Hauser 2010 N = 277 (2 RCTs)	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion, units to 24 hrs blunt penetrating RBC transfusion, units to 48 hrs	(n = 221) 6.9 ± 10.4 (184) 4.5 ± 7.3 (37)	(n = 255) 8.1 ± 10.9 (222) 6.2 ± 6.5 (33)	MD -1.35 (-3.09, 0.40)	No significant difference p = 0.13 ^d p = 0.04 Favours rFVIla p = 0.11 No difference No significant difference
*additional data sought from primary studies	Boffard 2005b				blunt	NR (69) NR (70) Estimated reduction. H transfusion amount fro who died within 48 ho	NR (74) NR (64) łodges-Lehmann point om placebo to active gro urs were assigned the h	Est. 2.0 (0.0, 4.6) Est. 0.2 (-0.9, 2.4) estimate of the shift in oup, including 90% CI. Patients ighest rank (see Boffard 2009). ^d	<i>p</i> = 0.07 <i>p</i> = 0.24
Simpson 2012 SR High quality	N = 493 (1 RCT) Hauser 2010	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion volume, mL blunt penetrating	(n = 230) 2340 ± 3180 (191) 1500 ± 2220 (39)	(n = 263) 2730 ± 3390 (228) 2040 ± 2070 (35)	MD -434.02 (-963.64, 95.59) -390.00 (-1020.09, 240.09) -540.00 (-1517.62, 437.62)	No significant difference $p = 0.11^{d}$ No significant heterogeneity $l^2 = 0\% (p = 0.80)$

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results			
Study design ª	(no. of trials) included in analysis		(Location)			rFVIIa mean ± SD (n)	No rFVIIa mean ± SD (n)	Mean Difference (95% CI)	Statistical significance p-value Heterogeneity ^b
Yank 2011 SR High quality	N = 220 (2 RCTs) Boffard 2005a Boffard 2005b	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion, units (from dosing to 48 hrs) blunt penetrating (excludes patients who died within 48 hours)	(n = 109) 6.9 ± 6.2 4.5 ± 5.3 (57)	(n = 111) 10.9 ± 9.3 (59) 7.7 ± 9.9	NR NR	p = 0.02 Favours rFVIIa p = 0.10 No difference
Curry 2011 SR <i>Moderate</i> <i>quality</i>	N = 850 (3 RCTs) Boffard 2005a Boffard 2005b Hauser 2010 Hauser 2010	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion volume blunt penetrating blunt penetrating	(n = 412) NR (69) NR (70) NR (226) NR (46)	(n = 438) NR (74) NR (64) NR (255) NR (45)	NR	Favours rFVIIa (blunt) trend towards (penetrating)
					Rizoli 2006 coagulopathic patients	Post-noc analyses ((n = 60)	of Boffard 2005a&b (n = 76) NR	NR	<i>Favours rFVIIa</i> <i>p</i> = 0.02
Medical emerg	jency								
Simpson 2012 SR High quality	N = 418 (3 RCTs) Bosch 2004 Bosch 2008 Chuansumrit 2005	Adult patients with cirrhosis and GI haemorrhage or children with dengue haemorrhagic fever	Medical trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	RBC transfusion volume, mL	(n = 213) 450 ± 1110 (121) 764 ± 719 (76) 131 ± 812 (16)	(n = 205) 390 ± 570 (121) 990 ± 930 (75) 103 ± 102 (9)	MD -48.74 (-239.88, 142.40) 60.00 (-162.33, 282.33) -226.00 (-491.39, 39.39) 28.00 (-375.41, 431.41)	No significant difference $p = 0.62^{\text{ d}}$ Moderate heterogeneity $l^2 = 28\% (p = 0.62)$
Surgical settin	g - no comparati	ve evidence found						1	
Obstetrics and	l maternity settin	g							
Lavigne- Lissalde 2015 RCT High risk of bias	N = 84	Women (aged 18 years or older) with severe persistent primary PPH ^e after sulprostone treatment	Obstetrics (multicentre, France, Switzerland)	rFVIIa (therapeutic) vs standard care	RBC transfusion volume, units	(n = 42) median (IQR) 2 (0, 3)	(n = 42) median (IQR) 2 (0, 4)	0 (NR)	No significant difference p = NR
Franchini 2010 SR (case series) High risk of bias	N = 272 (9 case series)	Women with severe PPH (≥ 500 mL after vaginal delivery and ≥ 1000 mL after caesarean delivery)	Obstetrics and gynaecology (multicountry, including Europe and Australia)	rFVIIa (therapeutic) vs placebo or no rFVIIa	Transfusion volume	The authors identified therefore attempted to reports (N >10) to provi PPH. Three retrospective co identified and discusse	no RCTs, case-control o o extract useful informal de recommendations fo hort studies (Ahonen 20 d in PBM Module 5 (TR	interventional cohort studies, tion from published case or the management of severe 007, Hossain 2007, Kalina 2011) vol. 1 Section 3.4.4).	Women who received rFVIIa were given more RBC than the comparator groups, but those women had more severe haemorrhaging.

Study ID	dy ID Sample size Patient	Patient population	Setting	Comparison	Outcome	Results					
Study design ª	(no. of trials) included in analysis		(Location)			rFVIIa mean ± SD (n)	No rFVIIa mean ± SD (n)	Mean Difference (95% CI)	Statistical significance p-value Heterogeneity ^b		

Paediatrics - no comparative evidence found

Abbreviations: CI, confidence interval; hrs, hours; IU, international units; M-H, Mantzel-Hentzel; NR, not reported; OR, odds ratio; PPH, postpartum haemorrhage; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. NRSIs not included in the review for this question.

d. Calculated post-hoc using RevMan 5.4. M-H random effects.

d. Exclusion of patients who died within 48 hours shows a significant reduction in total RBC transfusions in 48 hours among patients with blunt trauma (estimated reduction 2.6 units; 90% CI 0.7, 4.6; *p* = 0.02) but not patients with penetrating trauma (1.0 unit; 90% CI 0.0, 2.6; *p* = 0.10). (see Boffard 2009)

e. Defined as the loss of more than 1500 mL of blood within 24 hr after vaginal or caesarean delivery

Other blood components

A summary of the evidence relating to transfusion volumes of other blood components in patients with critical bleeding treated with rFVIIa is presented in Table 4.41.

A meta-analysis of usable data from RCTs included in this review (see Figure 4.25) revealed a significant (borderline) reduction in the transfusion of allogenic blood components at 24 hours in trauma patients with critical bleeding who received rFVIIa (n=237) compared with those who did not (n=263), with an overall MD of -4.17 units observed (95% CI -8.40, 0.07; p = 0.05; fixed effect, I² = 0%). This effect was significant for FFP (MD -2.14; 95% CI -3.54, -0.73; p = 0.003) but not platelets, fibrinogen concentrate or cryoprecipitate.

Data for patients with UGIB, paediatric patients with dengue haemorrhagic fever, patients with intractable bleeding after cardiac surgery or in women with severe PPH with persistent bleeding after sulprostone treatment were not able to be assessed.

Figure 4.25 Forest plot of comparison: rFVIIa vs placebo, outcome: transfusion volume (other blood components), Units

	Expe	rimen	tal	C	ontrol			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Fixed, 95% CI	IV, Fixed, 95% Cl
1.9.1 Allogenic									
Hauser 2010a (blunt)	17.1	26.8	198	20.7	25.7	228	71.6%	-3.60 [-8.61, 1.41]	←
Hauser 2010b (penetrating)	11.2	15	39	16.8	19.3	35	28.4%	-5.60 [-13.54, 2.34]	← ■
Subtotal (95% CI)			237			263	100.0%	-4.17 [-8.40, 0.07]	
Heterogeneity: Chi ² = 0.17, df =	: 1 (P = 0	0.68); I	²=0%						
Test for overall effect: Z = 1.93	(P = 0.0)	5)							
1.9.2 FFP									
Hauser 2010a (blunt)	47	64	160	69	86	188	79.3%	-2 20 [-3 78 -0 62]	 _
Hauser 2010b (penetrating)	3.8	0.4 6	29	57	6.4	33	20.7%	-1 90 [-4 99 1 19]	
Subtotal (95% CI)		-	189	•		221	100.0%	-2.14 [-3.54, -0.73]	◆
Heterogeneity: Chi ² = 0.03, df =	: 1 (P = 0	0.87); (²=0%						_
Test for overall effect: Z = 2.98	(P = 0.0	03)							
1.9.3 Platelets									
Hauser 2010a (blunt)	3.3	8.4	112	3.4	7	117	62.0%	-0.10 [-2.11, 1.91]	
Hauser 2010b (penetrating)	1.6	3.7	15	2.5	4.1	21	38.0%	-0.90 [-3.47, 1.67]	
Subtotal (95% CI)			12/			138	100.0%	-0.40 [-1.98, 1.18]	
Heterogeneity: Chif = 0.23, df =	:1 (P = 1	0.63);1 ~	*=0%						
Test for overall effect: $Z = 0.50$	(P = 0.6	2)							
1.9.4 FC									
Hauser 2010a (blunt)	1.5	6.7	28	1.3	4.7	28	75.7%	0.20 [-2.83, 3.23]	
Hauser 2010b (penetrating)	0.1	0.4	1	0.4	2.7	1	24.3%	-0.30 [-5.65, 5.05]	
Subtotal (95% CI)			29			29	100.0%	0.08 [-2.56, 2.72]	
Heterogeneity: Chi ² = 0.03, df =	: 1 (P = 0	0.87); I	= 0%						
Test for overall effect: Z = 0.06	(P = 0.9)	5)							
195 Cryoprecipitate									
Houser 2010e (blunt)	0.0	2.2	24	1.2	10	44	01500	100100101010	
Hauser 2010a (pluitt)	0.9	3.3	34 0	1.3	4.3	41	16 5 96	-0.40 [-2.12, 1.32]	
Subtotal (95% CI)	1.0	4.1	42	2	4.0	52	100.0%	-0.40 [-1.98, 1.18]	
Heterogeneity: Chi ² = 0.00. df =	: 1 (P = 1	1 00\· I	²= ∩%						
Test for overall effect: Z = 0.50	(P = 0.6	2)							
									Favours rEVIIa Favours placebo
Test for subgroup differences:	Chi ² = 6	i.37, df	= 4 (P	= 0.17),	, I ² = 37	7.2%			. Erodio II fild T drodio pracobo

Table 4.41 Results for rFVIIa versus no rFVIIa: Patients with critical bleeding – transfusion volume, other blood components

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ª	(no. of trials) included in analysis	population	(Location)			rFVIIa mean ± SD (n)	No rFVIIa mean ± SD (n)	Mean Difference (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting									
Updated in this review *additional data from primary	N = 573 (1 RCT) Hauser 2010	Adult patients with blunt or penetrating trauma who are critically bleeding	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Allogenic transfusions, units to 24 hrs blunt penetrating	17.1 ± 26.8 (198) 11.2 ± 15 (39)	20.7 ± 25.7 (228) 16.8 ± 19.3 (35)	NR NR	p = 0.09 No difference p = 0.03 Favours rFVIIa
					FFP, units to 24 hours blunt penetrating	4.7 ± 6.4 (160) 3.8 ± 6.0 (29)	6.9 ± 8.6 (188) 5.7 ± 6.4 (33)	NR NR	p < 0.001 Favours rFVIIa p = 0.04 Favours rFVIIa
					Platelets, units to 24 hours blunt penetrating	3.3 ± 8.4 (112) 1.6 ± 3.7 (15)	3.4 ± 7.0 (117) 2.5 ± 4.1 (21)	NR NR	p = 0.84 No difference p = 0.08 No difference
					FC, units to 24 hours blunt penetrating	1.5 ± 6.7 (28) 0.1 ± 0.4 (1)	1.3 ± 4.7 (28) 0.4 ± 2.7 (1)	NR NR	<i>p</i> = 0.68 No difference <i>p</i> = 0.92 No difference
					Cryoprecipitate, units to 24 hours blunt penetrating	0.9 ± 3.3 (34) 1.6 ± 4.1 (8)	1.3 ± 4.3 (41) 2.0 ± 4.8 (11)	NR NR	p = 0.66 No difference p = 0.33 No difference
	N = 277 (2 RCTs) Boffard 2005a&b	Adult patients with severe bleeding due blunt or penetrating trauma	Trauma (multicentre, multicountry)	rFVIIa (therapeutic) vs placebo	Transfusion volume, mL to 48 hours, FFP, platelets or cryoprecipitate	NR Among patients surviv favouring rFVIIa repor and a trend towards fr (See Boffard 2009, figu	NR ving 48 hours, a significa ted for FFP (p = 0.023), a or cryoprecipitate (p = 0.0 ure 2)	NR int difference nd platelets (p = 0.023) 053).	No significant difference
Curry 2011 SR	N = 277 (2 RCTs) Boffard 2005a&b	Adult patients with severe bleeding due	Trauma (multicentre,	rFVIIa (therapeutic) vs	Transfusion volume, mL to 48 hours	Post-hoc analyses	of Boffard 2005a&b l	by Rizoli 2006	
Moderate quality		blunt or penetrating trauma	multicountry)	placebo	FFP	(n = 60) NR	(n = 76) NR	NR	Favours rFVIIa p = 0.04
		patients)			platelets	(n = 60) NR	(n = 76) NR	NR	No significant difference p = 0.09

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results									
Study design ª	(no. of trials) included in analysis	population	(Location)			rFVIIa mean ± SD (n)	No rFVIIa mean ± SD (n)	Mean Difference (95% CI)	Statistical significance p-value Heterogeneity ^b						
Medical emergency - no comparative evidence found															
Surgical setting															
Yank 2011 SR High quality	N = 172 (1 RCT) Gill 2009	Adult patients who had undergone cardiac surgery and were bleeding. Adult patients who (multicentre, multicountry) Adult patients who (multicentre, multicountry)		rFVIIa (therapeutic) vs placebo	Total transfusion volume*, mL 40 ug/kg rFVIIa 80ug/kg rFVIIa *inclusive of all products	(n = 104) median (IQR) 640 (0, 1920) 500 (0, 1750)	(n = 68) median (IQR) 825 (326.5, 1893)		Favours rFVIIa p = 0.047 p = 0.042						
Obstetrics and maternity setting															
Lavigne- Lissalde 2015 RCT High risk of bias	N = 84	Women (aged 18 years or older) with severe persistent primary PPH ^d after sulprostone treatment	Obstetrics (multicentre, France, Switzerland)	rFVIIa (therapeutic) vs standard care	Transfusion volume, number of units FFP PC	(n = 42) median (IQR) 0 (0, 3) NR	(n = 42) median (IQR) 0 (0, 4) NR	NR NR	No significant difference p = NR p = NR						
Franchini 2010 SR (case series) High risk of bias	N = 272 (9 case series)	Women with severe PPH (≥ 500 mL after vaginal delivery and ≥ 1000 mL after caesarean delivery)	Obstetrics and gynaecology (various countries including Europe and Australia)	rFVIIa (therapeutic) vs placebo or no rFVIIa		The authors identified studies, therefore atte published case report: management of sever Three retrospective co Hossain 2007) identifie Section 3.4.4).	no RCTs, case-control of mpted to extract useful s (N>10) to provide recon e PPH. hort studies (Kalina 2011 ed and discussed in PBN	r interventional cohort information from nmendations for the , Ahonen 2007, 1 Module 5 (TR vol. 1	Women who received rFVIIa were given more fibrinogen and platelets than the comparator groups, but those women had more severe haemorrhaging.						

Paediatrics - no comparative evidence found

Abbreviations: CI, confidence interval; hrs, hours; FC, fibrinogen concentrate; M-H, Mantzel-Hentzel; NR, not reported; OR, odds ratio; PPH, postpartum haemorrhage; RR, relative risk; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

d. Defined as the loss of more than 1500 mL of blood within 24 hr after vaginal or caesarean delivery.

4.7 Blood components (Question 6)

Question 6 - (Interventional)

In patients with critical bleeding, what is the effect of fresh frozen plasma, cryoprecipitate, fibrinogen concentrate, prothrombin complex concentrate and/or platelet transfusion on RBC transfusion and patient outcomes?

4.7.1 Methods

This review assessed the evidence of fresh frozen plasma (FFP), cryoprecipitate (CRYO), fibrinogen concentrate (FC), platelet (PLT) and prothrombin complex concentrate (PCC) on red blood cell (RBC) transfusion and patient outcomes in patients with critical bleeding as outlined in Figure 4.26.

Figure 4.26 PICO criteria: Question 6 – effect of blood component therapy on patient outcomes



ARDS, acute respiratory distress syndrome; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ICU, intensive care unit; LOS, length of stay; PCC, prothrombin complex concentrate; PLT, platelets; RBC, red blood cells, TE, thromboembolism

a. 1 vs 1; 2 vs 2; etc.

b. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). c. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. d. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2009 for evidence of FFP, CRYO, FC and PLT and limited to studies published after 1990 for evidence of PCC. However, primary studies published prior to the date limits that had been identified in a systematic review were included. There were no restrictions applied in relation to study size for RCTs. The protocol outlined restrictions for observational studies (at least 500 participants in total), however, due to the evidence identified, this was not applied.

Assuming all relevant primary studies had been identified in the included systematic reviews; the systematic screen of RCTs was limited to studies published after 2015. This is based on the most recent identified systematic review (Cannon 2017), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting.

An updated literature search was conducted in August 2019 and again in September 2021 to identify any new studies meeting the eligibility criteria. In these updated searches, the focus was the identification of systematic reviews, with date limitations based on the most recent systematic reviews used to identify any new RCTs.

Assuming all relevant primary studies have been identified in the included systematic review studies; the systematic screen for RCTs was limited to studies published after the search date outlined in the systematic review.

- For FC, the date limit for RCTs was 2019 based on the most recent identified systematic review (Stabler 2020), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting.
- For PCC, the date limit for screening RCTs was 2020 based on the most recent identified systematic review (van den Brink 2020), which was assumed to have identified all relevant RCTs in the trauma and non-trauma setting.
- For all other blood component therapy, the date limit for screening RCTs was 2019.

4.7.2 Summary of evidence

4.7.2.1 Systematic reviews evidence

Eleven systematic reviews (66, 72, 102-110) were identified in the literature search that were relevant to the research question. The main characteristics and quality of these systematic reviews and relevant outcomes assessed are summarised in Table 4.42.

Four systematic reviews (van den Brink 2020, Fabes 2018, Lunde 2014, Warmuth 2012) assessed the effect of blood component therapy on patient outcomes in mixed clinical settings (including trauma, surgical and obstetrics); 3 systematic reviews assessed FC and one systematic review assessed PCC. One systematic review (Zaidi 2020) assessed the effect of FC on patient outcomes in the obstetric and maternity setting.

Six systematic reviews (Stabler 2020, Coccolini 2019, Rijnhout 2019, McQuilten 2018, Mengoli 2017, Aubron 2014) assessed the effect of blood component therapy on patient outcomes in the trauma setting; 3 systematic reviews assessed FC, 2 systematic reviews assessed FFP, one systematic review assessed FC and CRYO, and one systematic review assessed any blood component.

The reviews included 15 RCTs and 17 observational cohort studies that examined the effect of blood components in patients who were critically bleeding.

A matrix illustrating the overlap of RCTs identified in the included systematic reviews is provided in Table 4.43.

A matrix illustrating the overlap of cohort studies identified in the included systematic reviews is provided in Table 4.44.

Review ID <i>Review Quality</i>	Study design	Population	Intervention	Comparison	Outcomes				
Mixed trauma a	nd non-trauma s	etting							
van den Brink 2020 (103) <i>Low</i>	SR of RCTs and observational studies (17 studies)	Patients with critical bleeding	PCC	No PCC	Mortality Morbidity				
Fabes 2018 (106) High	SR of RCTs (31 studies)	Adults and children at risk of or with critical bleeding	FC	No FC or other blood components	Mortality Morbidity Transfusion volumes				
Lunde 2014 (109) <i>Critically low</i>	SR of RCTs and nonrandomised trials (30 studies)	Patients with critical bleeding	FC	No FC	Mortality Transfusion volume				
Warmuth 2012 (110) <i>Low</i>	SR of RCTs and nonrandomised trials (4 studies)	Adults/children with major haemorrhage	FC	No FC	Transfusion volume				
Trauma setting									
Rijnhout 2021 (66) <i>Low</i>	SR of RCTs and observational studies (9 studies)	Civilian blunt trauma patients	FFP	No FFP	Mortality				
Stabler 2020 (102) <i>Moderate</i>	SR of RCTs and observational studies (26 studies)	Patients older than 16 years of age with trauma-related bleeding	FC	No FC	Mortality Morbidity LOS				
Coccolini 2019 (105) <i>Moderate</i>	SR of RCTs (2 studies)	Severely injured adult trauma patients	FFP	Standard resuscitation protocol	Mortality Morbidity				
McQuilten 2018 (72) <i>Moderate</i>	SR of RCTs (16 studies)	Paediatric and adult trauma patients who had or were expecting to receive a massive transfusion	FC CRYO + MTP	No FC (n=45) MTP (n=41)	Mortality Morbidity LOS Transfusion volumes				
Mengoli 2017 (107) <i>Low</i>	SR of RCTs and observational studies (7 studies)	Trauma patients with critical bleeding	FC	No FC or other blood components	Mortality Morbidity Transfusion volume				
Aubron 2014 (108) <i>Critically low</i>	SR of observational studies (8 studies)	Trauma patients with critical bleeding	FC	No FC or other blood components	Mortality Morbidity Transfusion volume				
Obstetrics and r	maternity setting								
Zaidi 2020 (104) <i>Low</i>	SR of RCTs (5 studies)	Women with PPH	FC	No FC	Mortality Morbidity LOS Transfusion volume				

Table 4.42 Characteristics and quality of systematic reviews by clinical setting: blood components

CYRO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; LOS, length of stay; MTP, massive transfusion protocol; MOF, multiple organ failure; PCC, prothrombin complex concentrate; PPH, primary postpartum haemorrhage; RBC, red blood cells; RCTs, randomised controlled trials; SR, systematic review

Table 4.43	Overlap table of RCTs identified by included systematic reviews: blood
	components

		Tra	uma							Obs		Surgical								
	Study ID	iegler 2019	urry 2015	loore 2018	perry 2018	urry 2018	ascimento 2016	kbari 2018	Inerhofer 2017	ucena 2020	ollins 2017	vikkelsø 2015	ilecen 2017	ahe-Meyer 2013	ahe-Meyer 2016	alas 2014	anaka 2014	eppsson 2016	ance 2012	
	Fabes 2018	N	0	~	0)	✓	∠ ✓	4	_= ✓		√	~	√	√	√	✓	► ✓	X	X	
۵	McQuilten 2018		~				~													
≥	Coccolini 2019			~	~															
svie	Rijnhout 2019			~	~															
ž	Stabler 2020	1				~	~	1		~										
	Zaidi 2020										~	~								

Obs, obstetrics; RCT, randomised controlled trial

 \checkmark = study included in this review; X = study did not meet the inclusion criteria for this review.

		Tra	uma																										Surgical			Obs
	Study ID	Innerhofer 2013	Nienaber 2011	Wafaisade 2013	Schöchl 2011	Joseph 2016	Joseph 2014	O'Reilly 2014	Holcomb 2017	Shackelford 2017	Inokuchi 2017	Schlimp 2016	Almskog 2020	Zeeshan 2019	Jehan 2018	Yamamoto 2016	Bocci 2019	David 2016	Gonzalez-Guerrero 2017	Grassetto 2012	Hilbert 2013	ltagaki 2020	Javier 2019	Schöchl 2010	Schöchl 2014	Schlimp 2013	Schlimp 2016	Seebold 2019	Rahe-Meyer 2009a	Rahe-Meyer 2009b	Bilecen 2013	Ahmed 2012
	Stabler 2020		~	~	1						~	х	1			х	х	х	х	х	х	х	х	х		х		х				
0	van den Brink 2020					~	~							~	~																	
≥	Rijnhout 2019							~	×	~																						
vie	Mengoli 2017	1	~	~	~																											
Re	Aubron 2014	1	~	1	~																											
	Lunde 2014	~	~	~																									~	~	✓	~
	Warmuth 2012																												~	~		

Table 4.44 Overlap table of cohort studies identified by included systematic reviews: blood components

Obs, obstetrics

✓ = study included in this review

-- = no usable data

X = study did not meet the inclusion criteria for this review for reasons including comparator or intervention out of scope.

Fresh frozen plasma versus no FFP (or varying administration of)

Two systematic reviews (Coccolini 2019, Rijnhout 2019) identified 2 RCTs (Moore 2018, Sperry 2018) conducted in adult trauma patients relevant to this review question (see Section 4.7.2.2). There were discrepancies found across the 2 systematic reviews for the outcome of mortality. In one review (Rijnhout 2019), authors reported the outcome data for one RCT (Sperry 2018) as FFP combined with RBC and plasma. As this does not meet criteria for this question, data from Rijnhout 2019 was not used. Outcome data reported by Coccolini 2019 was used in the meta-analysis.

Four systematic reviews (Aubron 2014, Mengoli 2017, Lunde 2014, Rijnhout 2019) identified 4 cohort studies (Holcomb 2017, Shackelford 2017, O'Reilly 2014, Innerhofer 2013) in the trauma setting relevant to the review question (see Section 4.7.2.3).

The search did not find any additional SRs that examined the effect of FFP compared to no FFP (or varying administration of) in patients with critical bleeding in another setting (i.e. perioperative, obstetric, paediatric).

Cryoprecipitate versus no CRYO (or varying administration of)

One systematic review (McQuilten 2018) identified one RCT (Curry 2015) relevant to the review question (see Section 4.7.2.2).

Platelets

The search did not identify any SRs that assessed the use of PLT compared to no PLT (or varying administration of) in patients with critical bleeding on patient outcomes.

Fibrinogen concentrate versus no FC (or varying administration of)

Six systematic reviews (Fabes 2018, McQuilten 2018, Coccolini 2019, Rijnhout 2019, Stabler 2020, Zaidi 2020) identified 12 RCTs that were relevant to the review question (see Section 4.7.2.2). There were slight differences in reporting of outcome data across systematic reviews. Authors of 2 reviews (McQuilten 2018, Stabler 2020) reported per protocol mortality outcome data for one RCT (Nascimento 2016) and another review (Fabes 2018) reported intent-to-treat data. To reduce potential bias, mortality data reported by Fabes 2018 was used in the meta-analysis.

Authors of one review (Fabes 2018) reported lower patient numbers for the morbidity outcome of thrombosis for one RCT (Collins 2017) compared to another review (Zaidi 2020) which reported intent-to-treat numbers. Intent-to-treat morbidity (thrombosis) data reported by Zaidi 2020 was used in the meta-analysis.

Five systematic reviews (Stabler 2020, Aubron 2014, Mengoli 2017, Lunde 2014, Warmuth 2012) identified 9 cohort studies that were relevant to the review question. There were slight differences in reporting of outcome data across 2 systematic reviews. Authors of one review (Stabler 2020) reported organ failure outcome data for one cohort study (Wafaisade 2013) and another review (Aubron 2014) reported multiple organ failure outcome data. Morbidity data for multiple organ failure reported by Aubron 2014 was used in the meta-analysis to align with the key outcome of interest in this review.
Prothrombin complex concentrate versus no PCC (or varying administration of)

One systematic review (van den Brink 2020) identified 4 cohort studies (Johan 2018, Zeeshan 2019, Joseph 2014, Joseph 2016) that assessed the use of PCC and FFP versus FFP alone in the trauma setting (see Section 4.7.2.3).

4.7.2.2 Randomised controlled trials

The included systematic reviews identified 15 RCTs that were relevant to this review (Curry 2015, Moore 2018, Sperry 2018, Curry 2018, Nascimento 2016, Akbari 2018, Innerhofer 2017, Lucena 2020, Collins 2017, Wikkelsø 2015, Bilecen 2017, Rahe-Meyer 2013, Rahe-Meyer 2016, Galas 2014, Tanaka 2014). No additional RCTs were identified in the systematic review and handsearching process.

The main characteristics and quality of the included RCT and the relevant outcomes assessed are detailed in Table 4.45.

Review ID Risk of bias	Study design	Population N	Intervention	Comparison	Outcomes
Trauma sett	ing				
Lucena 2020 (111) <i>Low</i>	RCT Unblinded	Patients (18-80 years) admitted to ED with severe trauma (ISS \geq 15) N=32	FC *early administration (50 mg/kg)	No FC	Transfusion volume Morbidity LOS
Akbari 2018 (112) <i>High</i>	quasi-RCT	Patients with severe blunt trauma N=90	FC	FFP	Mortality LOS Transfusion volume Morbidity
Curry 2018 (E-FIT 1) (113) <i>Low</i>	RCT MC, double- blinded	Adult trauma patients who triggered MHP N=48	FC *within 45 minutes admission, maintaining fibrinogen levels ≥ 2 g/L during active haemorrhage	No FC	Transfusion volume Mortality Morbidity
Moore 2018 (COMBAT) (114) <i>Low</i>	RCT, SC	Trauma patients in haemorrhagic shock* N=125 * SBP ≤ 70 mmHg or 71-90 mmHg plus HR ≥ 108 bpm	FFP (prehospital)	Saline	Mortality Transfusion volume
Sperry 2018 (PAMPer) (115) <i>Low</i>	RCT, MC	Injured patients at risk of haemorrhagic shock N=501	FFP (prehospital)	No FFP (Standard care)	Mortality Morbidity Transfusion volumes
Innerhofer 2017 (RETIC) (116) <i>Low</i>	RCT SC, unblinded	Adult trauma patients (18-80 years) with ISS >15 and clinical signs or risk of haemorrhage N=100	FC	FFP	Morbidity LOS Transfusion volume
Nascimento 2016 (FiiRST) (117) <i>Low</i>	RCT SC, double- blinded	Adult patients (≥ 18 years) with blunt or penetrating trauma at risk for significant haemorrhage	FC	No FC	Mortality Morbidity Transfusion volume

Table 4.45 Characteristics and quality of RCTs by clinical setting: blood components

Review ID Risk of bias	Study design	Population N	Intervention	Comparison	Outcomes
		N=45			
Curry 2015 (CRYOSTAT) (118) <i>Low</i>	RCT Unblinded	Adult patients (\geq 16 years) with active bleeding and requiring activation of the MTP N=41	CRYO	No CRYO	Mortality Morbidity Transfusion volume
Surgical sett					
Bilecen 2017 (119) <i>High</i>	RCT	Patients (> 18 years) with intraoperative bleeding following high-risk cardiac surgery N=120	FC	No FC	Transfusion volume Mortality Morbidity
Rahe-Meyer 2016 (120) (REPLACE) <i>High</i>	RCT MC	Adult patients with intraoperative bleeding following cardiac surgery N=152	FC	No FC	Transfusion volume Mortality
Tanaka 2014 (121) <i>High</i>	RCT SC	Patients with moderate or severe bleeding following cardiac surgery N=20	FC		Morbidity Mortality
Rahe-Meyer 2013 (122) <i>High</i>	RCT SC, double- blinded	Patients (≥ 18 years) undergoing elective aortic replacement surgery with CPB N=61	FC	No FC	Transfusion volume
Surgical sett	ting, paediatr	ics			
Galas 2014 (123) High	RCT	Patients (< 7 years) with diffuse bleeding following elective cardiac surgery N=60	FC	CYRO	Transfusion volume Morbidity LOS Mortality
Obstetric an	d maternal se	etting			
Collins 2017 (124) <i>High</i>	RCT MC	Women (> 18 years) at 24 weeks gestation who had ongoing major PPH (1000–1500 mL blood loss) N=57	FC	Placebo	Mortality, Morbidity Transfusion volume LOS
Wikkelsø 2015 (125) <i>High</i>	RCT MC, double- blinded	Women with early postpartum haemorrhage N=244	FC	Saline	Mortality Morbidity Transfusion volume (RBC)

BPM, beats per minute; CPB, cardiopulmonary bypass; CRYO, cryoprecipitate; ED, emergency department; FC, fibrinogen concentrate; FFP, fresh frozen plasma; HR, heart rate; ISS, injury severity score; LOS, length of stay; MC, multicentre; MHP, major haemorrhage protocol; MT, massive transfusion; PLT, platelet; PPH, postpartum haemorrhage; RBC, red blood cell; RCT, randomised controlled trial; SBP, systolic blood pressure, SC, single centre; TEG, thromboelastography
 a source the source of the

a. as evidenced by a) SBP < 100 mmHg and b) requiring uncrossmatched RBC transfusion at any time from injury until 30 minutes after hospital arrival

Fresh frozen plasma

Two RCTs (Moore 2018, Sperry 2018) were included in this review. Both RCTs were conducted in US trauma centres and enrolled severely injured adults (aged 18 – 90 years) with systolic blood pressure 70 mmHg or lower or 71–90 mmHg and heart rate more than 108 beats per minute thought to be due to acute blood loss, either before the arrival of air medical transport or before arrival at the trauma centre. The RCTs assessed the use of 2 units of FFP compared with the standard resuscitation protocol according to local rules. Moore 2018 included a total of 125 patients in the analysis and Sperry 2018 included 501 patients. Both RCTs reported on the outcomes of mortality and morbidity (including acute lung injury and multiple organ failure) and were judged by the systematic review authors to be at low risk of bias.

Cryoprecipitate

One RCT (Curry 2015) evaluated the effect of CRYO on mortality, morbidity and transfusion volume in trauma patients with major haemorrhage requiring activation of the major haemorrhage protocol. The study included a total of 44 patients and was carried out in 2 civilian UK trauma centres. Risk of bias was judged by review authors as unclear due to small sample size and lack of blinding of participants, clinical staff and research staff.

Platelets

The search did not identify any RCTs that assessed the use of PLT compared to no PLT (or varying administration of) in patients with critical bleeding on patient outcomes.

Fibrinogen concentrate

Five RCTs were conducted in critically bleeding trauma patients (Lucena 2020, Curry 2018, Akbari 2017, Innerhofer 2017, Nascimento 2016), 2 RCTs were in the obstetrics setting (Collins 2017, Wikkelsø 2015), and 5 RCTs were in the surgical setting (Bilecen 2017, Rahe-Meyer 2016, Galas 2014, Tanaka 2014, Rahe-Meyer 2013).

Trauma setting

Five RCTs conducted in Austria, UK, Canada, Iran and Brazil were found that assessed the use of FC in adult patients with severe trauma. Three RCTs (Curry 2018, Nascimento 2016, Lucena 2020) compared the use of FC with saline or no FC, one RCT (Akbari 2017) compared FC to an active (FFP) and an inactive (no coagulation factor) comparator, and one RCT (Innerhofer 2017) compared FC to an active comparator (FFP) only.

The studies were assessed to be at overall moderate risk of bias due to lack of allocation concealment, blinding of study personnel and outcome assessors, incomplete outcome data and selective reporting.

Surgical setting

Four RCTs were conducted in the Netherlands, Germany and US and evaluated the therapeutic use of FC in the setting of cardiac surgery.

Three RCTs (Bilecen 2017, Rahe-Meyer 2013, Rahe-Meyer 2016) compared the use of FC with saline while one RCT (Tanaka 2014) compared the use of FC with one unit of PLT. All 4 RCTs were assessed by the systematic review authors to have overall no serious concerns of bias, however domains assessed to have some risk of bias included allocation concealment, blinding, incomplete outcome data and selective reporting.

One RCT (Galas 2014) evaluated the safety and efficacy of FC compared with CRYO in paediatric patients (aged less than 15 years) undergoing cardiopulmonary bypass surgery. The RCT was conducted in 63 children in Brazil, of which 30 patients were randomised to receive 60mg/kg FC and 33 patients were randomised to receive 10 mL/kg cryoprecipitate if they had diffuse bleeding requiring haemostatic therapy and plasma fibrinogen concentration less than 1 g/L. The primary outcome for this study was post-operative blood loss during 48 hours of surgery. The RCT was assessed by the systematic review to be at low risk of bias.

Obstetric and maternity setting

Two RCTs (Collins 2017, Wikkelsø 2015) were conducted in Denmark and the UK and evaluated the use of FC in the obstetrics setting. In both RCTs, adult women with postpartum haemorrhage were randomised to receive FC or saline and reported on the outcomes of mortality and morbidity. The RCTs were assessed by the review authors to be at overall low risk of bias for these outcomes.

Prothrombin complex concentrate

The search did not identify any RCTs that examined the effect of PCC compared to no PCC (or varying administration of) in patients with critical bleeding on patient outcomes.

4.7.2.3 Observational and cohort studies

The included systematic reviews identified 16 cohort studies that were relevant to this review (Almskog 2020, Zeeshan 2019, Jehan 2018, Holcomb 2017, Shackelford 2017, Joseph 2016, Joseph 2014, O'Reilly 2014, Innerhofer 2013, Wafaisade 2013, Nienaber 2011, Schöchl 2011, Rahe-Meyer 2009a, Rahe-Meyer 2009b, Bilecen 2013, Ahmed 2012).

The main characteristics and quality of these studies and the relevant outcomes assessed are detailed in Table 4.46.

The systematic review process identified 2 cohort studies (Inokuchi 2017, Yamamoto 2016) relevant to this review. Both studies were identified in the included systematic reviews, however one study (Yamamoto 2016) did not present analysis for comparator of interest (i.e. FC compared to no FC) and is not reported further.

Among the identified cohort studies, 12 were retrospective cohorts. As outlined in Section 3.1.4, retrospective cohort studies were to be excluded. However, given the lack of available evidence for blood component therapy, these studies were included in the review.

Table 4.46	Characteristics and quality of observational and cohort studies by clinical setting:
	the effect of blood component therapy on patient outcomes

Review ID Risk of bias	Study design	Population	Intervention	Comparison	Outcomes					
Trauma setting										
Almskog 2020 (126) <i>High</i>	Retrospective (matched) cohort	Adult patients with blunt trauma N=216	FC	No FC	Mortality Morbidity LOS					
Zeeshan 2019 (127)	Retrospective cohort	Severely injured trauma patients	4-factor PCC (+FFP)	FFP	Mortality Transfusion volumes					

Review ID Risk of bias	Study design	Population	Intervention	Comparison	Outcomes
Moderate		N=468			LOS Morbidity
Jehan 2018 (128) <i>Moderate</i>	Retrospective cohort	Adult trauma patients N=120	4-factor PCC (+FFP)	FFP	Transfusion volumes Mortality LOS Morbidity
Holcomb 2017 (129) <i>High</i>	Retrospective (matched) cohort	Adult civilian trauma patients N=109	FFP	Standard care	Mortality
Inokuchi 2017 (130) <i>Serious</i>	Retrospective (before and after) cohort SC	Patients with pelvic fractures stemming from blunt injury N=224	Early FC (+MHP)	MHP (without FC)	Transfusion volume Mortality
Shackelford 2017 (131) <i>High</i>	Retrospective (matched) cohort	Military trauma patients N=386	FFP	Standard care	Mortality
Joseph 2016 (132) <i>Moderate</i>	Retrospective cohort	Trauma patients N=81	3-factor PCC (+FFP)	FFP	Mortality Morbidity Transfusion volume LOS
Joseph 2014 (133) <i>Moderate</i>	Retrospective cohort	Trauma patients N=252	3-factor PCC (+FFP)	FFP	Mortality Transfusion volumes Morbidity
O'Reilly 2014 (134) High	Retrospective (matched) cohort	Military trauma patients N=194	FFP	Standard care	Mortality
Innerhofer 2013 (135) <i>High</i>	Retrospective cohort study	Adult trauma patients (≥18 years) with an ISS ≥15 N=144	FFP (+FC +/- PCC)	FC +/- PCC	Morbidity Transfusion volume
Wafaisade 2013 (136) High	Retrospective (matched) cohort	Severely injured trauma patients N=588	FC	No FC	Mortality Morbidity LOS Transfusion volume
Nienaber 2011 (137) High	Retrospective (matched) cohort	Adult trauma patients (≥18 to ≤70 years) with ISS ≥16. N=36	FC and/or PCC (ROTEM-guided)	Standard care guided MHP (1:1 RBC:FFP)	Mortality Morbidity LOS Transfusion volume
Schöchl 2011 (138) <i>High</i>	Prospective (unmatched) cohort	Trauma patients who received \geq 5 units of RBC concentrate within 24 hours. N=36	FC +/- PCC (ROTEM-guided)	Standard care guided transfusion in patients receiving >2 units FFP (no FC or PCC)	Mortality Transfusion volumes
Surgical setti	ng				
Bilecen 2013 (139) <i>High</i>	Prospective cohort SC	Patients who underwent complex cardiac surgery N=120	FC	No FC	Transfusion volume

Review ID Risk of bias	Study design	Population	Intervention	Comparison	Outcomes
Rahe-Meyer 2009a (140) <i>High</i>	Comparative prospective	Patients with TAAA N=18	FC	Standard transfusion algorithm	Mortality Transfusion volume
Rahe-Meyer 2009b (141) <i>High</i>	Comparative study with historical control	Patients undergoing AV- AA N=15 (Group B and C)	FC	Standard transfusion algorithm	Mortality Transfusion volume
Obstetric and	l maternal setting				
Ahmed 2012 (142) <i>High</i>	Retrospective cohort	Patients with major obstetric haemorrhage N=34	FC	CRYO	LOS Transfusion volume

AV-AA, aortic valve operation and ascending aorta replacement; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ISS, injury severity score; LOS, length of stay; PCC, prothrombin complex concentrate; RBC, red blood cell; SC, single centre; TAAA, thoracoabdominal aortic aneurysm

Fresh frozen plasma

Holcomb 2017 (129) was a multicentre, prospective cohort study conducted in the US that assessed the effect of prehospital transfusion of FFP alone, RBC alone or both compared with a propensity score matched group of 109 patients with penetrating trauma who received standard care (crystalloid resuscitation). A total of 26 patients received FFP only, 8 patients received RBC only and 75 patients received both FFP and RBC and constitute the interventional arm in this analysis. The study was found to be at high risk of bias due to imbalances in baseline characteristics which limited matching.

Innerhofer 2013 (135) was a single centre, prospective cohort study conducted in Austria that assessed the effect of FFP in 144 patients with blunt major trauma. All patients in the study received FC and PCC; 78 patients additionally received FFP transfusions and constitute the interventional arm in this analysis. Review authors judged the study to have high risk of bias due to small sample sizes, inadequate follow-up and lacked rigorous analyses.

Two retrospective cohort studies (Shackelford 2017, O'Reilly 2014) investigated the effect of prehospital transfusion of FFP compared to standard of care in military trauma patients in Afghanistan with gunshot wounds or explosive trauma. O'Reilly 2014 (134) assessed prehospital blood transfusion in 194 patients. A total of 97 patients received a median of one unit RBC and 2 units of FFP and 97 patients received standard of care. Shackelford 2017 (131) was a study of 386 US military combat casualties who received prehospital blood transfusion between 2012 to 2015. A total of 54 patients received RBC and FFP; 332 patients received standard of care.

Review authors judged the studies to be at high risk of bias due to retrospective analyses and a lack of uniform guidelines for initiating prehospital blood transfusion which makes it difficult to determine the effect of individual blood components.

Cryoprecipitate

The search did not identify any cohort studies that assessed the use of CRYO compared to no CRYO (or varying administration of) in patients with critical bleeding on patient outcomes.

Platelets

The search did not identify any cohort studies that assessed the use of PLT compared to no PLT (or varying administration of) in patients with critical bleeding on patient outcomes.

Fibrinogen concentrate

Five cohort studies (Almskog 2020, Wafaisade 2013, Inokuchi 2017, Schöchl 2011, Nienaber 2011) were conducted in the trauma setting, 3 studies (Bilecen 2013, Rahe-Meyer 2009a, Rahe-Meyer 2009b) were in the surgical setting, and one study (Ahmed 2012) was in the obstetrics setting.

Trauma setting

Five cohort studies were conducted in Europe and Japan and examined the effect of FC in trauma patients with critical bleeding. In 2 studies the comparator was no FC (Wafaisade 2013, Almskog 2020), while the remaining 3 cohort studies examined the effect of including fibrinogen concentrate as part of a MHP compared with an MHP without fibrinogen concentrate (Schöchl 2011, Nienaber 2011, Inokuchi 2017). For the purposes of this review, the patients who received FC were considered as the interventional arm for this analysis.

The cohort studies were judged by systematic reviews to be at high risk of bias due to missing data, absence of a clear objective criterion for the activation of MTP and lack of control for potential confounders.

Surgical setting

Three cohort studies were identified in the surgical setting that evaluated the use of FC in patients with major haemorrhage (Bilecen 2013, Rahe-Meyer 2009a, Rahe-Meyer 2009b). All 3 cohort studies were assessed to be at high risk of bias, predominately due to failure to in blinding, lack of information on the allocation of groups and insufficient information about comparability of groups at baseline and at the analysis stage.

Bilecen 2013 was a single centre prospective cohort study that assessed 1075 patients who underwent complex cardiac surgery in the Netherlands. A total of 264 patients received a median dose of 2g FC; the 811 patients that did not receive FC represent the control group. The authors note that due to the nonrandomised design of the study, the association between the infusion of FC and each of the outcomes were likely biased by potential confounders.

Rahe-Meyer 2009a was a pilot study that prospectively enrolled 15 patients undergoing aortic valve operation and ascending aorta replacement surgery in Germany. Five patients received transfusion according to the predefined blood components transfusion algorithm while the remaining 10 patients received FC before being transfused according to the algorithm. Rahe-Meyer 2009b was a retrospective group analysis of 18 patients who underwent elective thoracoabdominal aortic aneurysm surgery. All patients in the study were treated with allogenic blood components according to a predetermined algorithm; 6 patients also received a mean (SD) dose of 7.8 g (2.7) FC as a first step therapy. The small sample size prevents any meaningful analysis of the results.

Obstetric and maternity setting

One cohort study (Ahmed 2012) was found that evaluated the use of FC in women with major obstetric haemorrhage. Among 77 patients with major obstetric haemorrhage, 20 received a mean dose of 4 ± 0.8 g FC and 34 received a mean dose of 2.21 ± 0.35 pooled units of CRYO. Due to the nature of the active comparator, both treatment arms represent eligible interventions. For the purpose of this review, FC has been chosen as the interventional arm for this analysis.

Ahmed 2012 was assessed by review authors to be at serious risk of bias due to small sample size and inadequate follow-up.

Prothrombin complex concentrate

The search did not find any additional studies that examined the effect of PCC compared to no PCC (or varying administration of) in patients with critical bleeding in another setting (i.e. perioperative, obstetric, paediatric).

The 4 cohort studies were conducted in trauma patients presenting to the emergency department (total sample size 924). Two studies (Jehan 2018, Zeeshan 2019) investigated the effect of 4-factor PCC plus FFP compared to FFP only and 2 studies (Joseph 2014, Joseph 2016) investigated the effect of 3-factor PCC plus FFP compared to FFP only. Dose of PCC administered was 25 IU/kg for 3 studies and indication for administration was by clinical judgement for all 4 studies.

Review authors judged the studies as moderate risk of bias due to the retrospective study design, in which PCC was administered based on clinical judgement and may have resulted in confounding and bias. The authors also noted considerable variety in the type and dose for PCC that could lead to under or overrepresentation of the actual effects of PCC on the outcomes.

4.7.3 Results

4.7.3.1 Fresh frozen plasma

Mortality

A summary of the evidence relating to the effect of fresh frozen plasma (FFP) on mortality in patients with critical bleeding is presented in Table 4.47.

A meta-analysis of data from studies included in this review showed no significant difference in mortality at 24 hours (Figure 4.27) or latest reported timepoint (Figure 4.28) between patients who received FFP compared to those who did not.

24 hours

One RCT (Moore 2018) and 2 cohort studies (Holcomb 2017, Shackelford 2017) were identified that reported on the effect of FFP on the outcome of 24-hour mortality among patients with trauma. Combined data from all 3 studies (see Figure 4.27) showed the 24-hour mortality rate in patients who received FFP (16/162, 9.9%) to be lower than that observed among patients who did not receive FFP (83/458, 18.1%). The difference was not statistically significant (RR 0.66; 95% CI 0.28, 1.57; p = 0.35; random effects; I2 = 53%).

Latest timepoint

Two RCTs (Moore 2018, Sperry 2018) and 4 cohort studies (Holcomb 2017, Innerhofer 2013, O'Reilly 2014, Shackelford 2017) reported on the effect of FFP on the outcome of mortality, latest timepoint. All 6 studies were conducted in the trauma setting. Combined data from the 2 RCTs (see Figure 4.28) showed the mortality rate to be 26.4% (78/295) among those who received FFP compared to 31.4% (104/331) among those who did not. The difference was not statistically significant (RR 0.95; 95% CI 0.56, 1.59; p = 0.83; random effects, $l^2 = 38\%$), with moderate statistical heterogeneity observed.

Combined data from the 4 cohort studies (see Figure 4.28) suggested a significant association between FFP and mortality among trauma patients with critical bleeding (RR 0.65, 95%CI 0.43, 0.98; p = 0.04; random effects, $l^2 = 0\%$) with the mortality rate observed among those who received FFP (19.3%, 106/549) being lower than the mortality rate of those who did not receive FFP (24.4%, 218/892).

Figure 4.27 Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Mortality, all-cause (at 24 hours)

	FFP	1	No FFP (or varying)		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
1.1.1 RCTs							
Moore 2018 (RCT, trauma) Subtotal (95% CI)	8	65 65	6	60 60	34.6% 34.6%	1.23 [0.45, 3.34] 1.23 [0.45, 3.34]	
Total events	8		6				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.41 (P	= 0.68)						
1.1.2 Observational Holcomb 2017 (Coh, trauma) Shackelford 2017 (Coh, trauma)	5 3	43 54	10 67	66 332	34.5% 30.9%	0.77 [0.28, 2.09] 0.28 [0.09, 0.84]	
Subtotal (95% CI)		97		398	65.4%	0.47 [0.17, 1.34]	
Total events	8		77				
Heterogeneity: $Tau^2 = 0.27$; $Chi^2 =$ Test for overall effect: Z = 1.41 (P	1.93, df = = 0.16)	1 (P =	0.16); l² = 48%				
Total (95% CI)		162		458	100.0%	0.66 [0.28, 1.57]	
Total events 16 83 Heterogeneity: Tau ² = 0.31; Chi ² = 4.21, df = 2 (P = 0.12); l ² = 53% Test for overall effect: Z = 0.94 (P = 0.35) Test for subgroup differences: Chi ² = 1.68, df = 1 (P = 0.19), l ² = 40.6%							0.05 0.2 1 5 20 Favours FFP Favours no FFP

Figure 4.28 Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Mortality, all-cause (latest reported timepoint)

	FFF	•	No FFP (or vary	FP (or varying) Risk Ratio		Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	l	M-H, Random, 95% Cl
1.2.1 RCTs								
Moore 2018 (RCT, trauma)	10	65	6	60	5.0%	1.54 [0.60, 3.98]		
Sperry 2018 (RCT, trauma)	68	230	98	271	69.2%	0.82 [0.63, 1.05]		
Subtotal (95% CI)		295		331	74.2%	0.95 [0.56, 1.59]		•
Total events	78		104					
Heterogeneity: Tau ² = 0.08; Chi ² =	1.60, df =	1 (P =	0.21); l² = 38%					
Test for overall effect: Z = 0.21 (P =	= 0.83)							
1.2.2 Observational								
Holcomb 2017 (Coh, trauma)	8	43	14	66	7.4%	0.88 [0.40, 1.91]		
Innerhofer 2013 (Coh, trauma)	6	78	5	66	3.4%	1.02 [0.32, 3.18]		
O'Reilly 2014 (Coh, trauma)	8	79	19	97	7.6%	0.52 [0.24, 1.12]		
Shackelford 2017 (Coh, trauma)	6	54	76	332	7.4%	0.49 [0.22, 1.06]		
Subtotal (95% CI)		254		561	25.8%	0.65 [0.43, 0.98]		•
Total events	28		114					
Heterogeneity: Tau ² = 0.00; Chi ² =	2.05, df =	3 (P =	0.56); l² = 0%					
Test for overall effect: Z = 2.05 (P =	= 0.04)							
		540			400.00/	0 70 10 04 0 001		
Total (95% CI)		549		892	100.0%	0.79 [0.64, 0.98]		•
Total events	106		218					
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 =$	4.91, df =	5 (P =	0.43); l² = 0%				0.01	0.1 1 10 100
l est for overall effect: $Z = 2.13$ (P						Favours FFP Favours no FFP		
Test for subgroup differences: $Chi^2 = 1.24$, $df = 1$ (P = 0.27), $l^2 = 19.1\%$								

Table 4.47 Results for FFP versus no FFP: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			FFP n/N (%)	No FFP n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
Trauma setting											
Coccolini 2019 SR <i>Moderate quality</i>	N = 626 (2 RCTs) Moore 2018 Sperry 2018	Severely injured adults with acute blood loss	Trauma centre, civilian (US)	FFP vs standard resuscitation protocol	Mortality, 1 month	78/295 (26.4) 10/65 (15.4) 68/230 (29.6)	104/331 (31.4) 6/60 (10) 98/271 (36.3)	RR 0.86 (0.68, 1.11) 1.54 (0.60, 3.98) 0.82 (0.63, 1.05)	No significant difference p = 0.24 Minimal heterogeneity $l^2 = 38\% (p = 0.21)$		
Mengoli 2017 SR <i>Low quality</i>	N = 144 (1 Coh) Innerhofer 2013	Patients with severe trauma- related bleeding	Trauma centre (SC, Austria)	FFP vs no FFP (± FC and/or PCC in both groups)	Mortality, 30 days	6/78 (7.7)	5/66 (7.6)	NR	No significant difference p = 0.979		
Rijnhout 2019 SR <i>Low quality</i>	N = 495 (2 Coh) Shackelford 2017 Holcomb 2017	Civilian and military trauma patients	Trauma (US, Afghanistan)	FFP vs standard care	Mortality, 24 hours	8/97 (8.2) 3/54 (5.6) 5/43 (11.6)	77/398 (19.3) 67/332 (20.2) 10/66 (15.2)	RR 0.47 (0.17, 1.34) 0.28 (0.09, 0.84) 0.77 (0.28, 2.09)	No significant difference p = 0.16 Moderate heterogeneity $l^2 = 48\% (p = 0.16)$		
	N = 689 (3 Coh) O'Reilly 2014 Shackelford 2017 Holcomb 2017				Mortality, long- term	22/194 (11.3) 8/97(8.2) 6/54 (11.1) 8/43 (18.6)	109/495 (22) 19/97 (19.6) 76/332 (22.9) 14/66 (21.2)	OR 0.54 (0.32, 0.91) ^c 0.37 (0.15, 0.89) 0.42 (0.17, 1.02) 0.85 (0.32, 2.24)	Favours FFP $p = 0.02^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.53)^{\circ}$		

Cl, confidence interval; Coh, cohort study; FC, fibrinogen concentrate; FFP, fresh frozen plasma; NR, not reported; OR, odds ratio; PCC, prothrombin complex concentrate; RCT, randomised controlled trial; RR, relative risk; SC, single centre; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4.

Morbidity

A summary of the evidence reported in the identified systematic reviews relating to morbidity in patients with critical bleeding is presented in Table 4.48.

Three studies identified in the systematic reviews reported on the outcome of morbidity; 2 RCTs (Moore 2018, Sperry 2018) reported on multiple organ failure and acute lung injury and one cohort study (Innerhofer 2013) reported on multiple organ failure and thromboembolic events (Figure 4.29). The studies were not sufficiently powered to detect important differences in event rates, therefore evidence for morbidity outcomes should be considered with caution.

Thromboembolic events

One cohort study (Innerhofer 2013) reported a lower rate of thromboembolic events among patients who received FFP (7.7%, 6/78) compared with those who did not (9.0%, 6/66), but the difference between groups was not significant (RR 0.85, 95% CI 0.29, 2.50; p = 0.76).

Multiple organ failure

A meta-analysis of data from the included studies showed an increased risk of multiple organ failure among patients who received FFP (179/373, 48.0%) compared with those who did not (169/397, 42.6%). The difference between groups was not significant (RR 1.56, 95% CI 0.2, 2.96; p = 0.17; random effects; $I^2 = 68\%$); noting statistical heterogeneity is substantial. The results were not substantially different when only RCT evidence was considered (RR 1.76, 95% CI 0.40, 7.68); p = 0.45; random effects; $I^2 = 58\%$).

Acute lung injury

A meta-analysis of data from the RCTs showed no difference in the risk of acute lung injury between treatment groups. The rate of acute lung injury was 25.7% (76/295) among those who received FFP compared with 24.2% (80/331) among those who did not, corresponding to a RR of 0.99 (95% CI 0.76, 1.30; p = 0.97; random effects; $I^2 = 9\%$).

	FFP		No FFP (or vary	ing)		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl
1.3.1 Thromboembolic events							
Innerhofer 2013 (Coh, trauma)	6	78	6	66	100.0%	0.85 [0.29, 2.50]	——————————————————————————————————————
Subtotal (95% CI)		78		66	100.0%	0.85 [0.29, 2.50]	•
Total events	6		6				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.30 (P = 0.76)						
1.3.2 Multiple organ failure							
Innerhofer 2013 (Coh, trauma)	28	78	12	66	37.8%	1.97 [1.09, 3.57]	
Moore 2018 (RCT, trauma)	6	65	1	60	8.1%	5.54 [0.69, 44.68]	+
Sperry 2018 (RCT, trauma)	145	230	156	271	54.1%	1.10 [0.95, 1.26]	•
Subtotal (95% CI)		373		397	100.0%	1.56 [0.82, 2.96]	◆
Total events	179		169				
Heterogeneity: Tau ² = 0.19; Chi ²	= 6.32, df	= 2 (P =	= 0.04); l ² = 68%				
Test for overall effect: Z = 1.36 (P = 0.17)						
1.3.3 Acute lung injury							
Moore 2018 (RCT, trauma)	28	65	30	60	47.3%	0.86 [0.59, 1.26]	+
Sperry 2018 (RCT, trauma)	48	230	50	271	52.7%	1.13 [0.79, 1.61]	+
Subtotal (95% CI)		295		331	100.0%	0.99 [0.76, 1.30]	•
Total events	76		80				
Heterogeneity: Tau ² = 0.00; Chi ²	= 1.10, df	= 1 (P =	= 0.29); l² = 9%				
Test for overall effect: Z = 0.04 (P = 0.97)						
							Favours FFP Favours no FFP

Figure 4.29 Forest plot of comparison: FFP vs no FFP (or varying administration of), outcome: Morbidity

Test for subgroup differences: Chi² = 1.77, df = 2 (P = 0.41), l² = 0%

Table 4.48 Results for FFP versus no FFP: Patients with critical bleeding – Morbidity: any adverse outcome

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study designª	(no. of trials) included in analysis	population	(Location)			FFP n/N (%)	No FFP n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
Trauma setting											
Coccolini 2019 SR Moderate quality	N = 626 (2 RCTs) Moore 2018 Sperry 2018	Severely injured adults with acute blood loss	Trauma centre, civilian (US)	FFP vs standard resuscitation protocol	Acute lung injury	76/295 (25.8) 28/65 (43.1) 48/230 (20.9)	80/331 (24.2) 30/60 (50) 50/271 (18.5)	OR 1.03 (0.71, 1.50) OR 1.17 (0.75, 1.81)	No significant difference p = 0.87 Minimal heterogeneity $l^2 = 3\% (p = 0.31)$		
				FFP vs standard resuscitation protocol	MOF	149/295 (50.5) 4/65 (6.2) 145/230 (63.0)	157/331 (47.4) 1/60 (1.7) 156/271 (57.6)	OR 1.30 (0.92, 1.86) OR 3.87 (0.42, 35.63) OR 1.26 (0.88, 1.80)	No significant difference p = 0.14 No significant heterogeneity $l^2 = 0\% (p = 0.33)$		
Mengoli 2017 SR <i>Low quality</i>	N = 144 (1 Coh) Innerhofer 2013	Patients with severe trauma- related bleeding	Trauma centre (SC, Austria)	FFP ± FC ± PCC vs FC ± PCC	^c Thromboembolism	6/78 (7.7)	6/66 (10)	NR	No significant difference p = 0.772		
Lon quanty					Sepsis ^c	28/78 (35.9)	11/66 (16.7)	NR	No significant difference $p = 0.014$		
					MOF ^c	29/78 (37.2)	12/66 (18.2)	NR	No significant difference $p = 0.015$		

Cl, confidence interval; Coh, cohort; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ISS, injury severity score; MOF, multiorgan failure; NR, not reported; OR, odds ratio; PCC, prothrombin complex concentrate; RCT, randomised controlled trial; SC, single centre; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

Transfusion volumes

A summary of the evidence reported in the identified systematic reviews relating to transfusion volumes in patients with critical bleeding is presented in Table 4.49.

There was one cohort study (Innerhofer 2013) identified by 3 systematic reviews (Mengoli 2017, Aubron 2014, Lunde 2014) that was considered relevant to this review.

Red blood cells

One small cohort study (Innerhofer 2013) reported that the median (IQR) volume of RBC transfused (units to 24 hours) among the 78 patients who received FFP was 7 (4, 11) units, which was significantly higher than the median 2 (0, 6) units of RBC transfused among the 66 patients who did not receive FFP (p = 0.001).

Other blood components

One small cohort study (Innerhofer 2013) reported that the median (IQR) volume of PLT transfused (units to 24 hours) among the 78 patients who received FFP was 0 (0, 1) units, which was significantly higher than the median 0 (0, 0) units of PLT transfused among the 66 patients who did not receive FFP (p = 0.003).

There was no significant difference between treatment groups reported for the dose of FC (grams to 24 hours) and PCC (units to 24 hours) used.

Table 4.49 Results for FFP versus no FFP: Patients with critical bleeding – Transfusion volumes

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study designª	(no. Of trials) included in analysis	population	(Location)			FFP median (IQR)	No FFP median (IQR)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b		
Trauma setting											
Mengoli 2017 SR <i>Low</i>	N = 144 (1 Coh) Innerhofer 2013	Patients with severe trauma- related bleeding	Trauma centre (SC, Austria)	FFP vs no FFP (± FC ± PCC in both groups)	RBC transfusion volume, units to 24 hours	(n = 78) 7 (4, 11)	(n = 66) 2 (0, 6)	NR	Favours FFP p = 0.001 °		
					PLT transfusion volume, units to 24 hours ^c	(n = 78) O (O, 1)	(n = 66) O (O, O)	NR	Favours FFP p = 0.003		
Lunde 2014 SR	N = 144 (1 Coh) Innerhofer 2013	Patients with severe trauma-	Trauma centre (SC, Austria)	FFP vs no FFP (± FC and/or PCC in	FFP transfusion volume, units to 24 hours	(n = 78) 8 (5, 10)	(n = 66) O (O, O)	NR	Not tested		
Critically low		related bleeding requiring FC		both groups)	FC dose, grams to 24 hours	(n = 78) 4 (2, 6)	(n = 66) 4 (2, 4)	NR	No significant difference $p = 0.550$		
					PCC transfusion volume, international units to 24 hours ^c	(n = 78) O (0, 1200)	(n = 66) O (0, 1200)	NR	No significant difference p = 0.943		

Cl, confidence interval; Coh, cohort; FC, fibrinogen concentrate; FFP, fresh frozen plasma; IQR, interquartile range; NR, not reported; OR, odds ratio; PCC, prothrombin complex concentrate; PLT, platelets; RBC, red blood cells; SC, single centre; SR, systematic review

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

Length of stay

A summary of the evidence reported in the identified systematic reviews relating to length of stay (hospital and ICU) in patients with critical bleeding is presented in Table 4.50.

There was one cohort study (Innerhofer 2013) identified by 3 systematic reviews (Mengoli 2017, Aubron 2014, Lunde 2014) that was considered relevant to this review.

Hospital

One small cohort study (Innerhofer 2013) reported the median duration of hospital stay to be 29 days (IQR 16, 50) among 78 patients who received FFP which was longer than the median 24 days (IQR 12, 35) reported for the 66 patients who did not receive FFP. The difference was not statistically significant (p = 0.074).

Intensive care unit

One small cohort study (Innerhofer 2013) reported the median duration of ICU stay to be 14 days (IQR 7, 30) among 78 patients who received FFP which was longer than the median 12 days (IQR 6, 24) reported for the 66 patients who did not receive FFP. The difference was not statistically significant (p = 0.217).

Table 4.50 Results for FFP versus no FFP: Patients with critical bleeding – Length of stay

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. Of trials) included in analysis	population (Location)			FFP median (IQR)	No FFP median (IQR)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b		
Trauma setting	g									
Mengoli 2017	N = 144 (1 Coh)	Patients with	Trauma centre	FFP vs no FFP	LOS, In-patient days °	(n = 78)	(n = 66)	NR	No significant difference	
SR	Innerhofer 2013	severe trauma-	(SC, Austria)	(± FC ± PCC in both		29 (16-50)	24 (12-35)		<i>p</i> = 0.074	
Low		related bleeding		groups)	LOS, ICU days ^c	(n = 78)	(n = 66)	NR	No significant difference	
						14 (7-30)	12 (6-24)		<i>p</i> = 0.217	

Cl, confidence interval; Coh, cohort; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ICU, intensive care unit; IQR, interquartile range; LOS, length of stay; NR, not reported; PCC, prothrombin complex concentrate; SC, single centre; SR, systematic review

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

4.7.3.2 Cryoprecipitate

Mortality

A summary of the evidence reported in the identified systematic reviews relating to the effect of CRYO on mortality in patients with critical bleeding is presented in Table 4.51.

One systematic review (McQuilten 2018) reported the results of one RCT (Curry 2015) that contributed data relevant to critically bleeding patients in a trauma setting.

The RCT reported a lower rate of mortality among patients who received CRYO (2/20, 10.0%) compared with those who did not (6/21, 28.6%). The difference between treatment groups was not statistically significant (RR 0.35, 95% CI 0.08, 1.54; p = 0.14).

Figure 4.30 Forest plot of comparison: CRYO vs no CRYO (or varying administration of...), outcome: Mortality, latest timepoint.

	CRY	0	no CRYO (or varying)			Risk Ratio	Ris		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Ran	dom, 95% Cl	
Curry 2015 (RCT, trauma))	2	20	6	21	100.0%	0.35 [0.08, 1.54]			
Total (95% CI)		20		21	100.0%	0.35 [0.08, 1.54]			
Total events	2		6						
Heterogeneity: Not applicable Test for overall effect: Z = 1.3	9 (P = 0.1	6)					0.01 0.1 Favours CRYC	1 10 Favours no CRY	100 (O

Table 4.51 Results for CRYO versus no CRYO: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient population S	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis		(Location)			CRYO n/N (%)	No CRYO n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting										
McQuilten 2018	N = 41 (1 RCT)	Patients ≥ 16 years	Multi trauma	CRYO vs	Mortality, 28	2/20 (10)	6/21 (28.6)	RR 0.35 (0.08, 1.54)	No significant difference	
SR	Curry 2015	actively bleeding and	centres (UK)	standard	days				<i>p</i> = 0.14	
Moderate		required activation of		therapy (6 units						
quality		major haemorrhage		RBC vs 4 FFP)						
		protocol								

Cl, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; RBC, red blood cell; RCT, randomised controlled trial; RR, relative risk; SR, systematic review; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Morbidity

A summary of the evidence reported in the identified systematic reviews relating to morbidity in patients with critical bleeding is presented in Table 4.52.

One systematic review (McQuilten 2018) reported the results of one RCT (Curry 2015) that contributed data relevant to this outcome in critically bleeding patients in a trauma setting. The study was not sufficiently powered to detect important differences in event rates, therefore evidence for morbidity outcomes should be considered with caution

Thromboembolic events

One RCT (Curry 2015) reported no thromboembolic events among critically bleeding trauma patients who received CRYO compared with a total of 3 events in the placebo group (RR 0.15; 95% CI 0.01, 2.73; p = 0.20).

Specifically, a lower rate of deep vein thrombosis (DVT) was observed among patients who received CRYO (0/20, 0%) compared with those who did not (1/21, 4.8%) and a lower rate of pulmonary embolus (PE) was reported among patients who received CRYO (0/20, 0%) compared with those who did not (2/21, 9.5%).

The event rates for both outcomes were not significantly different (DVT: RR 0.35, 95% CI 0.02, 8.10; p = 0.51) and (PE: RR 0.21, 95% CI 0.01, 4.11; p = 0.30).

There were no events of myocardial infarction or stroke reported in the RCT.

Multiple organ failure

One RCT (Curry 2015) reported a higher rate of multiple organ failure among critically bleeding trauma patients who received CRYO (1/20, 5%) compared with those who did not (0/21, 0%), corresponding to a RR of 3.14 (95% CI 0.14, 72.92; p = 0.48).

Other adverse outcomes

One RCT (Curry 2015) reported a lower rate of ARDS among critically bleeding trauma patients who received CRYO (0/20, 0%) compared with those who did not (1/21, 4.8%); corresponding to a RR of 0.35 (95% CI 0.02, 8.10; p = 0.51). A higher rate of sepsis among patients who received CRYO (3/20, 15%) compared with those who did not (0/21, 0%) was also observed (RR 7.33, 95%CI 0.40, 133.57; p = 0.18).

Figure 4.31 Forest plot of comparison: CRYO vs no CRYO (or varying administration of...), outcome: Morbidity, thromboembolic events.

	CRY	0	no CRYO (or vary	/ing)		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% CI
2.2.1 Deep vein thrombosis							
Curry 2015 (RCT, trauma)) Subtotal (95% CI)	0	20 20	1	21 21	100.0% 100.0%	0.35 [0.02, 8.10] 0.35 [0.02, 8.10]	
Total events	0		1				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.66	6 (P = 0.5	1)					
2.2.2 Pulmonary embolus							_
Curry 2015 (RCT, trauma))	0	20	2	21	100.0%	0.21 [0.01, 4.11]	
Subtotal (95% CI)		20		21	100.0%	0.21 [0.01, 4.11]	
Total events	0		2				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 1.03$	3 (P = 0.3	0)					
2.2.3 Myocardial infarction							
Curry 2015 (RCT, trauma))	0	20	0	21		Not estimable	
Subtotal (95% CI)		20		21		Not estimable	
Total events	0		0				
Heterogeneity: Not applicable							
Test for overall effect: Not app	licable						
2.2.4 Stroke							
Curry 2015 (RCT, trauma))	0	20	0	21		Not estimable	
Subtotal (95% CI)		20		21		Not estimable	
Total events	0		0				
Heterogeneity: Not applicable							
Test for overall effect: Not app	licable						
							0.001 0.1 1 10 1000
T . ()	01.2	05 10					Favours CRYO Favours no CRYO
Test for subgroup differences:	$Chi^2 = 0.$	05, df =	$= 1 (P = 0.82), I^2 = 0^{\circ}$	%			

Figure 4.32 Forest plot of comparison: CRYO vs no CRYO (or varying administration of...), outcome: Morbidity, other.

	CRY	0	no CRYO (or vary	ying)		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	I M-H, Random, 95% CI
2.3.5 Multiorgan failure							
Curry 2015 (RCT, trauma))	1	20	0	21	100.0%	3.14 [0.14, 72.92]	
Subtotal (95% CI)		20		21	100.0%	3.14 [0.14, 72.92]	
Total events	1		0				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.7$	1 (P = 0.4	8)					
2.3.6 ARDS							_
Curry 2015 (RCT, trauma))	0	20	1	21	100.0%	0.35 [0.02, 8.10]	
Subtotal (95% CI)		20		21	100.0%	0.35 [0.02, 8.10]	
Total events	0		1				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.66	6 (P = 0.5	51)					
2.3.7 Sepsis							_
Curry 2015 (RCT, trauma))	3	20	0	21	100.0%	7.33 [0.40, 133.57]	
Subtotal (95% CI)		20		21	100.0%	7.33 [0.40, 133.57]	
Total events	3		0				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.35	5 (P = 0.1	8)					
							Favours CRYO Favours no CRYO
Test for subgroup differences:	Chi2 - 2	02 df -	2(P - 0.26) $12 - 1$	20/			

Test for subgroup differences: $Chi^2 = 2.03$, df = 2 (P = 0.36), $l^2 = 1.3\%$

Table 4.52 Results for CRYO versus no CRYO: Patients with critical bleeding – Morbidity

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis	population	(Location)			CRYO n/N (%)	No CRYO n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Trauma settin	g									
McQuilten 2018	n = 41 (1 RCT) Curry 2015	Patients ≥16 years actively bleeding and	Multiple trauma centres (UK)	CRYO vs standard	Deep vein thrombosis	0/20 (0)	1/21 (4.8)	RR 0.35 (0.02, 8.10)	NR	
SR		required activation of		therapy (6 U RBC	Pulmonary embolus	0/20 (0)	2/21 (9.5)	RR 0.21 (0.01, 4.11)	NR	
Moderate quality		protocol		VS 4 FFP)	Myocardial infarction	0/20 (0)	0/21 (0)	Not estimable	Not estimable	
					Stroke	0/20 (0)	0/21 (0)	Not estimable	Not estimable	
					MOF	1/20 (5)	0/21 (0)	RR 3.14 (0.14, 72.92)	NR	
			ARDS	0/20 (0)	1/21 (4.8)	RR 0.35 (0.02, 8.10)	NR			
			Sepsis	3/20 (15)	0/21 (0)	RR 7.33 (0.40, 133.57)	NR			

ARDS, acute respiratory distress syndrome; CI, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; MOF, multiorgan failure; NR, not reported; RBC, red blood cell; RCT, randomised controlled trial; RR, relative risk; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Transfusion volumes

A summary of the evidence reported in the identified systematic reviews relating to transfusion volumes in patients with critical bleeding is presented in Table 4.53 (RBC) and Table 4.54 (other blood component).

One systematic review (McQuilten 2018) reported the results of one RCT (Curry 2015) that contributed data relevant to this outcome in critically bleeding patients in a trauma setting.

Red blood cells

One small RCT (Curry 2015) reported no significant difference in the volume of RBC transfused up to 6 hours, 24 hours or 28 days among patients who received CRYO compared to those who did not. At 24-hours, participants in the control group had received a median (IQR) of 7 (6, 9) units of RBC compared to 8 (5,11) units given to those randomised to the CRYO group.

Other blood components

One small RCT (Curry 2015) reported no significant difference in the volume of FFP, PLT, or CRYO transfused up to 6 hours, 24 hours or 28 days among patients who received CRYO compared to those who did not.

At 24-hours, participants in the control group had received a median (IQR) of 6 (3, 8) units of FFP compared to 7 (4, 8) units given to those randomised to the CRYO group.

At 24-hours, participants in the control group had received a median (IQR) of 1 (1, 2) unit of PLT compared to 1 (0, 2) unit given to those randomised to the CRYO group.

At 24-hours, participants in the control group had received a median (IQR) of 2 (0, 2) unit of CRYO compared to 2 (2, 4) units given to those randomised to the CRYO group.

Table 4.53 Results for CRYO versus no CRYO: Patients with critical bleeding – RBC transfusion volume

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
design ^a <i>Trauma setting</i>	(no. Of trials) included in analysis	population	(Location)			CRYO median (IQR)	No CRYO median (IQR)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Trauma setting	I								
McQuilten 2018 SR <i>Moderate</i>	n = 41 (1 RCT) Curry 2015	RCT) Patients ≥16 years Curry 2015 actively bleeding and required activation of major baemorrhage	Multiple trauma centres (UK)	CRYO vs standard therapy (6 U RBC vs 4 FFP)	RBC transfusion volume, units to 6 hours RBC transfusion	(n = 20) 7 (4, 10) (n = 20)	(n = 21) 7 (4, 8) (n = 21)	Not estimable	No significant difference p = 0.49 No significant difference
		protocol	morrhage tocol		volume, units to 24 hours °	8 (5, 11)	7 (6, 9)		<i>p</i> = 0.83
					RBC transfusion volume, units to 28 days ^c	(n = 20) 9 (7, 15)	(n = 21) 8 (7, 11)	Not estimable	No significant difference <i>p</i> = 0.10

Cl, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; IQR, interquartile range; RBC, red blood cells; RCT, randomised controlled trial; SR, systematic review; UK, United Kingdom a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been

considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

Table 4.54	Results for CRYO versus no CRYO: Patients with critical bleeding	ng – Transfusion volume, other blood components
------------	--	---

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis	population	(Location)			CRYO median (IQR)	No CRYO median (IQR)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b	
Trauma setting	1									
McQuilten 2018 SR <i>Moderate</i>	n = 41 (1 RCT) Curry 2015	Patients ≥16 years actively bleeding and required	Multiple trauma centres (UK)	CRYO vs standard therapy (6 U RBC vs 4 FFP)	FFP transfusion volume, units to 6 hours	(n = 20) 7 (4, 8)	(n = 21) 5 (3, 8)	Not estimable	No significant difference <i>p</i> = 0.31	
quality		activation of major haemorrhage protocol			FFP transfusion volume, units to 24 hours ^c	(n = 20) 7 (4, 8)	(n = 21) 6 (3, 8)	Not estimable	No significant difference $p = 0.36$	
					FFP transfusion volume, units to 28 days ^c	(n = 20) 8 (4, 12)	(n = 21) 5 (3, 8)	Not estimable	No significant difference $p = 0.06$	
					PLT transfusion volume, units to 6 hours	(n = 20) 1 (0, 1)	(n =21) 1 (0, 1)	Not estimable	No significant difference p = 0.89	
					PLT transfusion volume, units to 24 hours ^c	(n = 20) 1 (0, 2)	(n = 21) 1 (1, 2)	Not estimable	No significant difference <i>p</i> = 0.56	
					PLT transfusion volume, units to 28 days ^c	(n = 20) 1 (0, 2)	(n = 21) 1 (1, 2)	Not estimable	No significant difference p = 0.82	
				CRYO transfusion volume, units to 6 hours	(n = 20) 2 (2, 4)	(n = 21) 2 (0, 2)	Not estimable	Favoured intervention p = 0.03		
					CRYO transfusion volume, units to 24 hours ^c	(n = 20) 2 (2, 4)	(n = 21) 2 (0, 2)	Not estimable	No significant difference $p = 0.23$	
					CRYO transfusion volume, units to 28 days ^c	(n = 20) 2 (2, 4)	(n = 21) 2 (0, 2)	Not estimable	No significant difference $p = 0.06$	

Cl, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; IQR, interquartile range; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; SR, systematic review; UK, United Kingdom a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been

considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

Length of stay

A summary of the evidence reported in the identified systematic reviews relating to LOS (hospital and ICU) in patients with critical bleeding is presented in Table 4.55.

One systematic review (McQuilten 2018) reported the results of one RCT (Curry 2015) that contributed data relevant to this outcome in critically bleeding patients in a trauma setting.

Hospital

One RCT (Curry 2015) reported the median (IQR) duration of hospital LOS to be 31 days (29, 33) among 20 patients who received CRYO compared to 30 days (22, 38) among the 21 patients who did not receive CRYO. The difference was not statistically significant (p = 0.66).

Intensive care unit

One RCT (Curry 2015) reported the median (IQR) duration of ICU LOS to be 11 days (5, 17) among 20 patients who received CRYO compared to 18 days (16, 20) among the 21 patients who did not receive CRYO. The difference was not statistically significant (p = 0.56).

Table 4.55 Results for CRYO versus no CRYO: Patients with critical bleeding – Length of stay

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results	No opvio		
Study designª	(no. Of trials) included in analysis		(Location)			CRYO Median (IQR)	No CRYO Median (IQR)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma settin	g								
McQuilten	n = 41 (1 RCT)	Patients ≥16 years	Multiple	CRYO vs standard	In-patient LOS, days	31 (29, 33)	30 (22, 38)	Not estimable	No significant difference
2018	Curry 2015	actively bleeding and	trauma centres	therapy					<i>p</i> = 0.66
SR		required activation of	(UK)	(6 U RBC vs 4 FFP)	ICU LOS, days	11 (5, 17)	18 (16, 20)	Not estimable	No significant difference
Moderate quality		major haemorrhage protocol							<i>p</i> = 0.56

Cl, confidence interval; CRYO, cryoprecipitate; FFP, fresh frozen plasma; ICU, intensive care unit; IQR, interquartile range; LOS, length of stay; RBC, red blood cells; RCT, randomised controlled trial; SR, systematic review; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

4.7.3.3 Platelets

No studies identified.

4.7.3.4 Fibrinogen concentrate

Mortality

A summary of the evidence reported in the identified systematic reviews relating to mortality (latest timepoint) in patients with critical bleeding is presented in Table 4.56.

There does not seem to be clear significant survival benefits for critically bleeding patients administered FC, but none of the RCTs and several nonrandomised cohort studies were sufficiently powered to detect differences in mortality.

Among critically bleeding trauma patients, a meta-analysis of data from the included RCTs (see Figure 4.33) showed the mortality rate (latest timepoint) among those who received FC (26/144, 18.1%) to be comparable to those who did not (25/139, 18.0%) with a RR of 1.12 observed (95% CI 0.53, 2.35; p = 0.77; random effects; $I^2 = 45\%$). Statistical heterogeneity was moderate.

Data from the included cohort studies (see Figure 4.33) suggests a non-significant association with higher mortality among trauma patients who received FC (131/615, 21.3%) compared with those who did not (152/1130, 13.5%) with the RR of 1.39 observed (95% CI 0.91, 2.13; p = 0.13; random effects; I² = 45%).

Among critically bleeding patients in the surgical setting, a meta-analysis of data from the included RCTs (see Figure 4.33) showed no significant difference in the rate of mortality (latest timepoint) between patients who received FC (4/177, 2.3%) compared to patients who did not (9/176, 5.1%) with a RR of 0.48 observed (95%Cl 0.08, 2.83; p = 0.42; random effects; $l^2 = 40\%$), noting the event rate was low across both treatment groups and statistical heterogeneity was moderate.

Data from the included cohort studies (see Figure 4.33) also suggested a non-significant association with higher mortality in patients who received FC (18/280, 6.4%) compared with those who did not (35/898, 3.9%), with a RR of 1.58 observed (95% CI 0.65, 3.85; p = 0.31; random effects; I² = 11%).

There were no deaths (up to 30 days) reported in the RCTs that examined the effect of FC on mortality in women with major postpartum haemorrhage (Collins 2017, Wikkelsø 2015).

Similarly, there were no deaths (up to 7 days) reported in the RCT that assessed the effect of FC on mortality in paediatric patients with diffuse bleeding after CPB.

Figure 4.33 Forest plot of comparison: FC vs no FC (or varying concentration of), outcome: Mortality, all-cause (latest timepoint)

Study cry bubycoup Events Total Events Total Wells M-H, Random, 95%, C1 M-H, Random, 95%, C1 Nascimento 2016 (RCT, trauma) 3 24 2 25 3.2% 1.56 (0.20, 8.55) Incendrei 2017 (RCT, trauma) 5 16 3 16 52% 1.87 (0.48, 6.83) Abbra 2016 (RCT, trauma) 10 24 7 24 9.8% 1.43 (0.65, 3.13) Subboal (95%, CL) 144 139 27.5% 1.12 (0.53, 2.35) 1.12 (0.53, 2.35) Total events 26 25 25 1.50 (0.28, 7.53) 1.50 (0.28, 7.53) Total events 28 25 1.32 (0.56, 1.31) 1.50 (0.28, 7.53) 1.50 (0.28, 7.53) Total events 28 20 2.51 (0.6, 1.50 (0.56) 1.50 (0.28, 7.53) 1.50 (0.28, 7.53) Insclaration events 18 2.08 (1.10, 6.56) 5.50 (0.10, 1.50 (0.56) 1.50 (0.26, 1.50 (0.56) Subchal (95% CL) 131 152 (0.56) (0.50 (0.56) 1.50 (0.52, 1.50 (0.5) 1.50 (0.52, 1.50 (0.5) Subchal (95% CL)		FC		No FC (or vary	/ing)		Risk Ratio	Risk Ratio
3.1.1 Trauma setting (RCTs) Naccimeta 2016 (RCT, trauma) 3 24 2 25 3.2% 1.56 [0.29, 8.55] Incerbidle 2017 (RCT, trauma) 5 60 2 44 3.5% 2.20 [0.45, 10.78] Locena 2020 (RCT, trauma) 3 30 11 30 5.8% 0.27 [10.8, 6.83] Atbair 2018 (RCT, trauma) 10 24 7 24 9.8% 1.43 [0.65, 5.13] Subtoal (95% CI) 144 139 27.5% 1.12 [0.53, 2.35] Total events 26 25 Heterogeneity: Tau ² = 0.32; Ch ² = 7.29, df = 4 (P = 0.12); P = 45% Test for overall effect Z = 0.29 (P = 0.77) 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Inockub 2017 (Coh, trauma) 17 115 6 109 8.4% 2.69 [1.10, 6.56] Schoch 12011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Inockub 2017 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Amsleg 2020 (Coh, trauma) 17 115 6 109 8.4% 2.09 [1.07, 407] Wafaisade 2013 (Coh, trauma) 23 108 11 108 11.6% 2.09 [1.07, 407] Wafaisade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 [0.86, 1.47] Subtoal (95% CI) 615 1130 52.3% Test for overall effect Z = 1.53 (P = 0.12); P = 45% Test for overall effect Z = 1.53 (P = 0.12); P = 45% Test for overall effect Z = 1.53 (P = 0.12); P = 45% Test for overall effect Z = 0.30 (P = 0.12); P = 40% Tanka 2014 (RCT, surgical) 1 29 4 22 2.1% 0.28 [0.02, 5.02.00] Rahe-Meyer 2016 (RCT, surgical) 1 78 5 74 2.1% 0.19 [0.02, 1.58] Subtoal (95% CI) 177 5 77 2 1.2% 0.37 [0.02, 6.71] Subtoal (95% CI) 20 0 10 0 5 Not estimable Bieleen 2017 (RCT, surgical) 0 10 0 5 Not estimable Bieleen 2017 (RCT, surgical) 0 10 0 5 Not estimable Bieleen 2017 (RCT, surgical) 0 10 0 5 Not estimable Bieleen 2017 (RCT, surgical) 1 224 33 881 13.6% 1.82 [1.04, 3.16] Subtoal (95% CI) 200 Ch, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Subtoal (95% CI) 200 Ch, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Subtoal (95% CI) 200 Ch, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Subtoal (95% CI) 200 Ch, surgical) 0 6 2 2 12 1.2% 0.37 [0.02, 6.71] Subtoal (95% CI) 200 Ch, surgical 10 6 0.2% 20 836 14.5% 1.58 [0.65, 3.58] Total events 18 5 35 Heterogeneity, Tau ² = 0.4(Ch	Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl
Naccimento 2016 (RCT, trauma) 3 24 2 25 3.2% 1.56 (0.29, 6.57) Innerhoffer 2017 (RCT, trauma) 5 60 2 44 3.5% 2.20 (0.48, 10.78) Lucena 2020 (RCT, trauma) 3 30 11 30 5.8% 0.27 (0.08, 0.88) Curry 2018 (RCT, trauma) 10 24 7 24 9.8% 1.43 (0.65, 3.13) Subtotal (8% CO) 144 7 139 27.5% 1.2 (0.53, 2.35) Total events 26 25 Heterogeneity: Tau ² = 0.32; $D^2 = 2.03$ d. 4 ($P = 0.12$); $P = 4.0\%$ Test for overall effect $Z = 0.29$ ($P = 0.77$) 3.1.2 Trauma setting (COh) Nenaeber 2011 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 118 3.3% 1.50 (0.28, 7.93) Inokuch 2017 (Coh, trauma) 3 18 2 119 52.3% 1.50 (0.28, 1.07) Subtotal (8% CO) 615 1130 52.3% 1.12 (0.66, 1.47) Subtotal (9% CO) 615 1130 52.3% 1.12 (0.66, 1.47) Subtotal (9% CO) 615 1130 52.3% 1.12 (0.68, 1.47) Tanaka 2014 (RCT, surgical) 1 77 176 5.4% 0.48 (0.08, 2.83] Heterogeneity: Tau ² = 0.10; Ch ² = 3.31, df = 2 (P = 0.12); P = 4.0% Test for overall effect. Z = 0.53 (P = 0.13) 3.1.3 Urgical setting (RCT) Test for overall effect. Z = 0.53 (P = 0.19); P = 4.0% Test for overall effect. Z = 0.53 (P = 0.42); P = 4.0%; Test for overall effect. Z = 0.50 (P = 0.42); P = 4.0%; Test for overall effect. Z = 0.50 (P = 0.42); P = 4.0%; Test for overall effect. Z = 0.50 (P = 0.42); P = 4.0%; Test for overall effect. Z = 0.50 (P = 0.42); P = 4.0%; Test for overall effect. Z = 0.50 (P = 0.42); 3.14.Supcial setting (COTh) Rahe-Meyer 2005a (Coh, surgical) 0 6 0 1 0 0 5 Net estimable Bile no 2017 (CD, surgical) 1 77 176 5.4% 0.48 [0.08, 2.83] Total events 18 (G COh) Rahe-Meyer 2005a (Coh, surgical) 0 6 10 0 5 Net estimable Bile no 2017 (CD, surgical) 1 8 2.64 33 881 13.6%; 1.52 [0.65, 3.85] Total events 18 2 0.3 881 13.6%; 1.52 [0.65, 3.85] Total events 18 2 0.3 881 13	3.1.1 Trauma setting (RCTs)							
Interchedre 2017 (RCT, trauma) 5 50 2 44 35% 2.20 [0.45, 10.78] Locera 2020 (RCT, trauma) 3 30 11 30 5.8% 0.27 [0.08, 0.88] Akbari 2018 (RCT, trauma) 10 2.4 7 24 9.8% 1.43 [0.68, 5.313] Subtotal (95% CI) 144 139 27.5% 1.12 [0.53, 2.35] Total events 26 25 Heterogeneity: Tau ² = 0.32; Ch ² = 7.29, df = 4 ($P = 0.12$); P = 45% Test for overall effect: Z = 0.29 ($P = 0.77$) 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Hochwich 2011 (Coh, trauma) 17 115 6 109 8.4% 2.26 [1.10, 6.56] Schoch 2011 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Schoch 2011 (Coh, trauma) 23 108 11 108 11.6% 1.20 [0.7, 407] Wafaisada 2013 (Coh, trauma) 23 108 11 108 11.6% 2.09 [1.07, 407] Wafaisada 2013 (Coh, trauma) 24 108 11 108 11.6% 2.09 [1.07, 407] Wafaisada 2013 (Coh, trauma) 13 152 Total events 13 1 52 Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 ($P = 0.12$); P = 45% Test for overall effect: Z = 1.53 ($P = 0.13$) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgica) 1 29 4 32 2.1% 0.28 [0.03, 2.33] Subtotal (95% CI) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.30 ($P = 0.42$) 3.1.4 Surgical setting (CCh) Rahe-Meyer 2013 (RCT, surgica) 1 7.8 5 74 2.1% 0.19 [0.03, 2.35] Subtotal (95% CI) 717 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.30 ($P = 0.42$) 3.1.4 Surgical setting (CCh) Rahe-Meyer 2003 (Coh, surgica) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2017 (RCT, surgica) 1 6 3 5 Heterogeneity: Tau ² = 0.19; P = 40% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.30 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.3	Nascimento 2016 (RCT, trauma)	3	24	2	25	3.2%	1.56 [0.29, 8.55]	
Lucena 2001 (RC1, trauma) 5 10 3 3 10 5.2% 1.12 (1.43 (0.48 , 0.88) Curry 2018 (RCT, trauma) 10 24 7 24 9.8% 1.43 (0.68 , 0.87) Curry 2018 (RCT, trauma) 10 24 7 24 9.8% 1.43 (0.68 , 0.87) Subtata (165% CI) 24 7 24 9.8% 1.42 (0.53 , 2.35] Total events 26 25 Heterogenety: Tau ² = 0.32; Ch ² = 7.29, df = 4 ($P = 0.12$); $P = 45\%$ Test for overall effect: 2 = 0.29 ($P = 0.77$) 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 7 115 6 109 8.4% 2.69 (1.10 , 6.56 , 3.47 , 1.29 (0.65 , 1.43) Subchat (165% CI) 244 7 3 244 9.8% 0.75 (0.24 , 1.68) Armskog 2020 (Coh, trauma) 23 108 11 108 11.8% 2.09 (1.10 , 6.56 , 1.47 , 1.39 (2.65 , 1.39 (0.65 , 1.37) Subchat (165% CI) 6 1130 52.3% 1.39 (0.91 , 2.13] Total events 13 1 152 Heterogenetity: Tau ² = 0.10; Ch ² = 7.31, df = 4 ($P = 0.12$); $P = 45\%$ Test for overall effect: 2 = 1.33 ($P = 0.13$) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgica) 1 29 4 32 2.1% 0.248 (0.03 , 2.33] Heterogenetity: Tau ² = 0.10; Ch ² = 7.31, df = 4 ($P = 0.12$); $P = 45\%$ Test for overall effect: 2 = 0.30 ($P = 0.42$) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2013 (RCT, surgica) 1 7.7 176 5.4% 0.48 (0.08 , 2.83] Total events 4 9 Heterogenetity: Tau ² = 0.07; Ch ² = 3.31, df = 2 ($P = 0.19$); $P = 400\%$ Test for overall effect: $Z = 0.30$ ($P = 0.42$) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2003 (Coh, surgica) 1 6 5 Not estimable Bilicen 2017 (RCT, surgica) 1 8 24 33 881 13.6\% 182 (1.04, 3.18] Subtatal (95% CI) 280 ($P = 0.42$) 3.1.4 Surgical setting (Coh) Test for overall effect: $Z = 1.31$ ($P = 0.22$); $P = 11\%$ Test for overall effect: $Z = 1.30$ ($P = 0.22$); $P = 11\%$ Test for overall effect: $Z = 1.30$ ($P = 0.22$); $P = 11\%$ Test for overall effect: $Z = 1.01$ ($P = 0.22$); $P = 11\%$ Test for overall effect: $Z = 1.01$ ($P = 0.22$); $P = 11\%$ Test for overall effect: $Z = 1.01$ ($P = 0.22$); $P = 1.05$ Total events 1 8 3.5% Test for overall effect: $Z = 1.01$ ($P = 0.22$); $P = 0.27$	Innerhofer 2017 (RCT, trauma)	5	50	2	44	3.5%	2.20 [0.45, 10.78]	
Adda 120 (KC, Hauma) 10 24 7 24 9.8% 14.3 (De5. 3.13) Subtoul (65% Cf) 144 139 27.5% 1.12 (D.53, 2.35) Total events 26 2 Heterogeneity: Tat ² = 0.32; Ch ² = 7.29, df = 4 ($P = 0.12$); P = 45% Test for overall effect: Z = 0.29 ($P = 0.7$) 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 (D.28, 7.93) Inokuchi 2017 (Coh, trauma) 17 115 6 109 8.4% 2.69 (1.10, 5.66) Schoch 2011 (Coh, trauma) 23 108 11 108 11.6% 2.09 (1.07, 4.07) Wafaisade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 (D.86, 1.47) Subtoat (65% Cf) 615 1130 52.3% 1.39 (D.91, 2.13) Total events 131 152 Heterogeneity: Tat ² = 0.12; P = 45% Test for overall effect: Z = 1.53 ($P = 0.13$): 3.1.3 Surgical setting (CGT) Tanaka 2014 (RCT ; surgical) 1 29 4 32 2.1% 0.28 (D.02, 1.59) Subtoat (65% Cf) 177 17 176 5.4% 0.48 (D.08, 2.83) Total events 4 9 Heterogeneity: Tat ² = 0.30; Ch ² = 3.31, df = 2 ($P = 0.19$); P = 40% Test for overall effect: Z = 0.80 ($P = 0.42$) 3.1.4 Surgical setting (CGT) Total events 4 9 Heterogeneity: Tat ² = 0.30; Ch ² = 3.31, df = 2 ($P = 0.19$); P = 40% Test for overall effect: Z = 0.80 ($P = 0.42$) 3.1.4 Surgical setting (CCh) Rahe-Meyer 20096 (Coh, surgical) 1 70 0 5 Not estimable Bilcen 2017 (Ch ² = 1.13, df = 1 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Test for Overall effect: Z = 1.01 ($P = 0.29$); P = 11% Tes	Lucena 2020 (RCT, trauma)	5	16	3	16	5.2%	1.67 [0.48, 5.83]	
Control 2017 (CC1, standing) 10 2^{-4} 1 2^{-4} 9 3^{-6} 1.15 (D05, 3.235) Total events 28 28 77.5% 1.12 (D.53, 2.35) Total events 28 28 77.5% 1.12 (D.53, 2.35) Total events 28 0.29 (P = 0.77) 3.1.2 Trauma setting (Coh) Nieraber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Inokuch 2017 (Coh, trauma) 6 80 60 001 9.5% 0.75 (D.34, 1.68] Amskog 2020 (Coh, trauma) 23 108 11 108 11.6% 2.09 [10.7, 4.07] Variatizade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 (D.66, 1.47] Subtotal (95% C) 615 1130 52.3% 1.39 [0.91, 2.13] Total events 131 152 Heterogeneity: Tav' = 0.10; Ch ² = 7.31; df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCT)s Total events 4 9 Heterogeneity: Tav' = 0.07; Ch ² = 3.31; df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.50 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2005 (Coh, surgical) 1 28 4 32 2.1% 0.29 [0.02, 2.159] Subtotal (95% Ci) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tav' = 0.29 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2005 (Coh, surgical) 1 0 0 5 Not estimable Bilecen 2017 (RCT, surgical) 1 2.26 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 1 8 264 33 881 13.6% 1.52 [1.04, 3.18] Subtotal (95% Ci) 280 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2005 (Coh, surgical) 1 8 264 33 881 13.6% 1.52 [1.04, 3.18] Subtotal (95% Ci) 280 (P = 0.42) 3.1.5 Distericic and maternity (RCT)s Total events 1 8 35 Heterogeneity: Tav' = 0.47 (Ch ² = 1.13, df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Distericic and maternity (RCTs) Total events 0 0 2 7 Not estimable Rahe-Meyer 2005 (Coh, surgical) 0 0 0 2 7 Not estimable Bilecen 2013 (Coh, surgical) 0 0 0 2 7 Not estimable Bilecen 2013 (Coh, surgical) 0 0 0 2 7 Not estimable Total events 1 8 35 Heterogeneity: Tav' = 0.47 (Ch ² = 1.13; df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Distericic and maternity (RCTs) Total events 0	Akban 2018 (RCT, trauma)	10	30	7	30	0.0%	0.27 [0.06, 0.66]	
Total events 26 25 25 Heterogeneity: Tau ² = 0.22 (Chi ² = 7.29, df = 4 (P = 0.12); P = 45% Test for overall effect $Z = 0.22 (P = 0.77)$ 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Inokuchi 2017 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Almskog 2020 (Coh, trauma) 23 108 11 108 11.6% 2.299 [1.07, 4.07] Wafaisade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 [0.86, 1.47] Subtotal (95% Ch) 615 1130 52.3% 1.39 [0.91, 2.13] Total events 1 31 152 Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.03, 2.33] Total events 0 17 (RCT, surgical) 1 29 4 32 2.1% 0.48 [0.08, 2.83] Total events 0 19 7 (Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Total events 0 7 (Ch, surgical) 1 70 5 74 2.1% 0.48 [0.08, 2.83] Total events 0 9.7 (Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.07; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 1.01 (P = 0.22); P = 11% Test for overall effect: Z = 1.01 (P = 0.21); P = 11% Test for overall effect: Z = 1.01 (P = 0.21); P = 11% Test for overall effect: Z = 1.01 (P = 0.21); P = 11% Test for overall effect: Z = 1.01 (P = 0.21); P = 11% Test for overall effect: Z = 0.01 (P = 0.21); P = 11% Test for overall effect: Z = 1.01 (P = 0.21	Subtotal (95% CI)	10	144	'	139	27.5%	1.12 [0.53, 2.35]	•
$ \begin{array}{c} \text{Test for overall effect: } 2 = 0.29 (P = 0.77) \\ \hline 3.1.2 Trauma setting (Coh) \\ \text{Nieraber 2011 (Coh, trauma)} & 3 & 18 & 2 & 18 & 3.3\% & 1.50 [0.28, 7.93] \\ \text{Inokubi 2017 (Coh, trauma)} & 17 & 115 & 6 & 109 & 8.4\% & 2.69 [1.10, 6.66] \\ \text{Schchd 2011 (Coh, trauma)} & 6 & 80 & 60 & 601 & 9.5\% & 0.75 [0.34, 1.68] \\ \text{Almskog 2020 (Coh, trauma)} & 22 & 108 & 11 & 108 & 11.8\% & 2.09 [1.07, 4.07] \\ \text{Variasiade 2013 (Coh, trauma)} & 22 & 294 & 73 & 294 & 19.6\% & 1.12 [0.86, 1.47] \\ \text{Subtotal (95% Ch)} & 615 & 1130 & 52.3\% & 1.39 [0.91, 2.13] \\ \text{Total events} & 131 & 152 \\ \text{Test for overall effect: } Z = 1.53 (P = 0.13) \\ \hline 3.1.3 \ \text{Surgical setting (RCTs)} \\ \text{Tanka 2014 (RCT, surgical)} & 1 & 29 & 4 & 32 & 2.1\% & 0.28 [0.03, 2.33] \\ \text{Total events} & 4 & 9 \\ \text{Heterogeneity: } Tal^2 = 0.16 (Ch)^2 = 3.31, df = 2 (P = 0.19); P = 40\% \\ \text{Test for overall effect: } Z = 0.80 (P = 0.42) \\ \hline 3.1.4 \ \text{Surgical setting (Coh)} \\ \text{Rahe-Meyer 2005 (Coh, surgical)} & 1 & 26 & 43 & 881 & 13.6\% & 1.58 [0.06, 2.8.5] \\ \text{Test for overall effect: } Z = 0.80 (P = 0.42) \\ \hline 3.14 \ \text{Surgical setting (Coh)} \\ \text{Rahe-Meyer 2009 (Coh, surgical)} & 1 & 62 & 43 & 881 & 13.8\% & 1.58 [0.65, 3.85] \\ \text{Test for overall effect: } Z = 1.01 (P = 0.29); P = 11\% \\ \text{Test for overall effect: } Z = 1.01 (P = 0.29); P = 11\% \\ \text{Test for overall effect: } Z = 1.01 (P = 0.29); P = 11\% \\ \text{Test for overall effect: } Z = 1.01 (P = 0.31) \\ \hline 3.15 \ \text{Dotsterics and maternity (RCTS)} \\ \hline \text{Cline 0.012 (RCT, surgical)} & 1 & 0 & 0 & 6 & 2 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 1 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 1 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 1 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ \hline \text{Cline 0.012 (RCT, surgical)} & 0 & 0 & 0 & 0 \\ $	Total events	26		25				T
Test for overall effect: $Z = 0.29 (P = 0.77)$ 3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 [0.28, 7.93] Inokuchi 2017 (Coh, trauma) 17 115 6 109 8.4% 2.69 [1.10, 6.56] Schochl 2011 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Amskog 2020 (Coh, trauma) 82 294 73 294 19.6% 1.12 [0.86, 1.47] Subtotal (9% Cl) 615 1130 52.3% 1.39 [0.91, 2.13] Total events 131 152 Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: $Z = 1.53 (P = 0.13)$ 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 29 4 32 2.1% 0.01 (0.25, 102.00] Rahe-Meyer 2013 (RCT, surgical) 1 77 5 74 2.1% 0.48 [0.08, 2.83] Rahe-Meyer 2013 (RCT, surgical) 1 77 5 74 2.1% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: $Z = 1.33$, $df = 2 (P = 0.19); P = 40\%$ Test for overall effect: $Z = 0.80 (P = 0.42)$ 3.1.4 Surgical setting (Coh) Rahe-Meyer 2008 (Coh, surgical) 0 6 2 112 1.2% 0.37 [0.02, 6.71] Bilcen 2017 (Ch ² , surgical) 1 8 264 33 881 13.6%, 1.82 [1.04, 3.18] Subtotal (9% Cl) 280 889 1.1.5% [0.56, 3.85] Total events 1 8 35 Heterogeneity: Tau ² = 0.11, Ch ² = -1.3, df = 1 (P = 0.29); P = 11% Test for overall effect: $Z = 1.01 (P = 0.31)$ 3.1.5 Dottevites and maternity (RCTs) Coline 0.207 (RCT, surgical) 0 0 2 9 0 0 27 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 75 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 89 1.1.5% [0.56, 3.85] Total events 1 8 35 Heterogeneity: Tau ² = 0.11, Ch ² = -1.3, df = 1 (P = 0.29); P = 11% Test for overall effect: $Z = 1.01 (P = 0.31)$ 3.1.5 Dottevites and maternity (RCTs) Celline 0.207 (RCT surgical) 0 0 0 2 7 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 2 7 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 2 7 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 2 7 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 2 7 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 2 7 Not estimable	Heterogeneity: Tau ² = 0.32; Chi ² = 7.	29, df = 4	(P = 0.1	12); l ² = 45%				
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Test for overall effect: Z = 0.29 (P = 0	0.77)						
3.1.2 Trauma setting (Coh) Nienaber 2011 (Coh, trauma) 17 115 6 109 8.4% 2.69[1.07, 4.07] Schoch 2011 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Schoch 2011 (Coh, trauma) 23 108 11 108 11.6% 2.09 [1.07, 4.07] Wataisade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 [0.86, 1.47] Subtotal (95% Ch) 615 1130 52.3% 1.39 [0.91, 2.13] Heterogeneity: Tau ² - 0.10; PT - 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.02, 2.53] Subtotal (95% Ch) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2008 (Coh, surgical) 0 10 0 5 Not estimable Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2008 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2017 (Ch, surgical) 1 8 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Ch) 1200 808 (1.8% 1.58 [0.65, 3.85] Total events 16 35 Heterogeneity: Tau ² = 0.17; Ch ² = 3.11, df = 2 (P = 0.29); P = 11%, Test for overall effect: Z = 1.01 (P = 0.29); P = 11%, Test for overall effect: Z = 1.01 (P = 0.29); P = 11%, Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Dotterics and maternity (RCTs) Colling 0.2017 (CT, burgical) 0 0 0 74 Not estimable Deterogeneity: Tau ² = 0.17; Ch ² = 3.13								
Nenaber 2011 (Coh, trauma) 3 18 2 18 3.3% 1.50 (0.28, 7.93) Incluchi 2017 (Coh, trauma) 17 115 6 109 8.4% 2.69 [1.10, 6.56] Schochl 2011 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Almskog 2020 (Coh, trauma) 23 108 11 108 11.6% 2.09 [1.07, 407] Variatsade 2013 (Coh, trauma) 82 294 73 294 19.6% 2.09 [1.07, 407] Subtotal (95% CI) 615 1130 52.3% 1.39 [0.51, 2.13] Total events 131 152 Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCTs) Total events 1 2 2 60 0 60 1.1% 5.00 [2.55, 102.00] Rahe-Meyer 2016 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.03, 2.33] Rahe-Meyer 2016 (RCT, surgical) 1 78 5 74 2.1% 0.19 [0.02, 1.59] Subtotal (95% CI) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 79 Heterogeneity: Tau ² = 0.37; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2008 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 1 8 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.15 Obstetrics and maternity (RCTS) Cellera 2017 (PCT, surgical) 0 0 2% Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 2% Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 0 0 0 5 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 6 2 12 1.2% 0.37 (0.02, 6.71] Bilicen 2013 (CDT, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.15 Obstetrics and maternity (RCTS)	3.1.2 Trauma setting (Coh)							
Inokuchu 2017 (Coh, trauma) 17 115 b 109 8.4% 2.69 [1.0, 5.56] Schoch 2017 (Coh, trauma) 6 80 60 601 9.5% 0.75 [0.34, 1.68] Almskog 2020 (Coh, trauma) 23 108 11 108 11.6% 2.09 [1.07, 4.07] Wafaisade 2013 (Coh, trauma) 82 294 73 294 19.6% 1.12 [0.86, 1.12 [0.86, 1.12] Subtotal (95% CI) 615 1130 52.3% 1.139 [0.91, 2.13] Total events 131 152 Heterogeneity. Tau ² e.0.10, Chi ² = 7.31, df = 4 ($P = 0.12$); $P = 45\%$ Test for overall effect: Z = 1.53 ($P = 0.13$) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 0 10 0 10 Not estimable Bilecen 2017 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.03, 2.33] Rahe-Meyer 2016 (RCT, surgical) 1 78 5 74 2.1% 0.19 [0.02, 1.59] Subtotal (95% CI) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity. Tau ² = 0.97, Chi ² = 3.31, df = 2 ($P = 0.19$); $P = 40\%$ Test for overall effect: Z = 0.80 ($P = 0.42$) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2008 (Coh, surgical) 0 6 2 112 1.2% 0.37 (0.02, 6.71] Bilicen 2013 (Cch, surgical) 1 8 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 5 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 ($P = 0.29$); $P = 11\%$ Test for overall effect: Z = 1.01 ($P = 0.31$) 3.1.5 Dotsetrics and maternity (RCTS) Collers 2014 (PCT, barterion) 0 0 28 0 27 Not estimable Rahe-Meyer 2008 (Coh, surgical) 0 10 0 75 Not estimable Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 5 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 ($P = 0.29$); $P = 11\%$ Test for overall effect: Z = 1.01 ($P = 0.31$) 3.1.5 Dotsetrics and maternity (RCTS)	Nienaber 2011 (Coh, trauma)	3	18	2	18	3.3%	1.50 [0.28, 7.93]	
School and Tricola, addition 2011 (Columnation 2011) Columnation 2011 (Columnation 2011)	Inokuchi 2017 (Con, trauma)	17	115	60	109	8.4%	2.69 [1.10, 6.56]	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Almskog 2020 (Cob. trauma)	23	108	60 11	108	9.5%	2.09 [1.07 4.07]	·
Total events 131 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Wafaisade 2013 (Coh. trauma)	82	294	73	294	19.6%	1 12 [0 86 1 47]	- - -
Total events 131 152 Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 0 10 Not estimable Bilecen 2017 (RCT, surgical) 1 2 60 0 1.1% 5.00 (0.25, 102.00) Rahe-Meyer 2016 (RCT, surgical) 1 2.9 4 32 2.1% 0.28 (0.03, 2.33) Subtact (85% CI) 1 177 176 5.4% 0.48 (0.08, 2.83) Total events 4 9 Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 (0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 38 881 13.6% 1.82 [1.04, 3.18] Subtact (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); P = 11%<	Subtotal (95% CI)	02	615		1130	52.3%	1.39 [0.91, 2.13]	◆
Heterogeneity: Tau ² = 0.10; Ch ² = 7.31, df = 4 (P = 0.12); P = 45% Test for overall effect: Z = 1.53 (P = 0.13) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 0 0 10 0 10 Not estimable Bilicen 2017 (RCT, surgical) 2 60 0 60 1.1% 5.00 [0.25, 102.00] Rahe-Meyer 2013 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.03, 2.33] Rahe-Meyer 2016 (RCT, surgical) 1 78 5 74 2.1% 0.19 [0.02, 1.59] Subtotal (95% Cl) 177 T 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) Celline 0214 (QfCT, exterimentic) 0 0 0 27 Not estimable	Total events	131		152				
Test for overall effect: $Z = 1.53$ (P = 0.13) 3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 1 0 10 0 10 Not estimable Bilecen 2017 (RCT, surgical) 2 60 0 60 1.1% 5.00 [0.25, 102.00] Rahe-Meyer 2013 (RCT, surgical) 1 29 4 32 2.1% 0.28 [0.32, 2.33] Rahe-Meyer 2013 (RCT, surgical) 1 78 5 74 2.1% 0.19 [0.02, 1.59] Subtotal (95% Cl) 1 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.37; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: $Z = 0.80$ (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 1 8 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 1 8 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); P = 11% Test for overall effect: $Z = 1.01$ (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) Celline 021/2 (CoT e heating) 0 0 - 28 0 0 77 Net estimable	Heterogeneity: Tau ² = 0.10; Chi ² = 7.	31, df = 4	(P = 0.1	12); l² = 45%				
3.1.3 Surgical setting (RCTs) Tanaka 2014 (RCT, surgical) 0 10 Not estimable Bilecen 2017 (RCT, surgical) 2 60 0 60 1.1% $5.00 [0.25, 102.00]$ Rahe-Meyer 2013 (RCT, surgical) 1 29 4 32 2.1% $0.20 (0.3, 2.33]$ Rahe-Meyer 2013 (RCT, surgical) 1 78 5 74 2.1% $0.19 [0.02, 1.59]$ Subtotal (95% CI) 1777 176 5.4% $0.48 [0.08, 2.83]$ Otal events 4 9 Heterogeneity: Tau ² = 0.97; Chi ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% $0.37 [0.02, 6.71]$ Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% $0.37 [0.02, 6.5, 3.85]$ Subtotal (95% CI) 280 898 14.8% $1.58 [0.65, 3.85]$ Subtotal (95% CI) 280 898 14.8% $1.58 [0.65, 3.85]$ Total events 18 35 5 5	Test for overall effect: $Z = 1.53$ (P = 0	0.13)						
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3.1.3 Surgical setting (RCTe)							
The fact length of the second	Tanaka 2014 (RCT surgical) (1)	0	10	0	10		Not estimable	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Bilecen 2017 (RCT, surgical)	2	60	0	60	1 1%	5 00 [0 25 102 00]	
Rahe-Meyer 2016 (RCT, surgical) 1 78 5 74 2.1% $0.19 [0.02, 1.59]$ Subtotal (95% CI) 177 176 5.4% $0.48 [0.08, 2.83]$ Total events 4 9 Heterogeneity: Tau ² = 0.97; Chi ² = 3.31, df = 2 (P = 0.19); l ² = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009b (Coh, surgical) 0 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 2.64 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% CI) 280 898 1.4.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 (P = 0.29); l ² = 11% 1.50 [0.55, 3.85] 3.1.5 Obstetrics and maternity (RCTs) Coline 0.012 (PGT destriction) 0 28	Rahe-Mever 2013 (RCT, surgical)	1	29	4	32	2.1%	0.28 [0.03, 2.33]	
Subtotal (95% Cl) 177 176 5.4% 0.48 [0.08, 2.83] Total events 4 9 Heterogeneity: Tau ² = 0.37; Ch ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilder 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13; df = 1 (P = 0.29); P = 11% Test for overall effect: Z = 1.01 (P = 0.31) 31.5 Obstetrics and maternity (RCTs) Colling 012 (PG C) ebstetrics 0 27 Not estimption	Rahe-Meyer 2016 (RCT, surgical)	1	78	5	74	2.1%	0.19 [0.02, 1.59]	
Total events 4 9 Heterogeneity: Tau ² = 0.97; Chi ² = 3.31, df = 2 (P = 0.19); P = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009b (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Image: Comparison of the tau of the tau of tau o	Subtotal (95% CI)		177		176	5.4%	0.48 [0.08, 2.83]	
Heterogeneity: Tau ² = 0.97; Ch ² = 3.31, df = 2 (P = 0.19); l ² = 40% Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Ch ² = 1.13, df = 1 (P = 0.29); l ² = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs)	Total events	4		9				
Test for overall effect: Z = 0.80 (P = 0.42) 3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau² = 0.14; Chi² = 1.13, df = 1 (P = 0.29); I² = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) 0 28 0 27	Heterogeneity: Tau ² = 0.97; Chi ² = 3.	31, df = 2	(P = 0.1	19); l ² = 40%				
3.1.4 Surgical setting (Coh) Rahe-Meyer 2009a (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau² = 0.14; Chi² = 1.13, df = 1 (P = 0.29); P = 11% Total events 1.50 Stetrics and maternity (RCTs) Solitos (2/4 CC) textution 0 -2% 0 -2% Not estimable	Test for overall effect: $Z = 0.80$ (P = 0	0.42)						
Rahe-Meyer 2009a (Coh, surgical) 0 10 0 5 Not estimable Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau² = 0.14; Chi² = 1.13, df = 1 (P = 0.29); P = 11% Total events 1.50 Stottrics and maternity (RCTs) Solitos (21/COT ebettrics) 0 -28 0 27 Not estimable	3.1.4 Surgical setting (Coh)							
Rahe-Meyer 2009b (Coh, surgical) 0 6 2 12 1.2% 0.37 [0.02, 6.71] Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% Cl) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 (P = 0.29); l ² = 11% 7 Test for overall effect: Z = 1.01 (P = 0.31) 31.5 Obstetrics and maternity (RCTs)	Rahe-Mever 2009a (Coh. surgical)	0	10	0	5		Not estimable	
Bilicen 2013 (Coh, surgical) 18 264 33 881 13.6% 1.82 [1.04, 3.18] Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau'z = 0.14; Chi² = 1.13, df = 1 (P = 0.29); l² = 11% 158 Test for overall effect: Z = 1.01 (P = 0.31) 31.5 Obstetrics and maternity (RCTs) Celling 012/4 (CFC) relations 0 27	Rahe-Meyer 2009b (Coh, surgical)	0	6	2	12	1.2%	0.37 [0.02, 6.71]	
Subtotal (95% CI) 280 898 14.8% 1.58 [0.65, 3.85] Total events 18 35 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 (P = 0.29); P = 11% 1 Test for overall effect: Z = 1.01 (P = 0.31)	Bilicen 2013 (Coh, surgical)	18	264	33	881	13.6%	1.82 [1.04, 3.18]	
Total events 18 35 Heterogeneity: Tau ² = 0.14; Chi ² = 1.13, df = 1 (P = 0.29); l ² = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) Colline 01/2 (PCT) estimate	Subtotal (95% CI)		280		898	14.8%	1.58 [0.65, 3.85]	-
Heterogeneity: Tau ² = 0.14, Chi ² = 1.13, df = 1 (P = 0.29); l ² = 11% Test for overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) Colling 0017 (PCT extetrice) = 0 = 28 = 0 = 27 = 0 = 27	Total events	18		35				
1est tor overall effect: Z = 1.01 (P = 0.31) 3.1.5 Obstetrics and maternity (RCTs) Colling 2017 (PCT, obstetrics) 0 27	Heterogeneity: $Tau^2 = 0.14$; $Chi^2 = 1$.	13, df = 1	(P = 0.2	29); l² = 11%				
3.1.5 Obstetrics and maternity (RCTs)	Test for overall effect: $Z = 1.01$ (P = 0	J.31)						
Colline 2017 (PCT obstatring) 0 29 0 27 Not astimable	3.1.5 Obstetrics and maternity (RC	Ts)						
	Collins 2017 (RCT, obstetrics)	0	28	0	27		Not estimable	
Wikkelso 2015 (RCT, obstetrics) 0 123 0 121 Not estimable	Wikkelso 2015 (RCT, obstetrics)	0	123	0	121		Not estimable	
Subtotal (95% Cl) 151 148 Not estimable	Subtotal (95% CI)		151		148		Not estimable	
Total events 0 0	Total events	0		0				
Heterogeneity: Not applicable	Heterogeneity: Not applicable							
Test for overall effect: Not applicable	Test for overall effect: Not applicable							
3.1.6 Pediatrics (RCTs)	3.1.6 Pediatrics (RCTs)							
Galas 2014 (RCT, paediatrics) (2) 0 30 0 33 Not estimable	Galas 2014 (RCT. paediatrics) (2)	0	30	0	33		Not estimable	
Subtotal (95% Cl) 30 33 Not estimable	Subtotal (95% CI)		30		33		Not estimable	
Total events 0 0	Total events	0		0				
Heterogeneity: Not applicable	Heterogeneity: Not applicable							
Test for overall effect: Not applicable	Test for overall effect: Not applicable							
Total (95% Cl) 1397 2524 100.0% 1.26 (0.91.1.75)	Total (95% CI)		1397		2524	100.0%	1.26 [0.91, 1.75]	•
Total events 179 221	Total events	179		221			·	*
Heterogeneity: Tau ² = 0.12; Chi ² = 22.85, df = 14 (P = 0.06); l ² = 39%	Heterogeneity: Tau ² = 0.12; Chi ² = 22	2.85, df = 1	4 (P =	0.06); l ² = 39%				
Test for overall effect: Z = 1.39 (P = 0.17) 0.005 0.1 1 0 200 Eavourse fibriogen concentrate	Test for overall effect: Z = 1.39 (P = 0	0.17)	`					U.UU5 U.1 1 10 200
Test for subgroup differences: Chi ² = 1.65, df = 3 (P = 0.65), l ² = 0%	Test for subgroup differences: Chi ² =	1.65, df =	3 (P =	0.65), l ² = 0%				avours normogen concentrate in avours no normogen concentrate
Footnotes	Footnotes							

(2) FC vs CRYO

Table 4.56 Results for FC versus no FC: Patients with critical bleeding - Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis	population	(Location)			FC n/N (%)	No FC n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Trauma setting		'								
Fabes 2018 SR High quality	N = 97 (2 RCTs) Curry 2018 Nascimento 2016	Adult trauma patients (≥16 years, blunt or penetrating) with active bleeding and in/at risk of haemorrhagic shock.	Trauma, MC (UK, Canada)	FC vs placebo (normal saline)	Mortality, 28 days	13/48 (27) 10/24 (42) 3/24 (12.5)	9/49 (18) 7/24 (29) 2/25 (8)	RR 1.46 (0.71, 2.99) RR 1.43 (0.65, 3.13) RR 1.56 (0.29, 8.55)	No significant difference p = 0.30 No significant heterogeneity $l^2 = 0\%$	
	N = 94 (1 RCT) Innerhofer 2017	Adults with ISS > 15 and clinical signs or risk of substantial haemorrhage	Trauma (Austria)	FC vs FFP	Mortality, 30 days	5/50 (10)	2/44 (4.5)	OR 2.20 (0.45, 10.78)	NR	
Stabler 2020 SR	N = 216 (1 Coh) Almskog 2020	Adult patients with major blunt trauma	Trauma (Sweden)	FC vs no FC	Mortality, 24 hours	7/108	1/108	NR	<i>p</i> = 0.494	
Moderate quality	N = 896 (2 RCTs, 2 Coh) RCTs Akbari 2018 Lucena 2020 Observational Wafaisade 2013 Almskog 2020	Adult patients with major blunt trauma	Trauma (Iran, Brazil, Germany, Sweden)	FC vs no FC	Mortality, latest timepoint	NR 3/30 (10) 5/16 (31.2) 82/294 (27.9) 23/108 (21.3)	NR 11/30 (10) 3/16 (31.2) 73/294 (24.8) 11/108 (10.2)	NR NR NR NR	NR <i>p</i> = 0.029 <i>p</i> = 0.456 <i>p</i> = 0.40 <i>p</i> = 0.859	
	N = 224 (1 Coh) Inokuchi 2017	Patients with pelvic fractures from blunt trauma requiring activation of MTP	Trauma (Japan)	FC + FFP vs FFP	Mortality, 28 days	17/115 (15)	6/109 (6)	NR	Favours no FC p <0.05	
	N = 717 (2 Coh) Schöchl 2011 Nienaber 2011	Adult trauma patients	Trauma centre (Germany, Austria)	FC vs FFP	Mortality, overall in-hospital	6/80 (7.5) 3/18 (16.7)	60/601 (10) 2/18 (11.1)	NR	No significant difference p = 0.69 p = 0.50	
Aubron 2014 SR <i>Critically low</i> <i>quality</i>	N = 588 (1 Coh) Wafaisade 2013	Trauma patients \geq 16 years of age with ISS \geq 16 + at least 1 RBC + TASH score \geq 9	Trauma (MC, multicountry)	FC vs No FC	Mortality, 24 hours	41/294 (13.9) ^c	54/294 (18.4)°	RR 1.12 (0.86, 1.46)	No significant difference p = 0.15 °	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results						
Study (no. Of trials design ^a included in analysis		population	(Location)			FC n/N (%)	No FC n/N (%)		Risk estimate (95% Cl)		Statistical significance p-value Heterogeneity ^b	
Surgical setting												
Fabes 2018 SR High	N = 120 (1 RCT) Bilecen 2017	Adults >18 years undergoing elective high-risk cardiac surgery	Surgery (SC, Netherlands)	FC vs placebo (albumin in normal saline)	Mortality, 30 days	2/60 (3.3)		0/60		RR 5.00 (0.25, 102.00)	No significant difference <i>p</i> = 0.30	
	N = 213 (2 RCTs) Rahe-Meyer 2013 Rahe-Meyer 2016	Aged ≥18 with elective aortic valve replacement surgery	Surgery (Germany)	FC vs 0.9% saline	Mortality, 46 days	2/107 (1.9)	1/29 (3.4) 1/78 (1.3)	9/106 (8.5)	4/32 (12.5) 5/74 (6.8)	RR 0.23 (0.05,1.01) 0.28 (0.03, 2.33) 0.19 (0.02, 1.59)	No significant difference p = 0.052 No significant heterogeneity $l^2 = 0.0\%$	
	N = 20 (1 RCT) Tanaka 2014	Patients undergoing elective CPB procedures	Surgery (US)	FC vs PLT	Mortality, 28 days	0/10		0/10		NR	NR	
Lunde 2014 SR <i>Critically low</i>	N = 1075 (1 Coh) Bilecen 2013	Patients undergoing surgery	Surgery (Netherlands)	FC vs No FC	Mortality, 30 days ^c	18/264 (7)		33/811 (4)		RR 0.96 (0.48- 1.92)	No significant difference p = 0.07	
Warmuth 2012 SR <i>Low</i>	N = 33 (2 Coh) Rahe-Meyer 2009a Rahe-Meyer 2009b	Adult patients undergoing surgery with major haemorrhage	Surgery (Germany)	FC vs standard infusion (FFP + PLT)	Mortality, 30 day	0/16	0/10 0/6	2/17 (11.8)	0/5 2/12 (17)	NR	NR	
Paediatrics, surg	gical setting											
Fabes 2018 SR High	N = 63 (1 RCT) Galas 2014	Patients <15 years undergoing CPB, intraoperative bleeding and hypofibrinogenemia	Paediatric surgery (Brazil)	FC vs CRYO	Mortality, 7 days	0/30		O/33		NR	NR	
Obstetrics and maternity setting												
Zaidi 2020 SR <i>Low</i>	N = 299 (2 RCTs) Collins 2017 Wikkelsø 2015	Women with major PPH	Obstetrics (Denmark and UK)	FC vs saline	Mortality, 30 days	O/151	0/28 0/123	0/148	0/27 0/121	NR	NR	

Cl, confidence interval; Coh, cohort; CPB, cardiopulmonary bypass; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ISS, injury severity score; MC, multicentre; NR, not reported; OR, odds ratio; PLT, platelets; PPH, postpartum haemorrhage; RCT, randomised controlled trial; RR, relative risk; SC, single centre; SR, systematic review; TASH, trauma associated severe haemorrhage; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

Morbidity

A summary of the evidence reported in the identified systematic reviews relating to morbidity (thromboembolic events, multiple organ failure, ARDs) in patients with critical bleeding is presented in Table 4.57 and Table 4.58.

Thromboembolic events

Among patients with critical bleeding in the trauma setting, a meta-analysis of data from 4 RCTs (see Figure 4.34) showed that the rate of thromboembolic events was comparable between patients who received FC (12/107, 11.2%) and those who did not (12/103, 11.7%). This corresponds to a RR of 0.90 (95% CI 0.42, 1.91; p = 0.78; random effects; $l^2 = 0$ %).

Data from the cohort studies (see Figure 4.34) also suggested no significant association with thromboembolic events among patients who received FC (53/511, 10.4%) compared with those who did not (49/517, 9.5%). This corresponds to a RR of 1.26 (95% CI 0.64, 2.49; p = 0.51; random effects; $I^2 = 58\%$), noting there was substantial statistical heterogeneity.

Among patients with critical bleeding in the surgical setting (see Figure 4.34) the rate of thromboembolic events was higher in patients who received FC (8/99, 8.0%) compared with those who did not (4/102, 3.9%) but the difference was not statistically significant (RR 2.03; 95% CI 0.63, 6.58; p = 0.24; random effects; $I^2 = 0$ %). It is noted that the evidence for thromboembolic events was limited by small patient numbers, with the included studies not sufficiently powered to detect important differences in event rates.

Among women with major postpartum haemorrhage (see Figure 4.34), the rate of thromboembolic events was comparable between patients who received FC (1/151, 0.7%) and those who did not (1/148, 0.7%); corresponding to a RR of 0.96 (95% CI 0.06, 14.65; p = 0.98; random effects; $I^2 =$ not applicable). The RCTs were small and not sufficiently powered to detect this outcome with one study (Wikkelsø 2015) reporting no thromboembolic events.

In paediatric patients with diffuse bleeding after CPB, a lower rate of thromboembolic events was reported among those who received FC (2/30, 6.7%) compared with those who did not (5/33, 12.2%) but the difference was not statistically significant (RR 0.44; 95% CI 0.09, 2.10; p = 0.3; random effects; $I^2 =$ not applicable).

Multiple organ failure

Three RCTs and 3 cohort studies were identified in the included systematic reviews that reported on the outcome of multiple organ failure (MOF) in the trauma setting.

A meta-analysis of data from the RCTs (see Figure 4.35) showed that the rate of MOF was lower among patients who received FC (29/97, 30%) compared with those who did not (38/98, 38.8%), but the difference did not reach statistical significance (RR 0.74; 95% CI 0.53, 1.03; p = 0.07; random effects; $l^2 = 0$ %).

Data from the cohort studies showed no significant difference in MOF among patients who received FC (184/420, 43.8%) compared with those who did not (156/420, 37.1%), corresponding to a RR of 0.70 (95%CI 0.21, 2.36; p = 0.57; random effects; $I^2 = 73\%$), noting the heterogeneity was substantial.

Acute respiratory distress syndrome

One RCT (Nascimento 2016) reported a lower event rate of ARDS among patients who received FC (0/21, 0%) compared with patients who did not receive FC (2/24, 8.3%), but the sample size was small and therefore no further analysis was performed.

Figure 4.34 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Morbidity, thromboembolic events

	FC	C No FC (or varying)			Risk Ratio	Risk Ratio			
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% Cl		
3.2.1 Trauma setting (RCTs)									
Lucena 2020 (RCT, trauma)	0	16	0	16		Not estimable			
Nascimento 2016 (RCT, trauma)	2	21	1	24	1.7%	2.29 [0.22, 23.44]			
Curry 2018 (RCT, trauma)	3	20	2	19	3.2%	1.43 [0.27, 7.61]			
Innerhofer 2017 (RCT, trauma)	7	50	9	44	11.1%	0.68 [0.28, 1.68]			
Tatal events	10	107	10	103	10.0%	0.90 [0.42, 1.91]			
Hotorogonoitur Tou2 – 0.00: Chi2 – 1.	12 26 df _ 2	(P - 0	1Z 52):12 - 09/						
Therefore the trace $z = 0.00$, $c_{11} = 1.20$, $c_{12} = 1.20$, $c_{13} = 2.17 = 0.53$; $t^2 = 0.\%$ Test for overall effect: $Z = 0.28$ (P = 0.78)									
3.2.2 Trauma setting (Coh)									
Almskog 2020 (Coh, trauma)	5	108	3	108	4.6%	1.67 [0.41, 6.80]			
Wafaisade 2013 (Coh, trauma)	20	294	10	294	16.4%	2.00 [0.95, 4.20]			
Inokuchi 2017 (Coh, trauma)	28	109	36	115	51.5%	0.82 [0.54, 1.25]			
Subtotal (95% CI)		511		517	72.5%	1.26 [0.64, 2.49]	-		
Total events	53		49						
Heterogeneity: Tau ² = 0.20; Chi ² = 4.77, df = 2 (P = 0.09); l ² = 58% Test for overall effect: Z = 0.67 (P = 0.51)									
3.2.3 Surgical setting (RCTs)									
Tanaka 2014 (RCT, surgical) (1)	0	10	0	10		Not estimable			
Rahe-Meyer 2013 (RCT, surgical)	1	29	1	32	1.2%	1.10 [0.07, 16.85]			
Bilecen 2017 (RCT, surgical)	7	60	3	60	5.3%	2.33 [0.63, 8.60]			
Subtotal (95% CI)		99		102	6.5%	2.03 [0.63, 6.58]			
Total events	8		4						
Heterogeneity: Tau ² = 0.00; Chi ² = 0.2	24, df = 1	(P = 0.	63); l² = 0%						
Test for overall effect: $Z = 1.18$ (P = 0	.24)								
3.2.4 Obstetrics and maternity (RC	Ts)								
Wikkelso 2015 (RCT, obstetrics)	0	123	0	121		Not estimable			
Collins 2017 (RCT, obstetrics)	1	28	1	27	1.2%	0.96 [0.06, 14.65]			
Subtotal (95% CI)		151		148	1.2%	0.96 [0.06, 14.65]			
Total events	1		1						
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.03 (P = 0	.98)								
3.2.5 Paediatric setting (RCTs)									
Galas 2014 (RCT, paediatrics) (2)	2	30	5	33	3.7%	0.44 [0.09, 2 10]			
Subtotal (95% CI)	-	30	0	33	3.7%	0.44 [0.09, 2.10]			
Total events	2		5						
Heterogeneity: Not applicable									
Test for overall effect: $Z = 1.03$ (P = 0	.30)								
Total (95% CI)		898		903	100.0%	1.03 [0.77, 1.40]	➡		
Total events	76		71						
Heterogeneity: Tau ² = 0.00; Chi ² = 8.8	80, df = 9	(P = 0.	46); l ² = 0%						
Test for overall effect: Z = 0.22 (P = 0	.83)						Eavours fibringen concentrate Eavours no fibringen concentrate		
Test for subgroup differences: $Chi^2 = 2.80$, df = 4 (P = 0.59), l ² = 0%							- avoirs nonnogen concentrate - r avoirs no nonnogen concentrate		
Footnotes									

(1) FC vs Platelets (2) FC vs CRYO

Figure 4.35 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Morbidity, multiple organ failure

	FC		No FC (or varyi	ng)		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	I	M-H, Random, 95% Cl
3.3.1 Trauma setting (RCTs)								
Nascimento 2016 (RCT, trauma)	2	21	2	24	6.0%	1.14 [0.18, 7.42]		
Akbari 2018 (RCT, trauma)	2	26	7	30	8.8%	0.33 [0.07, 1.45]		
Innerhofer 2017 (RCT, trauma)	25	50	29	44	32.0%	0.76 [0.53, 1.08]		
Subtotal (95% CI)		97		98	46.8%	0.74 [0.53, 1.03]		◆
Total events	29		38					
Heterogeneity: Tau ² = 0.00; Chi ² =	1.43, df =	2 (P = 0	0.49); l ² = 0%					
Test for overall effect: Z = 1.79 (P =	= 0.07)							
3.3.2 Trauma setting (Coh) Almskog 2020 (Coh, trauma) Nienaber 2011 (Coh, trauma)	1 3	108 18	1 11	108 18	3.0% 13.5%	1.00 [0.06, 15.78] 0.27 [0.09, 0.82]		
Wafaisade 2013 (Coh, trauma) Subtotal (95% CI)	180	294 420	144	294 420	36.7% 53.2%	1.25 [1.08, 1.45] 0.70 [0.21, 2.36]		*
Total events Heterogeneity: Tau ² = 0.75; Chi ² = Test for overall effect: $Z = 0.57$ (P :	184 7.45, df = = 0.57)	2 (P = 0	156 0.02); l² = 73%					
Total (95% CI)		517		518	100.0%	0.76 [0.46, 1.26]		-
Total events	213		194					
Heterogeneity: Tau ² = 0.17; Chi ² = 16.14, df = 5 (P = 0.006); l ² = 69% Test for overall effect: Z = 1.06 (P = 0.29) Test for subgroup differences: Chi ² = 0.01, df = 1 (P = 0.94), l ² = 0%								0.1 1 10 100 Favours fibrinogen concentrate
Table 4.57 Results for FC versus no FC: Patients with critical bleeding – Morbidity: critical complications

Study ID	Sample size	Patient S	Setting	Comparison	Outcome	Results					
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC n/N (%)	No FC n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b		
Trauma setting											
Fabes 2018 SR	N = 39 (1 RCT) Curry 2018	Adults ≥16 years with active bleeding	Trauma centre (UK)	FC vs saline	TE (arterial or venous)	3/20 (15)	2/19 (10.5)	NR	NR		
High quality	N = 94 (1 RCT) Innerhofer 2017	Adults with ISS >15 and clinical signs or risk of substantial haemorrhage	Trauma centre (Austria)	FC vs FFP		7/50 (14)	9/44 (20.5)	RR 0.63 (0.21, 1.87)	NR		
McQuilten 2018 SR <i>Moderate</i>	N = 45 (1 RCT) Nascimento 2016	Patients at risk for significant haemorrh age	Trauma (Canada)	FC vs placebo (normal saline)	Pulmonary embolus	2/21 (9.5)	1/24 (4.2)	RR 2.3 (0.2, 23.4)	NR		
quality					Symptomatic DVT	0	0	Not estimable	NR		
					DVT on leg doppler	2/15 (13.3)	3/14 (21.4)	RR 0.62 (-0.1, 3.2)	NR		
					Myocardial Infarction	0	0	Not estimable	NR		
					Stroke	0	0	Not estimable	NR		
					Acute kidney injury	3/21 (14.3)	2/24 (8.3)	RR 1.7 (-0.3, 9.3)	NR		
					MOF	2/21 (9.5)	2/24 (8.3)	RR 1.1 (-0.2, 7.4)	NR		
					Infection	5/21 (23.8)	8/24 (33.3)	RR 0.7 (-0.3, 1.8)	NR		
Stabler 2020 SR <i>Moderate</i>	N = 836 (1 RCT, 2 Coh)	Adult patients with major blunt trauma	Trauma (Germany, Sweden)	FC vs no FC	TE RCTs Lucena 2020	0/16 (0)	0/16 (O)	NR	<i>p</i> = NR		
quality					Observational Wafaisade 2013 Almskog 2020	20/294 (6.8) 5/108 (4.6)	10/294 (3.4) 3/108 (2.8)	NR NR	Favours no FC <i>p</i> = 0.06 <i>p</i> = 0.47		
	N = 366 (2 RCT, 1 Coh)	Adult patients with major blunt trauma	Trauma centre (Iran, Sweden)	FC vs no FC	MOF RCTs Akbari 2018 Innerhofer 2017 Observational Almskog 2020	2/26 (7.6) 25/50 (50) 1/108 (0.9)	7/30 (23.3) 29/44 (66) 1/108 (0.9)	NR NR	p = 0.106 p = 0.15 p = 1.00		

Study ID	Sample size	Patient S	Setting	Comparison	Outcome	Results				
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC n/N (%)	No FC n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b	
Aubron 2014 SR <i>Critically low</i> <i>quality</i>	N = 588 (1 Coh) Wafaisade 2013	Adult trauma patients with ISS >16 + at least 1 RBC + TASH score >9	Trauma (Germany)	FC vs No FC	MOF	180/294 (61.2)	144/294 (49)	NR	<i>Favours FC</i> <i>p</i> = 0.003	
	N = 36 (1 Coh) Nienaber 2011	Adult trauma patients	Trauma (Germany, Austria)	FC vs FFP		3/18 (16.7)	11/18 (61)	NR	<i>Favours FC</i> <i>p</i> = 0.015	
Mengoli 2017 SR <i>Low quality</i>	N = 36 (1 Coh) Nienaber 2011	Patients with severe trauma-related bleeding	Trauma (Germany, Austria)	FC vs FFP	Sepsis ^c	3/18 (16.7)	6/18 (33.3)	NR	No significant difference <i>p</i> = 0.443	
Inokuchi 2017 Coh	N = 224 (1 Coh)	Patients with pelvic fractures from blunt	Trauma (SC, Japan)	FC (after revision of MTP)	Transarterial embolisation	28/109 (26)	36/115 (31)	NR	No significant difference p = 0.764	
Serious risk of bias		trauma requiring activation of MTP		vs MTP (before revision)	Patients requiring external fixation	14/109 (13)	13/115 (11)	NR	No significant difference p = 0.838	
					Patients requiring internal fixation	43/109 (39)	42/115 (36)	NR	No significant difference p = 0.681	
					Patients requiring pelvic packing	2/109 (2)	3/115 (3)	NR	No significant difference p = 1.000	
Surgical setting										
Fabes 2018 SR High quality	N = 120 (1 RCT) Bilecen 2017	Adults >18 undergoing elective high-risk cardiac	Surgery (Netherlands)	FC vs placebo	TE (arterial or venous)	7/60 (11.7)	3/60 (5)	RR 2.33 (0.63, 8.60)	NR	
		surgery			Allergic AE up to 30 days	0/60	0/60	RD 0.0 (-0.03, 0.03)	NR	
٦	N = 61 (1 RCT) Rahe-Meyer 2013	Adults >18 with elective aortic valve replacement surgery	Surgery (Germany)	FC vs saline	TE (arterial or venous)	1/29 (3.4)	2/32 (3.1)	NR	NR	
					Allergic AE up to 10 days	0/29	0/32	RD 0.0 (-0.06, 0.06)	NR	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC n/N (%)	No FC n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b		
	N = 20 (1 RCT) Tanaka 2014	Patients undergoing elective	Surgery (US)	FC vs PLT	TE (arterial or venous)	0/10	1/10 (10)	RR 0.33 (0.02, 7.32)	NR		
		cardiopulmonary bypass procedures			Post-operative atrial fibrillation	0/6	1/12 (8%)	NR	NR		
					Renal failure °	0/6	2/12 (17%)	NR	NR		
					Major neurologic events ^c	0/6	2/12 (17%)	NR	NR		
Paediatric, surgi	cal setting										
Fabes 2018 SR High	N = 63 (1 RCT) Galas 2014	Patients <15 years undergoing CPB, intraoperative bleeding and hypofibrinogenemia	Paediatric surgery (Brazil)	FC vs CRYO	TE (arterial or venous)	2/30 (6.7)	5/33 (12.2)	RR 0.44 (0.09, 2.10)	NR		
Obstetrics and n	naternity setting										
Zaidi 2020 SR <i>Low</i>	N = 299 (2 RCTs) Collins 2017 Wikkelsø 2015	Women with major PPH	Obstetrics (Denmark, UK)	FC vs saline	Thrombosis up to 6 weeks	1/28 (3.6 0/12) 1/27 (3.7) 3 0/121	NR	NR		

AE, adverse events; CI, confidence interval; Coh, cohort; CPB, cardiopulmonary bypass; CRYO, cryoprecipitate; DVT, deep vein thrombosis; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ISS, injury severity score; MOF, multiorgan failure; NR, not reported; PLT, platelets; PPH, postpartum haemorrhage; RBC, red blood cell; RCT, randomised controlled trial; RD, risk difference; RR, relative risk; SC, single centre; SR, systematic review; TASH, trauma associated severe haemorrhage; TE, thromboembolic events; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

d. Calculated post-hoc using RevMan 5.4

Table 4.58 Results for FC versus no FC: Patients with critical bleeding – Morbidity: ARDS

Study ID	Sample size (no. Of trials)	Patient Suppopulation (1	Setting (Comparison	Outcome	Results														
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC n/N (%)	No FC n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b											
Trauma setting					1				notorogononj											
McOuilton 2019		Dationts at risk for	Trauma SC	EC vs placaba	ADDS	0/21 (0)	2/24 (9.2)	NA	ND											
MCQuitteri 2016	N = 45 (I R C I)	Patients at fisk for	Hauffia, SC	FC vs placebo	ARDS	0/21(0)	2/24 (0.3)	INA	INK											
SR	Nascimento 2016	significant haemorr	(Canada)	(normal saline)																
Moderate		hage																		
quality		_																		
Surgical setting	- no identified stud	lies reported on outco	ome of interest																	
Paediatrics, surgical setting – no identified studies reported on outcome of interest																				
Obstetrics and r	naternity setting - i	no identified studies i	reported on outcome	of interest					Obstetrics and maternity setting - no identified studies reported on outcome of interest											

ARDS, acute respiratory distress syndrome; CI, confidence interval; FC, fibrinogen concentrate; NA, not applicable; NR, not reported; RCT, randomised controlled trial; SC, single centre; SR, systematic review

a. (18)Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Transfusion volumes

Red blood cells

A summary of the evidence reported in the identified systematic reviews relating to RBC transfusion volumes in patients with critical bleeding is presented in Table 4.59.

One RCT and 4 cohort studies reported the effect of FC on RBC transfusion volume in trauma patients with critical bleeding (see Figure 4.36). Data from Wafaisade 2013 suggested a higher volume of RBC was required for patients who received FC (n=294) compared with those who did not (n=294), but the difference was not significant (SMD 0.12; 95% CI –0.04, 0.28; p = 0.14). The other 4 studies (one RCT, 3 cohort studies) reporting median [IQR] values suggested there was no significant difference in the volume of RBC transfused (comparing patients who received FC compared with those who did not). Reported median values ranged from 3 to 12.8 units (FC) and 3 to 12.5 units (no FC).

Two cohort studies reported the effect of FC on RBC transfusion volume in the surgical setting (see Figure 4.36). Data from Rahe-Meyer 2009a suggested that patients who received FC had a lower volume of RBC transfused compared with patients who did not receive FC (SMD –1.69, 95% CI –2.49, –0.88; p < 0.0001). The other study (Rahe-Meyer 2009b) reported that there were significantly fewer (p < 0.05) median units of RBC transfused to 24 hours in patients who received FC compared with those who did not.

One cohort study (Ahmed 2012) reported the effect of FC on RBC transfusion volume among women with major postpartum haemorrhage. The study reported a lower volume of RBC transfused among women who received FC compared with those who did not (SMD –0.29; 95% CI –0.98, 0.40; p = 0.41) but the difference was not significant.

There were no studies that reported on the outcome of RBC transfusion volume in the paediatric setting.

Figure 4.36 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: RBC transfusion volume, units

		FC		No FC	(or varyi	ng)	:	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.4.1 Trauma setting (RCTs)									
Nascimento 2016 (RCT, trauma) (1) Subtotal (95% CI)	0	0	21 0	0	0	24 0		Not estimable Not estimable	
Heterogeneity: Not applicable									
Test for overall effect: Not applicable									
3.4.2 Trauma setting (Coh)									
Inokuchi 2017 (Coh, trauma) (2)	0	0	115	0	0	109		Not estimable	
Nienaber 2011 (Coh, trauma) (3)	0	0	18	0	0	18		Not estimable	
Schochl 2011 (Coh, trauma) (4)	0	0	80	0	0	601		Not estimable	
Wafaisade 2013 (Coh, trauma) Subtotal (95% CI)	12.8	14.3	294 294	11.3	10	294 294	37.7% 37.7%	0.12 [-0.04, 0.28] 0.12 [-0.04, 0.28]	₽
Heterogeneity: Not applicable									
Test for overall effect: Z = 1.47 (P = 0.14	1)								
x	,								
3.4.4 Surgical setting (Coh)									
Rahe-Meyer 2009b (Coh, surgical) (5)	0	0	6	0	0	12		Not estimable	
Rahe-Meyer 2009a (Coh, surgical)	0.5	1.1	16	2.4	1.1	17	30.3%	-1.69 [-2.49, -0.88]	_
Subtotal (95% CI)			16			17	30.3%	-1.69 [-2.49, -0.88]	\bullet
Heterogeneity: Not applicable									
Test for overall effect: Z = 4.08 (P < 0.00	001)								
3.4.5 Obstetrics and maternity (Coh)									
Ahmed 2012 (Coh, obstetrics) (6)	5.9	4.293	20	7.21	4.602	14	32.1%	-0.29 [-0.98, 0.40]	
Subtotal (95% CI)			20			14	32.1%	-0.29 [-0.98, 0.40]	-
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.83 (P = 0.41	I)								
Total (95% CI)			330			325	100.0%	-0.56 [-1.54, 0.43]	
	1 44 0	(D . 0 0	004\. 12	000/		323	100.0 /6	-0.30 [-1.34, 0.43]	
Therefore the provide the provided the prov	+, di = 2 (7)	(٣ < 0.0	1001); l²	= 90%					-4 -2 0 2 4
Test for overall effect: $Z = 1.11$ (P = 0.27))	a (n	0.004	12 00 7	n/				Favours fibrinogen concentrate Favours no fibrinogen concentrate
rest for subgroup differences: ChP = 19	.34, ai =	2 (P < l	J.UUU1)	, ı~ = ŏ9./	70				

Footnotes

(1) No significant difference for RBC transfusion volume (median [IQR]) to 24 hours between FC (3 [2, 5]) and no FC (3 [2, 4]).

(2) No significant difference for RBC transfusion volume (median [IQR]) up to 7 days between FC (10 [6, 20]) and no FC (10 [4, 22]).

(3) No significant difference for RBC transfusion volume (median [IQR]) to 24 hours between FC (3 [0, 5]) and no FC (12.5 [8, 20]).

(4) No difference for RBC transfusion volume (median [IQR]) between FC (5.5 [0, 9.5]) and no FC (6 [4, 11]). Timepoint and p-values not reported.

(5) Significantly fewer (p<0.05) median units of RBCs transfused to 24 hours in patients who recieved FC (1.0) compared with those who did not (4.1). IQR values not reported.

(6) SD calculated from SEM.

Table 4.59 Results for FC versus no FC: Patients with critical bleeding – RBC transfusion volume

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis	population	(Location)			FC Mean ± SD	No FC Mean ± SD	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Trauma settin	ng									
McQuilten 2018 SR Moderate quality	N = 45 (1 RCT) Nascimento 2016	Patients at risk for significant haemorrhage	Trauma (SC, Canada)	FC vs placebo (normal saline)	RBC transfusion volume, units to 24 hours	Median (IQR) 3 (2, 5) (n = 21)	Median (IQR) 3 (2, 4) (n = 24)	Not estimable	No significant difference p = 0.41	
Mengoli 2017 SR <i>Low quality</i>	N = 588 (1 Coh) Wafaisade 2013	Patients with severe trauma- related bleeding	Trauma (MC, multicountry)	FC vs No FC	RBC transfusion volume, units '	12.8 ± 14.3 (n = 294)	11.3 ± 10.0 (n = 294)	NR	No significant difference p = 0.20	
Aubron 2014 SR <i>Critically low</i> <i>quality</i>	N = 717 (2 Coh) Schochl 2011 Nienaber 2011	Adult trauma patients	Trauma (MC, Germany and Austria)	FC vs FFP Schochl 2011	RBC transfusion volume, units ^c	Median (IQR) 5.5 (0, 9.5) (n = 80)	Median (IQR) a.	NR (1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	NR	
				FC vs FFP Nienaber 2011	RBC transfusion volume, units to 24 hours ^c	Median (IQR) 3 (0, 5) (n = 18)	Median (IQR) 12.5 (8, 20) (n = 18)	NR	<i>p</i> < 0.005	
Inokuchi 2017 Coh Serious risk of bias	N = 224 (1 Coh)	Patients with pelvic fractures from blunt trauma requiring activation of MTP	Trauma (SC, Japan)	FC (after revision of MTP) vs MTP (before revision)	RBC transfusion volume, units to 7 days	Median (IQR) 10 (6, 20) (n = 109)	Median (IQR) 10 (4, 22) (n = 115)	NR	No significant difference <i>p</i> = 0.958	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC Mean ± SD	No FC Mean ± SD	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
Surgical settir	ng										
Lunde 2014 SR <i>Critically low</i> <i>quality</i>	N = 18 (1 Coh) Rahe-Meyer 2009b	Patients undergoing cardiac surgery with bleeding requiring FC	Surgery (SC, Germany)	FC vs standard infusion (FFP + PLT)	RBC transfusion volume, units to 24 hours	b.	4.1 (n = 12)	NR	Favoured FC p < 0.05		
					RBC transfusion volume, mL to 24 hours	449.2 (n = 6)	1092.5 (n = 12)	NR	Favoured FC p < 0.05		
Warmuth 2012 SR <i>Low quality</i>	N = 33 (2 Coh)	Adult patients undergoing surgery with major haemorrhage	Surgery (Germany)	FC vs standard infusion (FFP + PLT)	RBC transfusion volume, units Rahe-Meyer 2009a Rahe-Meyer 2009b	(n = 16) 0.5 ± 1.1 1.0	(n = 17) 2.4 ± 1.1 4.1	NR	Favoured FC p < 0.05		
Paediatrics, su	urgical setting – no	identified studies re	ported on outc	ome of interest							
Obstetrics and	d maternity setting										
Lunde 2014 SR <i>Critically low</i>	N = 34 (1 Coh) Ahmed 2012	Women with PPH requiring FC	Obstetrics (Ireland)	FC vs CRYO	RBC transfusion volume, units ^c	(n = 20) 5.90 (SEM 0.96)	(n = 14) 7.21 (SEM 1.23)	NR	No significant difference p = 0.40		

CI, confidence interval; Coh; cohort; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; IQR, interquartile range; MC, multicentre; MTP, massive transfusion protocol; NR, not reported; PLT, platelets; PPH, postpartum haemorrhage; RBC, red blood cells; RCT, randomised controlled trial; SC, singe centre; SD, standard deviation; SEM, standard error of mean; SR, systematic review

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

d. Calculated post-hoc using RevMan 5.4

Other blood components

A summary of the evidence reported in the identified systematic reviews relating to transfusion volumes (other blood components) in patients with critical bleeding is presented in Table 4.60.

One RCT and 4 cohort studies reported on the effect of FC on the volume of FFP transfused in the trauma setting (see Figure 4.37).

Data from Wafaisade 2013 showed a statistically significant increase in the volume of FFP transfused among patients who received FC (n=294) compared with those who did not (n=294) (SMD 0.19, 95% CI 0.03, 0.35; p = 0.02).

Among the other 4 studies (one RCT, 3 cohort studies), 2 studies reporting median [IQR] values suggested there was no significant difference in the volume of FFP transfused between patients who received FC compared with those who did not (Inokuchi 2017, Nascimento 2016). One study found a decrease in the volume of FFP transfused among patients who received FC compared with those who did not (Nienaber 2011), and one study did not report comparative data for this outcome.

One RCT and 3 cohort studies reported on the effect of FC on the volume of PLT transfused in the trauma setting. Among the 3 studies that reported comparative data, 2 studies suggested there was no significant difference in the volume of PLT transfused between patients who received FC compared with those who did not (Nascimento 2016, Inokuchi 2017). One cohort study (Nienaber 2011) reported a significant reduction (p < 0.005) in platelet transfusion among patients who received FC compared with those who did not, but no further data was provided.

One RCT reported on the effect of FC on the volume of CRYO transfused in the trauma setting and found no significant difference between treatment groups (p = 0.18).

Among critically bleeding patients in the surgical setting, there was a significant reduction in the volume of FFP transfused among patients who received FC compared to those who did not (SMD -4.78, 95%CI -7.04, -2.51; p < 0.0001). Two cohort studies also found a statistically significant reduction in the volume of PLT and PCC transfused among patients who received FC compared to those who did not (p < 0.05) (see Figure 4.38).

Among women with major postpartum haemorrhage, no significant difference in the volume of FFP or PLT transfused between treatment groups was observed.

One systematic review (Zaidi 2020) reported the effect of FC on transfusion volume among women with major postpartum haemorrhage. The systematic review authors identified one RCT (Collins 2017) that they used to determine the total volume of blood transfused per patient at 7 days (inclusive of RBC, FFP, CRYO, FC, PLT, PC) between women who received TEG guided early administration of FC compared with those who did not. An adjusted rate ratio 0.72 (95% CI 0.30, 1.70) was reported (p = 0.45).

Figure 4.37 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Transfusion volume, other blood components, FFP (trauma)

		FC		No FC (or varyi	ng)	:	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% Cl
3.5.1 FFP, units									
Inokuchi 2017 (Coh, trauma) (1)	0	0	115	0	0	109		Not estimable	
Nascimento 2016 (RCT, trauma) (2)	0	0	21	0	0	24		Not estimable	
Nienaber 2011 (Coh, trauma) (3)	0	0	18	0	0	18		Not estimable	
Schochl 2011 (Coh, trauma) (4)	0	0	80	0	0	601		Not estimable	
Wafaisade 2013 (Coh, trauma) Subtotal (95% CI)	10.6	11.4	294 528	8.7	8.2	294 1046	100.0% 100.0%	0.19 [0.03, 0.35]	
Heterogeneity: Not applicable Test for overall effect: Z = 2.31 (P = 0.0	02)								
Total (95% CI)			528			1046	100.0%	0.19 [0.03, 0.35]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 2.31 (P = 0.0	02)								-0.5 -0.25 0 0.25 0.5
Test for subgroup differences: Not appl	licable								Favours indiniogen concentrate Favours no librinogen concentrate
Ecotrotos									

Ecothotes (1) No significant difference (median [IQR]) for FFP transfusion volume up to 7 days comparing FC (8 [6, 20]) with no FC (10 [6, 20]). (2) No significant difference (p=0.72) in median [IQR] FFP transfusion volume up to 24 hours among patients who received FC (2.73 [2.4, 3.6]) compared with no FC 1.75 [1.4, 2.0]). (3) FFP transfusion volume (units) up to 24 hours reported to be lower in patients who received FC (0) compared with no FC (10). IQR and p-value not reported. (4) No data reported for the outcome of transfusion volume, other blood products.

Figure 4.38 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Transfusion volume, other blood components, FFP (surgical)

FC				No FC (or varying)			5	Std. Mean Difference	e Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	CI IV, Random, 95% CI
3.6.3 FFP, units (Coh)									
Rahe-Meyer 2009a (Coh, surgical)	0.2	0.6	10	4.2	1.1	5	100.0%	-4.78 [-7.04, -2.51]	A]
Rahe-Meyer 2009b (Coh, surgical)	1	0	6	9.1	0	12		Not estimable	le
Subtotal (95% CI)			16			17	100.0%	-4.78 [-7.04, -2.51]	1]
Heterogeneity: Not applicable									
Test for overall effect: Z = 4.14 (P < 0	0.0001)								
			16			17	100.0%	4 79 [7 04 2 54]	11
10tal (95% CI)			10			17	100.0%	-4.78 [-7.04, -2.51]	
Heterogeneity: Not applicable									-10 -5 0 5 10
Test for overall effect: Z = 4.14 (P < 0 Test for subgroup differences: Not ap	0.0001) oplicable	•							Favours fibrinogen concentrate Favours no fibrinogen concentrate

Table 4.60 Results for FC versus no FC: Patients with critical bleeding – Transfusion volume, other blood components

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. Of trials) included in analysis	population	(Location)			FC Mean ± SD	No FC Mean ± SD	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Trauma setting										
McQuilten 2018 SR <i>Moderate</i>	N = 45 (1 RCT) Nascimento 2016	Patients at risk for significant haemorrhage	Trauma (SC, Canada)	FC vs placebo (normal saline)	Plasma transfusion volume, units to 24 hours Median (IQR)	(n = 21) 2.73 (2.4, 3.6)	(n = 24) 1.75 (1.4, 2.0)	Not estimable	No significant difference p = 0.72	
quality					PLT transfusion volume, units to 24 hours Median (IQR)	(n = 21) 2.81 (2.5, 3.6)	(n = 24) 2.32 (1.9, 2.7)	Not estimable	No significant difference p = 0.53	
					CRYO transfusion volume, units to 24 hours Median (IQR)	(n = 21) 4.0 (3.1, 4.6)	(n = 24) 3.5 (2.9, 4.0)	Not estimable	No significant difference p = 0.18	
Aubron 2014 SR	N = 717 (2 Coh)	Adult trauma patients	Trauma (MC, Germany and	FC vs FFP Schöchl 2011	FFP transfusion volume, units ^c	(n = 80) NA	(n = 601) 3	NR	NR	
Critically low quality	Schöchl 2011 Nienaber 2011		Austria)		PLT transfusion volume, units ^c	(n = 80) 1 or 2	(n = 601) NR	NR	NR	
					PCC transfusion volume, international units ^c	(n = 80) 1200	(n = 601) NR	NR	NR	
					FC dose, grams ^c	(n = 80) 6	(n = 601) NA	NR	NR	
				FC vs FFP Nienaber 2011	FFP transfusion volume, units to 6 hours ^c	(n = 18) O	(n = 18) 6	NR	NA	
					FFP transfusion volume, units to 24 hours ^c	(n = 18) O	(n = 18) 10	NR	NA	
					PLT transfusion volume, units to 24 hours ^c	(n = 18) O	(n = 18) 2	NR	p < 0.005	
					FC dose, grams to 6 hours c	(n = 18) 4	(n = 18) O	NR	NA	
					FC dose, grams to 24 hours °	(n = 18) 4	(n = 18) O	NR	NA	
					PCC transfusion volume, international units to 6 hours ^c	(n = 18) 1200	(n = 18) O	NR	NA	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC Mean ± SD	No FC Mean ± SD	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
					PCC transfusion volume, international units to 24 hours ^c	(n = 18) 1200	(n = 18) O	NR	NA	
Mengoli 2017 SR <i>Low</i>	N = 588 (1 Coh) Wafaisade 2013	Adult trauma patients	Trauma (MC, multicountry)	FC vs No FC	FFP transfusion volume, units ^c	(n = 294) 10.6 ± 11.4	(n = 294) 8.7 ± 8.2	NR	No significant difference p = 0.07	
Inokuchi 2017 Coh <i>Serious risk of</i>	N = 224 (1 Coh)	Patients with pelvic fractures from blunt	Trauma (SC, Japan)	FC (after revision of MTP) vs MTP (before	FFP transfusion volume, units to 7 days	Median (IQR) (n = 109) 8 (6, 20)	(n = 115) 10 (6,20)	NR	No significant difference <i>p</i> = 0.685	
bias		trauma requiring activation of MTP		revision)	PLT transfusion volume, units to 7 days	Median (IQR) (n = 109) 20 (20, 20)	(n = 115) 20 (20, 37.5)	NR	No significant difference p = 0.251	
Surgical setting										
Lunde 2014 SR	N = 18 (1 Coh) Rahe-Meyer 2009b	Patients undergoing	Surgery (SC, Germany)	FC vs standard infusion (FFP +	FFP transfusion volume, units to 24 hours ^c	(n = 6) 1.0	(n = 12) 9.1	NR	Favoured FC p < 0.05	
Critically low		cardiac surgery with bleeding requiring FC		PLT)	PLT transfusion volume, units to 24 hours ^c	(n = 6) 0.5	(n = 12) 3.2	NR	Favoured FC p < 0.05	
Warmuth 2012 SR <i>Low</i>	N = 33 (2 Coh)	Adult patients undergoing surgery with major	Surgery (Germany)	FC vs standard infusion (FFP + PLT)	FFP transfusion volume, units to 24 hours Rahe-Meyer 2009a	(n = 16) 0.2 ± 0.6	(n = 17) 4.2 ± 1.1	NR	<i>Favoured FC</i> <i>p</i> < 0.05	
		haemorrhage			PCC transfusion volume, international units to 24 hours Rahe-Meyer 2009a Rahe-Meyer 2009b	(n = 16) 0.0 0.5	(n = 17) 1.6 ± 0.9 3.2	NR	<i>Favoured FC</i> <i>p</i> < 0.05	
Paediatrics, sur	gical setting			·			·			
Obstetrics and	maternity setting									
Zaidi 2020 SR <i>Low</i>	N = 299 (2 RCTs) Collins 2017	Women with major PPH	Obstetrics (UK)	FC vs saline	Total volume of blood transfused (inclusive of RBC, FFP, CRYO, FC, PLT, PCC)	(n=28) NR	(n=27) NR	NR	No significant difference p = 0.45	
Lunde 2014 SR	N = 34 (1 Coh) Ahmed 2012	Patients with PPH requiring FC	Obstetrics (Ireland)	FC vs CRYO	FFP transfusion volume, units ^c	(n = 20) 3.15 (SEM 0.65)	(n =14) 4.07 (SEM 0.74)	NR	No significant difference <i>p</i> = 0.36	

Study ID	Sample size I (no. Of trials)	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. Of trials) included in analysis	population	(Location)			FC Mean ± SD	No FC Mean ± SD	Risk estimate (95% CI)	Statistical significance	
	-								<i>p</i> -value	
									Heterogeneity ^b	
Critically low					PLT transfusion volume,	(n = 20)	(n =14)	NR	No significant difference	
					pools ^c	1.00 (SEM 0.30)	1.00 (SEM 0.36)		<i>p</i> = 0.99	
					FC dose, grams post	(n = 20)	(n =14)	NR	No significant difference	
					treatment ^c	3.34 SEM (0.22)	3.05 (SEM 0.19)		<i>p</i> = 0.35	

Cl, confidence interval; Coh, cohort; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; IQR, interquartile range; MC, multicentre; MTP, massive transfusion protocol; NR, not reported; PLT, platelets; PPH, postpartum haemorrhage; RCT, randomised controlled trial; SC, single centre; SD, standard deviation; SEM, standard error of mean; SR, systematic review; UK, United Kingdom

a. (18)Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

d. Calculated post-hoc using RevMan 5.4

Length of stay

Hospital

A summary of the evidence reported in the identified systematic reviews relating to hospital LOS in patients with critical bleeding is presented in Table 4.61.

Four RCTs and 3 cohort studies reported the effect of FC on hospital LOS in the trauma setting (see Figure 4.39). Data were available for 2 studies (reported as mean [SD]), that showed FC has no significant impact on the duration of hospital stay comparing patients who received FC with those who did not (RR –1.30; 95% CI –6.76, 4.16; p = 0.64; random effects; I² = 69%), noting the heterogeneity was substantial. The remaining studies reported data as median (IQR) that also suggested there is no significant difference inhospital LOS between patients who received FC and those who did not.

Among critically bleeding patients with postpartum haemorrhage, no significant difference was reported for hospital LOS between treatment groups.

Intensive care unit (ICU)

A summary of the evidence reported in the identified systematic reviews relating to ICU LOS in patients with critical bleeding is presented in Table 4.62.

Two RCTs and 4 cohort studies reported the effect of FC on ICU LOS (days) in the trauma setting (see Figure 4.40). Complete data were not available, but 5 of the 6 studies suggested that there is no significant difference in the duration of ICU stay for patients who received FC compared to those who did not. One RCT (Lucena 2020) suggested that the length of ICU stay among patients who received FC was lower (p = 0.021) than the length of ICU stay among patients who did not.

There was one cohort study in the surgical setting (Rahe-Meyer 2009b) that reported on ICU LOS (hours) which suggested FC is associated with a reduction in the length of ICU stay among patients who received FC compared with those who did not (MD – 3.27, 95% CI –4.82, –1.71; p < 0.0001; [hours converted to days]); however, the sample size is small and survivorship bias may have influenced the results.

Among women with major postpartum haemorrhage, one cohort study (Ahmed 2012) reported that there was no significant difference in the length of ICU stay between patients who received FC compared with those who did not (p = 0.95).

Figure 4.39 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Length of stay, hospital (days)

		FC		No FC	(or varyi	ng)		Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.7.1 Trauma setting (RCTs)									
Curry 2018 (RCT, trauma) (1)	0	0	24	0	0	24		Not estimable	
Innerhofer 2017 (RCT, trauma) (2)	0	0	50	0	0	44		Not estimable	
Lucena 2020 (RCT, trauma) (3)	0	0	16	0	0	16		Not estimable	
Akbari 2018 (RCT, trauma)	11	6.1	30	14.8	7.6	30	55.3%	-3.80 [-7.29, -0.31]	
Subtotal (95% CI)			96			90	55.3%	-3.80 [-7.29, -0.31]	\bullet
Heterogeneity: Not applicable									
Test for overall effect: Z = 2.14 (P = 0	0.03)								
0.7.0.7.									
3.7.2 Trauma setting (Con)	-								
Nienaber 2011 (Con, trauma) (4)	0	0	18	0	0	18		Not estimable	
Schochi 2011 (Con, trauma) (5)	0	0	80	0	0	601	44 70/	Not estimable	
Subtotal (95% CI)	34.0	33.3	294	32.8	28.4	294	44.7%	1.80 [-3.20, 6.80]	
Heterogeneity: Net applicable			002			0.0		1100 [0120, 0100]	
Test for overall effect: Z = 0.71 (P = 0	1 4 9 1								
	5.40)								
3.7.3 Obstetrics and maternity									
Collins 2017 (RCT, obstetrics) (6)	0	0	28	0	0	27		Not estimable	
Ahmed 2012 (Coh, obstetrics) (7)	6.55	3.6224	20	5.21	1.2347	14		Not estimable	
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not applicable									
Test for overall effect: Not applicable									
			400			1002	100.0%	1 20 [6 76 4 46]	
10tal (95% CI)			400			1003	100.0%	-1.30 [-0.70, 4.10]	
Heterogeneity: 1au ² = 10.84; Chi ² = 3	3.24, df =	= 1 (P =)	0.07); l²	= 69%					-20 -10 0 10 20
Test for overall effect: $Z = 0.47$ (P = 0	0.64)	4 (5	0.07)						Favours fibrinogen concentrate Favours no fibrinogen concentrate
Lest for subgroup differences: Chi ² =	: 3.24, df	'= 1 (P =	: 0.07), I	² = 69.19	%				

Footnotes (1) No data reported for hospital LOS

(2) No difference (p=0.61) in median [IQR] reported for hospital LOS for patients who received FC (28 [18, 28]) compared with those who did not (27 [16, 28]).
(3) No difference (p=NR) in median [IQR] reported for hospital LOS for patients who received FC (28 [18, 28]) compared to those who did not (27 [16, 28]).
(4) No difference (p=0.48) in median [IQR] reported for hospital LOS for patients who received FC (28 [18, 28]) compared with those who did not (27 [16, 28]).

(c) Hospital LOS (median [IQR]) was significantly lower (p=0.05) for patients who received FC (23 [14,5, 40.5)) compared with those who did not (3 [2, 4]).
 (c) No difference (p=0.19) in median [IQR]) reported for hospital LOS for patients who received FC (3 [2, 5]) compared with those who did not (3 [2, 4]).
 (c) SD calculated from SEM.

Figure 4.40 Forest plot of comparison: FC vs no FC (or varying administration of), outcome: Length of stay, ICU (days)

		FC		No FC	(or varyi	ng)		Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.8.1 Trauma setting (RCTs)									
Innerhofer 2017 (RCT, trauma) (1)	0	0	50	0	0	44		Not estimable	
Lucena 2020 (RCT, trauma) (2)	0	0	16	0	0	16		Not estimable	
Subtotal (95% CI)			66			60		Not estimable	
Heterogeneity: Not applicable									
Test for overall effect: Not applicable									
3.8.2 Trauma setting (Coh)									
Almskog 2020 (Coh, trauma) (3)	0	0	108	0	0	108		Not estimable	
Nienaber 2011 (Coh, trauma) (4)	0	0	18	0	0	18		Not estimable	
Schochl 2011 (Coh, trauma) (5)	0	0	80	0	0	601		Not estimable	
Wafaisade 2013 (Coh, trauma)	17.2	17.6	294	17.3	17.9	294	42.4%	-0.10 [-2.97, 2.77]	
Subtotal (95% CI)			500			1021	42.4%	-0.10 [-2.97, 2.77]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.07 (P = 0.95	5)								
3.8.3 Surgical setting (Coh)									
Rahe-Mever 2009b (Coh. surgical) (6)	1.542	0.7875	6	4.808	2.5083	12	57.6%	-3.27 [-4.82, -1.71]	— — —
Subtotal (95% CI)			6			12	57.6%	-3.27 [-4.82, -1.71]	◆
Heterogeneity: Not applicable									
Test for overall effect: Z = 4.12 (P < 0.00	001)								
Total (95% CI)			572			1093	100.0%	-1.92 [-4.99, 1.14]	
Heterogeneity: Tau ² = 3.63; Chi ² = 3.62,	df = 1 (F	e = 0.06);	$l^2 = 72^4$	%				-	
Test for overall effect: Z = 1.23 (P = 0.22	2)								-10 -5 U 5 10 Favours fibringgen concentrate
Test for subgroup differences: Chi ² = 3.6	62, df = 1	(P = 0.06)	6), l ² = 1	72.4%					avours nonnogen concentrate - 1 avours no libilitogen concentrate

Test for subgroup dimeterizes: Chi^{*} = 3.52, di = 1 (F = 0.06), F = 72.4%
<u>Foolnotes</u>:
(1) No difference (p=0.65) in median [IQR] ICU LOS for patients who received FC (9 [4, 22]) compared with those who did not (10 [4.8, 23.3]).
(2) Median [IQR] ICU LOS was significant lower (p=0.021) for patients who received FC (8 [5.75, 10]) compared to those who did not (11 [8.5, 16]).
(3) No difference (p=0.628) in median [IQR] ICU LOS for patients who received FC (19 [9, 33]) compared to those who did not (5 [1, 16]).
(4) No difference (p=0.95) in median [IQR] ICU LOS for patients who received FC (14.5 [8.5, 21]) compared to those who did not (14 [6, 23]).
(5) No difference (p=0.95) in median [IQR] ICU LOS for patients who received FC (14.5 [8.5, 21]) compared to those who did not (14 [6, 23]).

(6) Data converted from hours to days

Table 4.61 Results for FC versus no FC: Patients with critical bleeding – Hospital LOS

Study ID Sample size		Patient	Setting	Comparison	Outcome	Results				
Study	(no. Of trials)	population	(Location)			FC	No FC	Risk estimate	Statistical significance	
design	analysis					Mean ± SD	Mean ± SD	(95% CI)	<i>p</i> -value	
									Heterogeneity ^b	
Trauma setting	g									
Stabler 2020	N = 822	Adult patients with	Trauma (MC,	FC vs no FC	Hospital LOS, days			NR		
SR	(4 RCTs, 1 Coh)	major blunt	multi-country)		Curry 2018	NR 11 (C1)	NR 14.0 (7.0)		p = NR	
Moderate		trauma			Wafaisade 2013 (Coh)	34.6 (33.3)	32.8 (28.4)		p = 0.045	
						Median (IQR)	Median (IQR)			
					Lucena 2020	12 (10-22)	18.5 (17-21)		p = 0.61	
	N 717 (2 0 - h)	Detients with	Tanuara (NAC	50 m 550	Innerhofer 2017	28 (18-28)	7 (16-28)	ND	p = 0.96	
Mengoli 2017	N = /1/ (2 CON)	Patients with	multi-country)	FCVSFFP	In-patient, days *	Median (IQR)	Median (IQR)	NR	p = 0.005	
		related bleeding	mail country)		Schoem zon	23 (14.5-40.5)	32 (20-49)		p = 0.003	
2011						(n = 18)	(10)			
					Nienaber 2011	26 (19-50)	(n = 18) 38 (21-48)		No significant difference p = 0.481	
	N = 588 (1 Coh)	Patients with	Trauma (MC,	FC vs No FC	In-patient, days °	(n = 294)	(n = 294)	NR	No significant difference	
	Wafaisade 2013	severe trauma-	multi-country)			34.6 ± 33.3	32.8 ± 28.4		<i>p</i> = 0.96	
		related bleeding								
Surgical settin	g – no identified st	udies reported on o	utcome of interes	st						
Paediatrics, su	ırgical setting - no	identified studies re	eported on outco	me of interest						
Obstetrics and	I maternity setting	I								
Zaidi 2020	N = 55 (1 RCT)	Women with major	Obstetrics (UK)	FC vs saline	Hospital LOS, days	Median (IQR)	Median (IQR)	NR	No significant difference	
SR	Collins 2017	PPH				3 (2, 5)	3 (2, 4)		<i>p</i> = 0.13	
Low										
Lunde 2014	N = 34 (1 Coh)	Patients with PPH	Obstetrics (UK)	FC vs CRYO	In-patient, days ^c	(n = 20)	(n =14)	NR	No significant difference	
SR	Ahmed 2012	requiring FC				6.55 (SEM 0.81)	5.21 (SEM 0.33)		<i>p</i> = 0.19	
Critically low										

Cl, confidence interval; Coh, cohort; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; IQR, interquartile range; LOS, length of stay; MC, multicentre; NR, not reported; PPH, postpartum haemorrhage; RCT, randomised controlled trial; SEM, standard error of the mean; SD, standard deviation; SR, systematic review; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study

Table 4.62 Results of FC versus no FC: Patients with critical bleeding – ICU LOS

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis		(Location)		F(M		No FC Mean ± SD	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting				1	1					
Stabler 2020 SR <i>Moderate quality</i>	N = 930 (2 RCTs, 2 Coh)	Adult patients with major blunt trauma	Trauma (MC, multi- country)	FC vs no FC	ICU LOS, days Wafaisade 2013 (Coh) Lucena 2020 Innerhofer 2017 Almskog 2020 (Coh)	Mean ±SD 17.2 ± 17.6 Median (IQR) 8 (5.75-10) 9 (4-22) 7 (1-20)	Mean ± SD 17.3 ± 17.9 Median (IQR) 11 (8.5-16) 10 (4.8-23.3) 5 (1-16)	NR	p = 0.68 p = 0.021 p = 0.65 p = 0.97	
Mengoli 2017 SR <i>Low quality</i>	N = 717 (2 Coh) Schöchl 2011 Nienaber 2011	Patients with severe trauma-related bleeding	Trauma (MC, multi- country)	FC vs FFP	ICU LOS, days ^c Schöchl 2011 Nienaber 2011	Median (IQR) 14.5 (8.5-21) (n = 80) 19 (9-33) (n = 18)	Median (IQR) 14 (6-23) (n = 601) 16 (13-25) (n = 18)	NR	No significant difference p = 0.95 p = 0.628	
	N = 588 (1 Coh) Wafaisade 2013	Patients with severe trauma-related bleeding	Trauma (MC, multi- country)	FC vs No FC No FC	ICU LOS, days °	(n = 294) 17.2 ± 17.6	(n = 294) 17.3 ± 17.9	NR	No significant difference p = 0.68	
Surgical setting										
Lunde 2014 SR <i>Critically low quality</i>	N = 18 (1 Coh) Rahe-Meyer 2009b	Patients undergoing cardiac surgery with bleeding requiring FC	Surgery (SC, Germany)	FC vs standard infusion (FFP + PLT)	ICU LOS, hours ^c	(n = 6) 37 ± 18.9	(n =12) 115.4 ± 60.2	NR	p < 0.05	
Paediatrics, surgica	l setting – no identi	fied studies reported or	n outcome of in	terest						
Obstetrics and mate	ernity setting									
Lunde 2014 SR <i>Critically low quality</i>	N = 34 (1 Coh) Ahmed 2012	Patients with PPH requiring FC	Obstetrics	FC vs CRYO	High dependency unit, hours ^c	(n =20) 33.6 (SEM 5.44)	(n = 14) 34.1 (SEM 4.32)	NR	No significant difference p = 0.95	
Zaidi 2020 SR <i>Low quality</i>	N = 55 (1 RCT) Collins 2017	Women with major PPH	Obstetrics (MC, UK)	FC vs saline	ICU LOS, days	Median (IQR) 16 (12-25)	Median (IQR) 20.5 (10.5-28.5)	Difference 0.90	NR	

Cl, confidence interval; Coh, cohort; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ICU, intensive care unit; IQR, interquartile range; LOS, length of stay; MC, multicentre; NR, not reported; PLT, platelets; PPH, postpartum haemorrhage; RCT, randomised controlled trial; SC, single centre; SD, standard deviation; SEM, standard error of mean; SR, systematic review; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data extracted from primary study.

4.7.3.5 Prothrombin complex concentrate

Mortality

A summary of the evidence reported in the identified systematic reviews relating to mortality in patients with critical bleeding is presented in Table 4.63.

A meta-analysis of data from the 4 retrospective cohort studies identified in the systematic review by van den Brink 2020 (see Figure 4.41) revealed a significant reduction in mortality among patients who received PCC (72/364, 19.8%) compared with those who did not (159/557, 28.5%), representing an odds ratio (OR) of 0.64 (95%CI 0.46, 0.88; p = 0.007; random effects; $l^2 = 0$ %).

Figure 4.41 Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: Mortality (trauma setting)

	PCC	;	no PCC (or var	ying)		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
4.1.1 trauma setting							
Jehan 2018 (Coh, trauma) (1)	10	40	26	80	14.2%	0.69 [0.29, 1.63]	
Joseph 2014 (Coh, trauma) (2)	15	63	53	189	23.8%	0.80 [0.41, 1.55]	
Joseph 2016 (Coh, trauma) (3)	6	27	15	54	8.8%	0.74 [0.25, 2.20]	
Zeeshan 2019 (Coh, trauma) (4) Subtotal (95% CI)	41	234 364	65	234 557	53.2% 1 00.0%	0.55 [0.35, 0.86] 0.64 [0.46, 0.88]	-
Total events	72		159				
Heterogeneity: Tau ² = 0.00; Chi ² =	0.98, df =	3 (P =	0.81); l² = 0%				
Test for overall effect: Z = 2.72 (P =	= 0.007)						
Total (95% CI)		364		557	100.0%	0.64 [0.46, 0.88]	•
Total events	72		159				
Heterogeneity: Tau ² = 0.00; Chi ² =	0.98, df =	3 (P =	0.81); l² = 0%				
Test for overall effect: Z = 2.72 (P = 0.007)							0.01 0.1 1 10 100 Eavours PCC Eavours no PCC
Test for subgroup differences: Not	applicable	e					
Footnotes							
(4) Other is a second and second in a second							

(1) Study carried out in any setting

(2) Study carried out in any setting

(3) Study carried out in any setting

(4) Study carried out in any setting

Table 4.63 Results for PCC versus no PCC: Patients with critical bleeding – Mortality, latest timepoint

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			PCC n/N (%)	No PCC n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting									
van den Brink 2020	N = 921 (4 Coh)	Patients ≥18 years with	Trauma (US)	4 factor PCC + FFP vs FFP	Mortality, overall	72/364 (19.8)	159/557 (28.5)	OR 0.64 (0.46, 0.88)	<i>Favours PCC</i> p = 0.007
SR <i>Low quality</i>		active bleeding		3 factor PCC + FFP vs FFP	Jehan 2018 Zeeshan 2019 Joseph 2014 Joseph 2016	10/40 (25) 41/234 (17.5) 15/63 (23.8) 6/27 (22.2)	26/80 (32.5) 65/234 (27.8) 53/189 (28.0) 15/54 (27.8)	0.69 (0.29, 1.63) 0.55 (0.35, 0.86) 0.80 (0.41, 1.55) 0.74 (0.25, 2.20)	No heterogeneity 1² = 0% (<i>p</i> = 0.81)

Cl, confidence interval; Coh, cohort; FFP, fresh frozen plasma; OR, odds ratio; PCC, prothrombin complex concentrate; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Morbidity

Thromboembolic events

A summary of the evidence reported in the identified systematic reviews relating to morbidity (critical complications) in patients with critical bleeding is presented in Table 4.64.

A meta-analysis of data from the 4 retrospective cohort studies identified in the systematic review by van den Brink 2020 (see Figure 4.42) showed no significant difference in thromboembolic events between treatment groups (OR 0.90, 95%Cl 0.49, 1.67; p = 0.74; random effects; $l^2 = 0$ %).

Figure 4.42 Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: Morbidity, thromboembolic events (trauma setting)

	PCC no l		no PCC (or var	no PCC (or varying)		Odds Ratio		Odds Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl		M-H, Rand	om, 95% Cl	
Jehan 2018 (Coh, trauma) (1)	1	40	2	80	6.5%	1.00 [0.09, 11.37]				
Joseph 2014 (Coh, trauma) (2)	2	63	3	189	11.6%	2.03 [0.33, 12.45]				
Joseph 2016 (Coh, trauma) (3)	4	27	5	54	19.3%	1.70 [0.42, 6.95]				
Zeeshan 2019 (Coh, trauma) (4)	11	234	17	234	62.6%	0.63 [0.29, 1.38]			-	
Total (95% CI)		364		557	100.0%	0.90 [0.49, 1.67]				
Total events	18		27							
Heterogeneity: Tau ² = 0.00; Chi ² = 2	2.38, df =	3 (P =	0.50); l² = 0%							100
Test for overall effect: Z = 0.33 (P =	0.74)						0.01	Favours PCC	Favours no PCC	100
Footnotes										
(1) Study carried out in any setting										
(2) Study carried out in any setting										
(3) Study carried out in any setting										
(4) Study carried out in any setting										

Acute respiratory distress or other adverse outcomes

No comparative evidence for PCC versus no PCC was reported in the systematic review by van den Brink 2020 for the outcome of ARDS or other adverse outcomes.

The 4 retrospective cohort studies may have measured and reported these outcomes, but because retrospective studies were considered to be inappropriate for inclusion (see Section 3.1.4), we did not retrieve for inspection.

Table 4.64 Results for PCC versus no PCC: Patients with critical bleeding - Morbidity: critical complications

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			PCC n/N (%)	No PCC n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting									
van den Brink	N = 921 (4 Coh)	Patients ≥ 18 years	Trauma (US)	4 factor PCC +	TEs	18/364 (4.9)	27/557 (4.8)	OR 0.90 (0.49, 1.67)	No significant difference
2020		of age with active		FFP vs FFP				1.00 (0.09, 11.37)	<i>p</i> = 0.74
SR		bleeding			Jehan 2018	1/40 (2.5)	2/80 (2.5)	2.03 (0.33, 12.45)	No heterogeneity
Low quality					Zeeshan 2019	2/63 (3.2)	3/189 (1.6)		$l^2 = 0\% (D < 0.50)$
				3 factor PCC +	Joseph 2014	4/27 (14.8)	5/54 (9.3)	1.70 (0.42, 6.95)	
				FFP vs FFP	Joseph 2016	11/234 (4.7)	17/234 (7.3)	0.63 (0.29, 1.38)	

Cl, confidence interval; Coh, cohort study; FFP, fresh frozen plasma; PCC, prothrombin complex concentrate; OR, odds ratio; SR, systematic review; TEs, thromboembolic events; US, United States a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Transfusion volumes

Red blood cells

A summary of the evidence reported in the identified systematic reviews relating to RBC transfusion volumes in patients with critical bleeding is presented in Table 4.65.

A meta-analysis of data from the 4 retrospective cohort studies identified in the systematic review by van den Brink 2020 (see Figure 4.43) showed a significant reduction in the volume of RBC transfused among patients that received PCC compared with those who did not (standardised MD –0.65; 95%Cl –0.98, –0.32; p = 0.0001; random effects; $l^2 = 77\%$), noting the heterogeneity was substantial.

Figure 4.43 Forest plot of comparison: PCC vs no PCC (or varying administration of), outcome: RBC transfusion volume, Units (trauma setting)

	PCC			no PCC (or varying)			5	Std. Mean Difference	Std. Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
Jehan 2018 (Coh, trauma) (1)	7	3	40	9	5	80	23.1%	-0.45 [-0.83, -0.06]	_	
Joseph 2014 (Coh, trauma) (2)	6.6	4.1	63	10	8.3	189	26.8%	-0.45 [-0.74, -0.17]		
Joseph 2016 (Coh, trauma) (3)	3.2	1.9	27	5.4	4.1	54	19.9%	-0.62 [-1.09, -0.15]		
Zeeshan 2019 (Coh, trauma) (4)	6	4	234	10	4	234	30.2%	-1.00 [-1.19, -0.81]		
Total (95% CI)			364			557	100.0%	-0.65 [-0.98, -0.32]	◆	
Heterogeneity: Tau ² = 0.08; Chi ² = 13.17, df = 3 (P = 0.004); l ² = 77%					7%					
Test for overall effect: Z = 3.87 (P =	= 0.0001)							Favours PCC Favours no PCC	
Footnotes										
(1) Study carried out in any setting										
(2) Study carried out in any setting										
(3) Study carried out in any setting										
(4) Study carried out in any setting										

Other blood components

No comparative evidence for PCC versus no PCC was reported in the SR by van den Brink 2020 for the outcome of transfusion volume (other blood components).

The 4 retrospective cohort studies may have measured and reported these outcomes, but because retrospective studies were considered to be inappropriate for inclusion (see Section 3.1.4), we did not retrieve for inspection.

Table 4.65 Results for PCC versus no PCC: Patients with critical bleeding – RBC transfusion volume

Study ID Sample size		Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)			PCC Mean ± SD	No PCC Mean ± SD	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting										
van den Brink	N = 921 (4 Coh)	Patients ≥ 18 years	Trauma (US)	4 factor PCC + FFP	RBC utilisation,	N = 364	N = 557	MD -2.99 (-4.06, -1.91)	Favours PCC	
2020		of age with active		vs FFP	units				<i>p</i> <0.00001	
SR		bleeding			Jehan 2018	7±3 (n=40)	9±5 (n=80)	-2.00 (-2.44, -0.56)	Significant heterogeneity	
Low quality					Zeeshan 2019	6±4 (n=234)	10±4 (n=234)	-4.00 (-4.72, -3.28)	I ² = 68% (p<0.0001)	
				3 factor PCC + FFP vs FFP	Joseph 2014 Joseph 2016	6.6±4.1 (n=63) 3.2±1.9 (n=27)	10±8.3 (n=189) 5.4±4.1 (n=54)	-3.40 (-4.96, -1.84) -2.20 (-3.51, -0.89)		

Cl, confidence interval; Coh, cohort study; FFP, fresh frozen plasma; MD, mean difference; PCC, prothrombin complex concentrate; RBC, red blood cell; SD, standard deviation; SR, systematic review; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

Length of stay

No comparative evidence for PCC versus no PCC was reported in the systematic review by van den Brink 2020 regarding length of stay (hospital or ICU).

The 4 retrospective cohort studies may have measured and reported this outcome, but because retrospective studies were considered to be inappropriate for inclusion (see Section 3.1.4), we did not retrieve for inspection.

4.8 Antifibrinolytics (Question 7)

Question 7 - (interventional)

In patients with critical bleeding, what is the effect of antifibrinolytics on blood loss, RBC transfusion and patient outcomes?

4.8.1 Methods

Question 7 examined the effect of antifibrinolytics (TXA, aprotinin, or EACA) on patient outcomes compared to no antifibrinolytics in patients with critical bleeding (i.e. major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion) as outlined in Figure 4.44.

This question focused on intravenous delivery of TXA³¹ with the timing of delivery being at onset of bleeding (i.e. therapeutic use). Patients admitted in any setting were eligible for inclusion including trauma, obstetrics, and perioperative (e.g. cardiothoracic, liver transplant), with a subgroup analysis of evidence in patients who received a massive transfusion to be conducted where possible. Studies where bleeding status was not assessed at the time of enrolment were excluded (such as those that randomised patients prior to elective cardiac surgery). No age limits were applied, however studies in neonates (newborns up to 28 days) and studies in individuals with hereditary bleeding disorders were not eligible for inclusion.

Figure 4.44 PICO criteria: Question 7 – antifibrinolytics



ARDS, acute respiratory distress syndrome; EACA, epsilon-aminocaproic acid; RBC, red blood cell; TE, thromboembolic event; TXA, tranexamic acid

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. c. Newborns up to 28 days following birth.

³¹ EACA is not available or licensed for use in Australia and aprotinin, although on the Australian Register of Therapeutic Goods, is not being supplied or marketed by the Australian sponsor.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2000, noting primary studies published prior to 2000 and identified within a systematic review were also eligible for inclusion. All RCTs were eligible for inclusion regardless of sample size, however, nonrandomised or observational cohort studies were required to enrol at least 500 participants.

Assuming all relevant primary studies had been identified in the included systematic reviews³²; the screening of primary studies for this question was not conducted. This is because the latest literature search date of the most comprehensive identified systematic reviews was 2018 (El-Menyar 2018, Gayet-Ageron 2018, Shakur 2018).

The literature search was updated in August 2019³³ and again in September 2021³⁴ to identify any new studies meeting the eligibility criteria. In these updated searches the focus was the identification of systematic reviews (of RCTs or cohort studies). Based on the latest literature search date of the most comprehensive identified systematic reviews, the screening for additional primary studies was not conducted.

It was noted that there is an ongoing international multicentre, randomised, doubleblind, placebo-controlled trial of prehospital treatment with TXA for severely injured patients at risk of acute traumatic coagulopathy. The study aims to determine the effects of early administration of TXA on survival and recovery of severely injured patients treated within advanced trauma systems.

4.8.2 Summary of evidence

4.8.2.1 Systematic reviews

Thirteen systematic reviews (61, 143-154) were included that assessed the effects of antifibrinolytics compared to no antifibrinolytics in patients with critical bleeding. The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.66. A matrix illustrating the overlap of studies included in each review is provided in Table 4.67.

Eight systematic reviews were identified for inclusion (155-162), but were later excluded. One review (Bennett 2014) was in adults undergoing emergency or urgent surgery for upper gastrointestinal bleeding, but the studies were confounded by the administration of oral TXA (in combination with IV TXA) and were not reflective of current standard of care. Two other reviews (Burke 2021, Lee 2021), also in adults with gastrointestinal bleeding, included the same studies identified by Bennett 2014, plus one additional RCT (HALT-IT) that was relevant to this review. As HALT-IT was the only RCT to meet the inclusion criteria for this review, the primary study was retrieved and included (see Section 4.8.2.2).

³² Nine SRs identified (El-Menyar 2018, Gayet-Ageron 2018, Shakur 2018, Cannon 2017, Huebner 2017, Nishida 2017, Ausset 2015, Ker 2015, Bennett 2014)

³³ One additional SR (Chornenki 2019) and two observational studies identified (Marsden 2019, Myers 2019)

³⁴ Four additional SRs identified (Al-Jeabory 2021, Almuwallad 2021, Ageron 2020, Della Corte 2020)

Five reviews (Baskaran 2018, Gausden 2017, Wang 2017, Zhang 2017, Perel 2013) looked for studies in patients with orthopaedic trauma, but the patients within the included studies were not critically bleeding at study entry and therefore did not meet the inclusion criteria for this review (noting that the intervention was being delivered to prevent, rather than treat, perioperative bleeding and was not given in response to haemorrhage).

A list of studies that *met* the PICO criteria for this question but are excluded or awaiting classification is provided in **Appendix B** (technical report, volume 2).

Review ID Review quality	Study design <i>Risk of bias</i>	Population	Intervention	Comparator	Outcomes
Any setting					
Ageron 2020 (163) <i>Moderate</i>	MA of individual patient-level data from RCTs involving over 1000 patients (2 studies)	Patients with acute severe haemorrhage (trauma, PPH) N=28 333	Antifibrinolytics (TXA, aprotinin, EACA)	Placebo or SoC	Mortality* (due to bleeding) Morbidity* (TE) *By baseline risk of death
Chornenki 2019 (146) <i>High</i>	SR / MA of RCTs (22 studies)ª	Patients with haemorrhage, TBI or non-specific trauma injury N= 49 538	ТХА	Placebo or no TXA	Mortality Morbidity (TE) ^b
Gayet-Ageron 2018 (149) <i>High</i>	MA of individual patient-level data from RCTs involving over 1000 patients (2 studies) ^c	Patients with acute severe haemorrhage (traumatic, PPH, emergency) N=40 138	Antifibrinolytics (TXA, aprotinin, EACA)	Placebo or SoC	Mortality (due to bleeding) Morbidity (TE)
Trauma settin	g				
Al-Jeabory 2021 (143) <i>High</i>	SR / MA of RCTs and cohort studies (17 studies)	Adult patients following acute traumatic injury	ТХА	Placebo or no TXA	Mortality Morbidity (TE, MOF) Transfusion volume
Almuwallad 2021 (144) <i>High</i>	SR / MA of RCTs and cohort studies (17 studies)	Adult patients following acute traumatic injury	TXA (prehospital administration)	Placebo or no TXA	Mortality Morbidity (TE)
El-Menyar 2018 (148) <i>Moderate</i>	SR / MA of RCTs (2 studies)	Traumatic injury patients presenting to the ED requiring blood transfusions N=769	TXA (prehospital administration)	Placebo or no TXA	Mortality Morbidity (TE)
Nishida 2017 (152) <i>Moderate</i>	SR of RCTs and observational studies (8 studies) ^d	Adult patients with trauma- induced coagulopathy N=23 117	TXA	Placebo or no TXA	Morbidity (TE)

Table 4.66	Characteristics and quality of systematic review evidence: Antifibrinolytics versus
	no antifibrinolytics

Review ID Review quality	Study design <i>Risk of bias</i>	Population	Intervention	Comparator	Outcomes
Huebner 2017 (151) <i>Critically low</i>	Narrative review of RCTs and cohort studies (8 studies)	Haemorrhaging paediatric and adult trauma patients	TXA (early and prehospital administration)	Placebo	Mortality
Cannon 2017 (61) <i>Moderate</i>	SR of RCTs and cohort studies (4 studies)	Adult patients with severe traumatic haemorrhage	TXA	Placebo or no TXA	Mortality Transfusion rates Morbidity (TE)
Ker 2015 (154) <i>High</i>	SR of RCTs (3 studies) ^{d. e}	People of any age following acute traumatic injury N=20 528	Antifibrinolytics (aprotinin, TXA, EACA or AMBA)	Placebo or no TXA	Mortality Morbidity (TE) Transfusion volume Transfusion rate
Ausset 2015 (153) <i>Critically low</i>	Narrative review	Trauma management in the prehospital setting	TXA	Placebo or no TXA	Mortality Morbidity (TE)
Obstetrics and	l maternity				
Della-Corte 2020 (145) <i>Moderate</i>	SR / MA of RCTs	Women with PPH after vaginal birth N=20 212 (2 studies)	Antifibrinolytics (TXA, aprotinin, EACA)	Placebo or SoC	Mortality (all-cause, due to bleeding) Morbidity (MOF, hysterectomy, TE, shock) Transfusion volume
Shakur 2018 (150) <i>High</i>	SR / MA of RCTs (3 studies) ^e	Women with PPH after birth (vaginal or caesarean section) following a pregnancy of at least 24 weeks' gestation N=20 212	Antifibrinolytics (TXA, aprotinin, EACA)	Placebo or SoC	Mortality Morbidity (MOF, hysterectomy, TE, shock) Transfusion volume

AMBA, aminomethylbenzoic acid; EACA, epsilon-aminocaproic acid; ED, emergency department; MA, meta-analysis; RCT, randomised controlled trial; SoC, standard of care; SR, systematic review; TBI, traumatic brain injury; TE, thromboembolic events; TXA, tranexamic acid

a. 20 studies did not meet eligibility criteria for this review as they were in adults with non-surgical indications (including TBI) where TXA was used for prevention of bleeding as part of a planned protocol.

b. TE includes vascular occlusive events, myocardial infarct, deep vein thrombosis, pulmonary embolism, stroke, etc.

c. The authors conducted logistic regression models assessing overall treatment effect and effect of treatment delay. All models were controlled for systolic blood pressure (5 mm Hg interval) and age (10-yr intervals), which are strong risk factors for death due to bleeding.

d. One RCT (Yutthakasemsunt 2013) was in patients with traumatic brain injury and is not relevant to this review.

e. One RCT (McMichan 1982) examined the effect of aprotinin in patients with a combination of hypovolaemic shock and major fractures of the lower limb and or pelvis and did not provided any data relevant to this review.

f. One RCT (Sahaf 2014) did not contribute data for outcomes relevant to this review (only estimated blood loss) and is not further discussed.

Three reviews assessed patients in both the trauma and non-trauma setting (Ageron 2020, Chornenki 2019, Gayet-Ageron 2018). Two of these reviews (Ageron 2020, Gayet-Ageron 2018) were secondary analyses of individual patient data from 2 large RCTs conducted in trauma (CRASH-2) and obstetrics (WOMAN) patients. One review (Chornenki 2019) included studies that administered TXA for the prevention of bleeding as part of a planned surgical protocol or as planned medical management (including brain injury) therefore the reported analysis was not relevant for this review.

Eight reviews focused on patients in the trauma setting (Al-Jeabory 2021, Almuwallad 2021, El-Menyar 2018, Nishida 2017, Huebner 2017, Cannon 2017, Ker 2015, Ausset 2015). The most recent and comprehensive review was that by Al-Jeabory 2021, which included and reported data from 3 RCTs and 14 prospective or retrospective cohort studies. Two reviews (Almuwallad 2021, El-Menyar 2018) focused on the impact of prehospital TXA administered to bleeding trauma patients during air and/or ground medical transport, or upon arrival to designated trauma centres. One review (Heubner 2017) included paediatric patients (one cohort study found) and the remaining 4 reviews (Nishida 2017, Cannon 2017, Ker 2015, Ausset 2015) reported additional studies that were not covered in the review by Al-Jeabory 2021.

There were 2 systematic reviews (Della-Corte 2020, Shakur 2018) that focused on women in the obstetric setting, both of which reported data from 2 RCTs conducted in women after birth (vaginal or caesarean section) with postpartum haemorrhage.

		Trauma: RCTs				Trauma: Observational cohort studies											Obstetric: RCTs								
	Study ID	Guyette 2020 (STAAMP)	Kakaei 2017	Shakur 2010 (CRASH-2) a	Rivas 2021	Adair 2020	Cole 2020	El-Menyar 2020	Myers 2019	Ng 2019	Neeki 2018	Howard 2017	Neeki 2017	Wafaisade 2016	Cole 2015	Harvin 2015	Eckert 2014	Lipsky 2014	Haren 2014	Morrison 2013 (MATTERs II)	Swendsen 2013	Valle 2014	Morrison 2012 (MATTERs I)	Arulkumaran 2017 (WOMAN)	Ducloy-Bouthers 2011
	Ageron 2020			~		_									-							-		~	
	Chornenki 2019			~																				~	
	Gayet- Ageron 2018			~																				~	
	Al-Jeabory 2021	~	~	~	~	~	~	~	~	~	~	~	~	~				~			~	~	~		
0	Almuwallad 2021	~						~			~			~											
iew II	El-Menyar 2018												~	~											
Rev	Nishida 2017			~										•	•	•			✓		•		•		
-	Huebner 2017			~										~	~	~	~				~	~	~		
	Cannon 2017			~											~					~			~		
	Ker 2015			~																					
	Ausset 2015			~											~					~		~	~		
	Della-Corte 2020																							~	~
	Shakur 2018																							~	~

Table 4.67Overlap table of studies identified by included systematic reviews:Antifibrinolytics

4.8.2.2 Randomised controlled trials

There were 6 RCTs identified through the included systematic reviews that met the inclusion criteria and were considered relevant to this review. The main characteristics and quality of these studies and the relevant outcomes assessed are summarised in Table 4.68.

CRASH-3 (a landmark study similar to CRASH-2) was not included as it examines the effects of tranexamic acid on death, disability, vascular occlusive events and other morbidities in patients with acute traumatic brain injury (TBI)³⁵.

Study ID	Study design	Population	Intervention	Comparator	Outcomes						
Trauma settin	Trauma setting										
Guyette 2020 (STAAMP) (164)	RCT, multicentre (US)	Civilian trauma patients	TXA 1 g TXA loading dose prehospital	No TXA	Mortality						
Kakaei 2017 (165)	RCT, SC (Iran)	Civilian trauma patients *with potentially life- threatening injuries or evidence of critical illness, which could include respiratory and cardiac arrest.	TXA 1 g TXA loading dose over 10 minutes followed by infusion of 1g over 8 hours	No TXA	Mortality						
Shakur 2010 (CRASH-2)ª (166)	RCT	Civilian trauma patients with wide range of injury severity scores *68% with blunt mechanism of injury, 18% with Glasgow Coma Score of ≤ 8.	TXA 1 g TXA loading dose over 10 minutes followed by infusion of 1g over 8 hours	No TXA	Mortality Morbidity (VE) Transfusion volume						
Medical emerg	gency		-								
Roberts 2020 (HALT-IT) (167, 168)	RCT	Adults with significant gastrointestinal bleeding *defined as a risk of bleeding to death ^b	TXA 1 g TXA loading dose over 10 minutes followed by infusion of 3g over 24 hours	No TXA	Mortality Morbidity (VE)						
Obstetric and	maternity										
Arulkumaran 2017 (WOMAN 2017) (169)	RCT	Women with clinically diagnosed PPH	TXA 1 g TXA as soon as possible ^c	Placebo ^d	Mortality Morbidity (vascular events, MOF)						

Table 4.68 Characteristics and quality of RCT evidence: Antifibrinolytics versus no antifibrinolytics

³⁵ The definition of critical bleeding for this review did not include haemorrhage of a smaller volume in a critical area or organ (e.g. intracranial, intraspinal or intraocular)

Study ID	Study design	Population	Intervention	Comparator	Outcomes
		(estimated blood loss after vaginal birth > 500 mL, or > 1000 mL after caesarean section or estimated blood loss enough to compromise the haemodynamic status of the woman)			
Ducloy- Bouthors 2011 (170)	RCT	Women with clinically diagnosed PPH after vaginal delivery of a baby or caesarean section.	TXA 4g TXA in 1 hour (loading dose) then 1g TXA per hour over 6 hours	No TXA	Mortality

PPH, primary postpartum haemorrhage; RCT, randomised controlled trial; TE, thromboembolic events; TXA, tranexamic acid; VE, vascular events

a. The previous guidelines referred to the CRASH-2 study (published after the literature search date), and guidance regarding the use of TXA was provided (not a Recommendation or a PP). The CRASH-2 study population does not fully align with the population of interest for the Guidelines.

b. including patients with hypotension, tachycardia, or signs of shock, or those likely to need transfusion or urgent endoscopy or surgery.

c. a second dose could be given if bleeding continued after 30 minutes or if bleeding stopped and restarted within 24 hours after first dose.

d. other interventions allowed included: oxytocin, ergometrine, misoprostol, prostaglandin, uterine massage, bladder catheter, manual removal of retained placenta (if necessary), intrauterine tamponade.

Trauma

Three RCTs (Guyette 2020, Kakaei 2017, CRASH-2) examined the effect of TXA in civilian trauma patients with critical bleeding. Participants were typically administered a loading dose of 1 g TXA as soon possible, followed by a maintenance dose of 1 g TXA over 8 hours. Patients included in the largest study (CRASH-2, ~20 000 participants) were classified as being at risk of significant bleeding, in addition to being diagnosed with major haemorrhage. Around 50% of enrolled patients did not receive a blood product. Participants had a wide range of injury severities, with most enrolled from over 40 low-income countries. The severity of diagnosis and life-threatening nature of haemorrhage for these patients was not specified and there was no systematic adverse event reporting, making it difficult to interpret results relating to thrombotic risk, and blood loss.

Medical emergency

One RCT (HALT-IT 2020) examined the effect of TXA in patients with acute upper or lower gastrointestinal bleeding (approximately 45% of patients had suspected variceal bleeding due to liver disease). The trial included 12 009 participants from 15 countries, who were randomised to receive either 1g TXA (IV infusion loading dose) followed by 3 g TXA maintenance dose (infused over 24 hours) or matching placebo (0.9% sodium chloride). Around 12% of patients did not have suspected active bleeding at enrolment and around 30% of patients did not receive a blood product. The primary outcome was death due to bleeding within 5 days of randomisation, and diagnosis of thromboembolic events was made using strict definitions and diagnostic criteria.

Obstetrics and maternity

Two RCTs (Ducloy-Bouthors 2011, WOMAN 2017) assessed the safety and effectiveness of TXA given to women with primary postpartum haemorrhage (PPH). Participants were typically administered a loading dose of 1 g TXA as soon possible after randomisation, and

if bleeding continued after 30 minutes, or stopped and restarted within 24 hours after first dose, a second dose could be given.

The largest study (WOMAN 2017, ~20 000 participants) enrolled women aged 16 years or older with clinically diagnosed PPH (estimated blood loss after vaginal birth more than 500 mL, or more than 1000 mL after caesarean section or estimated blood loss enough to compromise the haemodynamic status of the woman). Approximately 50% of participants had an estimated volume of blood loss less than 1000 mL and 41% had no clinical signs of haemodynamic instability. Around 54% of women received a blood product. There was no systematic adverse event reporting, making it difficult to interpret results relating to thrombotic risk and blood loss.

4.8.2.3 Observational and cohort studies

There were 16 cohort studies identified through the included systematic reviews that met the inclusion criteria and were considered relevant to this review. The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.70.

No additional retrospective cohort studies were identified in the systematic review and handsearching process.

The included cohort studies examined the effect of TXA in patients with critical bleeding after trauma (mixed combat and civilian trauma, including one paediatric trauma). All had concerns of bias relating to confounding (related to the co-administration of other products) and patient selection bias. There was also concerns for reporting bias with a lack of detail regarding injury severity, and protocols for adverse event reporting.

Study ID	Study design	Population	Intervention	Comparator	Outcomes	
Trauma settin	g					
Rivas 2021	Retrospective cohort	Civilian trauma patients	ТХА	No TXA	Mortality	
El-Menyar 2020	Retrospective cohort	Civilian trauma patients	ТХА	No TXA	Mortality Morbidity (VE)	
Myers 2019	Retrospective cohort	Civilian trauma patients	ТХА	No TXA	Mortality Morbidity (VE) RBC transfusion volume	
Neeki 2018	Prospective cohort, MC	Civilian trauma patients	TXA	No TXA *retrospective control group	Mortality	
Howard 2017	Retrospective cohort	Combat trauma	ТХА	No TXA	Mortality	
Neeki 2017	Prospective cohort, MC	Civilian trauma patients, with blunt or penetrating trauma* *resulting in signs and symptoms of haemorrhagic shock	TXA, prehospital	No TXA *retrospective control group	Mortality Morbidity (VE)	

Table 4.69Characteristics and quality of cohort evidence: Antifibrinolytics versus no
antifibrinolytics

Study ID	Study design	Population	Intervention	Comparator	Outcomes
Wafaisade 2016	Retrospective cohort	Trauma registry (German) Patients with potentially life-threatening injuries or evidence of critical illness, which could include respiratory and cardiac arrest	TXA, prehospital	No TXA *matched control	Mortality Morbidity (VE)
Cole 2014	Prospective cohort	Civilian trauma patients, (UK) ISS > 15 SBP < 90 mm Hg, poor response to fluids, suspected active haemorrhage)	TXA in trauma protocol	No TXA in trauma protocol	Mortality Morbidity (VE) RBC transfusion volume
Eckert 2014	Retrospective cohort	Paediatric trauma (Afghanistan) with predominantly penetrating injury (mean age 11 years)	ТХА	No TXA	Mortality
Haren 2014	Retrospective cohort	Civilian trauma patients with hypercoagulable state defined as Greenfield's risk assessment profile ≥ 10	ТХА	No TXA	Morbidity (VE)
Harvin 2014	Retrospective cohort	Civilian trauma patients with hyperfibrinolysis determined by rapid thromboelastography	ТХА	No TXA	Mortality Morbidity (VE)
Lipsky 2014	Retrospective cohort	Civilian trauma patients	TXA	No TXA	Mortality
Valla 2014	Prospective cohort	Civilian trauma patients	ТХА	No TXA	Mortality
Morrison 2013 (MATTERS II)	Retrospective cohort	Combat trauma patients receiving ≥ 1 Unit RBC	TXA, administered within 48hrs of injury	No TXA	Mortality RBC transfusion volume
Swendson 2013	Retrospective cohort	Civilian trauma patients within 3 hours of injury with an SBP < 90 mm Hg, activation of MTP at ED or taken directly to operating theatre matched to historical controls	ТХА	No TXA	Mortality Morbidity (VE)
Morrison 2012 (MATTERS)	Retrospective cohort	Combat trauma patients receiving ≥ 1 unit RBC 30% injured by gunshot wound, 70% injured by explosion, 29% with Glasgow Coma Score of ≤ 8	TXA, administered within 48 hrs of injury	No TXA	Mortality Morbidity (VE) RBC transfusion volume

Abbreviations: hrs, hours; TXA, tranexamic acid; RBC, red blood cells, VE, vascular events

4.8.3 Results

4.8.3.1 Mortality

A summary of the evidence relating to mortality (all-cause) in patients with critical bleeding treated with TXA is presented in Table 4.70.

It is noted that, due to substantial heterogeneity, data from systematic reviews that considered or stratified patients according to baseline risk of death, or mortality due to bleeding were considered not informative for this review.

A meta-analysis of data from studies included in this review (see Figure 4.45), the RCT evidence showed a slight decrease in the risk of mortality (latest timepoint) among trauma patients who received TXA (1503/10 537, 14.26%) compared with those who did not 1660/10 550, 15.73%) (RR 0.91; 95% CI 0.85, 0.97; p = 0.003; random effect, $I^2 = 0\%$) (GRADE: Low).

Among the cohort studies conducted in critically bleeding trauma patients, the risk of mortality was not different between groups (19.4% vs 17.26%, RR 0.97; 95%Cl 0.75, 1.25; p = 0.80, $l^2 = 90\%$) *(GRADE: Very low).* Noting there was substantial heterogeneity with a wide variety of injury severity and bleeding risk in the included studies, with the results likely to differ after adjustments for confounders across all studies (e.g. patients who received TXA had higher incidence of shock, blood loss or transfusion requirements).

In a sensitivity analysis, the risk estimate for mortality moved towards favouring TXA (RR 0.83, 95% CI 0.64, 1.06; p = 0.13, $I^2 = 87\%$) when the 3 studies that reported adjustment for confounders (Harvin 2014, Eckert 2014, Cole 2015) were removed from the analysis.

In a subgroup analysis examining the effect of TXA among civilian trauma patients (see Figure 4.46), a total of 12 649 patients received TXA compared with 13 168 patients who did not; with the combined RCT and cohort evidence suggesting no difference between groups (15.2% vs 17.1%, RR 0.90; 95% CI 0.73, 1.11; p = 0.32, random effect, $l^2 = 87\%$). In a sensitivity analysis, the risk estimate for mortality suggested a slight decrease in the risk of mortality favouring TXA (RR 0.78, 95% CI 0.65, 0.93; p = 0.006, $l^2 = 76\%$) when the 2 studies that reported adjustment for confounders (Harvin 2014, Cole 2015) were removed from the analysis.

In the medical emergency setting (serious GI bleeding), the RCT evidence (see Figure 4.45) suggested the mortality rate among patients who received TXA (564/5956, 9.5%) was comparable to the mortality rate among patients who did not receive TXA (548/5981, 9.2%). This corresponded to a RR of 1.03 (95%CI 0.92, 1.16; p = 0.56; random effect, I² = not applicable) (*GRADE: Low*).

In the obstetric setting, the RCT evidence (see Figure 4.45) suggested the mortality rate among women who received TXA (227/10 111, 2.2%) was comparable to the mortality rate among women who did not receive TXA (255/10 051, 2.5%). This corresponded to a RR of 0.89 (95% CI 0.74, 1.06; p = 0.18; random effect, I² = not applicable) (*GRADE: Low*).

Figure 4.45 Forest plot of comparison: TXA vs no TXA, outcome: Mortality, latest timepoint

	Tranexami	c acid	Cont	rol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% Cl
.1.1 Trauma (RCTs)							
Kakaei 2017 (Civilian)	3	30	4	30	0.2%	0.75 [0.18, 3.07]	
Guyette 2020 (STAAMP) (Civilian)	37	447	43	453	2.3%	0.87 [0.57, 1.33]	
CRASH-2 2010 (Civilian) (1)	1463	10060	1613	10067	97.4%	0.91 [0.85, 0.97]	
Subtotal (95% CI)		10537		10550	100.0%	0.91 [0.85, 0.97]	•
otal events	1503		1660				
Heterogeneity: Tau ² = 0.00; Chi ² = 0.10, df = 2 (Fest for overall effect: Z = 2.99 (P = 0.003)	(P = 0.95); l ² =	: 0%					
.1.2 Trauma (Coh)							
larvin 2014 (adult trauma) (2)	39	98	157	924		Not estimable	
Cole 2015 (Civilian, ISS 15) (3)	30	160	36	225		Not estimable	
ckert 2014 (paediatric trauma) (4)	10	66	56	700		Not estimable	
ipsky 2014 (Coh, trauma)	6	28	0	12	0.8%	5.83 [0.35, 95.93]	
leeki 2017 (prehospital)	8	128	13	125	5.1%	0.60 [0.26, 1.40]	
Swendsen 2013 (adult trauma)	9	52	17	74	6.0%	0.75 [0.36, 1.56]	
Jeeki 2018 (Civilian)	13	362	30	362	6.9%	0.43 [0.23, 0.82]	
/alle 2014 (adult trauma)	25	109	14	105	7.2%	1.72 [0.95, 3.12]	
I-Menyar 2020 (Civilian)	25	102	30	102	8.7%	0.83 [0.53, 1.31]	
Vafaisade 2016 (prehospital, civilian)	38	258	42	258	9.3%	0.90 [0.60, 1.35]	
Iorrison 2013 (MATTERS II) (5)	57	406	179	758	10.6%	0.59 [0.45, 0.78]	
ivas 2021 (Civilian)	106	654	91	254	10.9%	0.45 [0.36, 0.57]	
loward 2017 (Coh, combat)	82	849	271	2924	11.0%	1.04 [0.82, 1.32]	
Iorrison 2012 (MATTERS) (6)	148	293	218	603	11.6%	1.40 [1.20, 1.63]	
Nyers 2019 (Civilian) Subtotal (95% CI)	136	189 3430	161	189 5766	11.9% 100.0%	0.84 [0.76, 0.94] 0.83 [0.64, 1.06]	→
otal events	653		1066				
Heterogeneity: Tau ² = 0.14; Chi ² = 86.78, df = 1	1 (P < 0.0000	1); l ² = 87	7%				
est for overall effect: $Z = 1.50$ (P = 0.13)							
.1.3 Medical emergency (RCT)							<u> </u>
Roberts 2020 (HALT-IT) (GI bleeding, IV only)	564	5956	548	5981 5981	100.0%	1.03 [0.92, 1.16]	—
	564		E19				
lateregeneity: Net applicable	504		040				
Test for overall effect: $Z = 0.58$ (P = 0.56)							
.1.4 Obstetrics and maternity (RCTs)							
uclov-Bouthers 2011	0	77	0	74		Not estimable	
VOMAN 2017 (PPH)	227	10034	255	9977	100.0%	0.89 [0.74, 1.06]	
ubtotal (95% CI)		10111		10051	100.0%	0.89 [0.74, 1.06]	
otal events	227		255				
leterogeneity: Not applicable							
est for overall effect: Z = 1.35 (P = 0.18)							
						_	
							Favours TXA Favours placebo or no

Test for subgroup differences: Chi² = 5.18, df = 3 (P = 0.16), I² = 42.0%

Footnotes (1) within 4 weeks of injury (2) in-hospital; non significant effect after adjustment for confounders (OR 0.74, 95% CI 0.380, 1.403; p=0.801). (3) not adjusted for confounders (4) Effect favouring TXA observed after adjusting for confounders (OR 0.27; 95% CI 0.85, 0.89; p=0.03) (5) within 4 between 5 inter-

(5) within 48 hours of injury(6) within 48 hours of injury
	Tranexami	c acid	Cont	rol	Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
1.2.1 Civilian							
CRASH-2 2010 (Civilian) (1)	1463	10060	1613	10067	8.6%	0.91 [0.85, 0.97]	-
Myers 2019 (Civilian)	136	189	161	189	8.5%	0.84 [0.76, 0.94]	-
Rivas 2021 (Civilian)	106	654	91	254	7.6%	0.45 [0.36, 0.57]	-
Harvin 2014 (adult trauma) (2)	39	98	157	924	7.2%	2.34 [1.77, 3.11]	-
Wafaisade 2016 (prehospital, civilian)	38	258	42	258	6.1%	0.90 [0.60, 1.35]	
Guyette 2020 (STAAMP) (Civilian)	37	447	43	453	6.0%	0.87 [0.57, 1.33]	
Cole 2015 (Civilian, ISS 15) (3)	30	160	36	225	5.8%	1.17 [0.75, 1.82]	- -
El-Menyar 2020 (Civilian)	25	102	30	102	5.6%	0.83 [0.53, 1.31]	
Valle 2014 (adult trauma)	25	109	14	105	4.5%	1.72 [0.95, 3.12]	
Neeki 2018 (Civilian)	13	362	30	362	4.2%	0.43 [0.23, 0.82]	
Swendsen 2013 (adult trauma)	9	52	17	74	3.6%	0.75 [0.36, 1.56]	
Neeki 2017 (prehospital)	8	128	13	125	3.0%	0.60 [0.26, 1.40]	
Kakaei 2017 (Civilian)	3	30	4	30	1.4%	0.75 [0.18, 3.07]	
Subtotal (95% CI)		12649		13168	72.2%	0.90 [0.73, 1.11]	•
Total events	1932		2251				
Heterogeneity: Tau ² = 0.09; Chi ² = 89.3	6, df = 12 (P ·	< 0.00001); l ² = 87	%			
Test for overall effect: Z = 1.00 (P = 0.3	2)						
1.2.2 Combat							
Morrison 2012 (MATTERS) (4)	148	293	218	603	8.2%	1.40 [1.20, 1.63]	-
Howard 2017 (Coh, combat)	82	849	271	2924	7.6%	1.04 [0.82, 1.32]	+
Morrison 2013 (MATTERS II) (5)	57	406	179	758	7.3%	0.59 [0.45, 0.78]	
Eckert 2014 (paediatric trauma) (6)	10	66	56	700	4.3%	1.89 [1.02, 3.53]	
Lipsky 2014 (Coh, trauma)	6	28	0	12	0.4%	5.83 [0.35, 95.93]	
Subtotal (95% CI)		1642		4997	27.8%	1.13 [0.73, 1.74]	—
Total events	303		724				
Heterogeneity: $Tau^2 = 0.17$; $Chi^2 = 34.3$	0, df = 4 (P <	0.00001)	; l² = 88%	þ			
Test for overall effect: $Z = 0.56$ (P = 0.5	8)						
Total (95% CI)		14291		18165	100.0%	0.95 [0.79, 1.14]	•
Total events	2235		2975				
Heterogeneity: Tau ² = 0.10; Chi ² = 136.	46, df = 17 (P	< 0.0000)1); l ² = 8	8%			
Test for overall effect: Z = 0.54 (P = 0.5	9)		,-				0.01 0.1 1 10 100
Test for subgroup differences: Chi ² = 0.	, 88, df = 1 (P :	= 0.35), l ²	= 0%				Favours (experimental) Favours (control)
Footnotes	, , ,	,,					
(1) within 4 weeks of injury							
(2) in-hospital; non significant effect after	er adjustment	for confo	unders (C	DR 0.74,	95% CI 0	.380, 1.403; p=0.801).	
(3) not adjusted for confounders	,		(-	,			

Figure 4.46 Forest plot of comparison: TXA vs no TXA, outcome: Mortality, latest timepoint (trauma only)

(4) within 48 hours of injury

(5) within 48 hours of injury

(6) Combat zone (Afghanistan). Effect favouring TXA observed after adjusting for confounders (OR 0.27; 95% CI 0.85, 0.89; p=0.03)

Table 4.70 Results for TXA versus no TXA: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Any setting				- -	- -			- -	
Ageron 2020 SR <i>Moderate</i>	N = 28 333 (2 RCTs) CRASH-2 WOMAN	CRASH-2 are trauma patients, the WOMAN trials are women with a	Trauma, Obstetrics (MC, over 40 countries)	TXA vs placebo	Mortality, all- cause	434/14270 (3) NR NR	597/14063 (4.3) NR NR	RR 0.72 (0.63, 0.81) NR NR	No significant difference <i>p</i> = 0.98
quality	N = 28 333 (2 RCTs) CRASH-2 WOMAN	postpartum haemorrhage	Trauma, Obstetrics		Mortality, occlusive event	27/14270 (0.00) NR NR	40/14063 (0.00) NR NR	NR NR NR	No significant difference <i>p</i> = 0.058
Chornenki 2019	N = 44 077 (10 RCTs)	Patients requiring treatment or	Trauma, Obstetrics and Medical (MC,	TXA vs placebo or no TXA	Mortality, all- cause	2087/22014 (9.5)	2269/22063 (10.3)		p = NR
SR High quality	Chewdhary 1986 Tsementzis 1990 Roos 2000 Hiilman 2002 Shakur 2010 Yutthakasemsunt 2013 Spring 2014	prevention of postpartum haemorrhage, ICH, subarachnoid haemorrhage, TBI or non-specific trauma iniury	over 40 countries)			5/65 (7.7) 22/50 (44.0) 76/229 (33.2) 27/254 (10.6) 1463/10060 (14.5) 12/120 (10.0) 3/16 (18.8)	8/64 (12.5) 14/50 (28.0) 75/233 (32.2) 32/251 (12.7) 1613/10067 (16.0) 17/118 (14.4) 2/8 (25.0)	RR 0.62 (0.27, 1.78) RR 1.57 (0.91, 2.71) RR 1.03 (0.79, 1.34) RR 0.83 (0.52, 1.35) RR 0.91 (0.85, 0.97) RR 0.69 (0.35, 1.39) RR 0.75 (0.16, 3.62)	
	Arulkumaran 2017 Sprigg 2018 Fakharian 2018					227/9985 (2.3) 250/1161 (21.5) 2/74 (2.7)	256/10033 (2.6) 249/1164 (21.4) 3/75 (4)	RR 0.89 (0.75, 1.06) RR 1.01 (0.86, 1.18) RR 0.68 (0.12, 3.93)	
Gayet-Ageron 2018 SR High quality	N = 40 138 (2 RCTs) CRASH-2 2010 WOMAN 2017	Patients with acute severe bleeding	Trauma, Obstetrics (MC, over 40 countries)	TXA vs placebo	Mortality, all- cause	1690/20094 (8.4) 1463/10060 (14.5) 227/10034 (2.3)	1868/20044 (9.3) 1613/10067 (16.0) 255/9977 (2.6)	RR 0.90 (0.85, 0.96) ^c RR 0.91 (0.85, 0.97) RR 0.89 (0.74, 1.06)	Favours TXA $p = 0.001^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.79)$
	N = 40 138 (2 RCTs) CRASH-2 2010 WOMAN 2017				Mortality, due to bleeding	644/20094 (3.2) 489/10060 (4.9) 155/10034 (1.5)	764/20044 (3.8) 574/10067 (5.7) 190/9977 (1.9)	RR 0.84 (0.76, 0.93) ^c RR 0.85 (0.76, 0.96) RR 0.81 (0.66, 1.00)	Favours TXA $p = 0.001^{\circ}$ No significant heterogeneity $l^2 = 0\% (p = 0.69)$

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
						Based on a logistic r found improved sur favouring immediat 1.42, 2.10; $p < 0.0001$). until 150 to 200 min estimates), at which (OR 1.00).	egression simulation vival (absence of de e administration of . Thereafter, survivan utes treatment delate point the models e	on model, the authors eath from bleeding) TXA (OR 1.72, 95% CI I benefits decreased ay (95% CI lower and estimated no benefit	Nonlinear association with increasing delay d p = 0.0109
	N = 40 138 (2 RCTs) CRASH-2 2010 WOMAN 2017				Mortality, not due to bleeding	1046/20094 (5.2) 974/10060 (9.7) 72/10034 (0.7)	1104/20044 (5.5) 1039/10067 (10.3) 65/9977 (0.7)	RR 0.95 (0.87, 1.03) ^c RR 0.94 (0.86, 1.02) RR 1.10 (0.79, 1.54)	No significant difference p = 0.18 ° No significant heterogeneity $l^2 = 0\%$ ($p = 0.36$)
	N = 40 138 (2 RCTs) CRASH-2 2010 WOMAN 2017				Mortality due to vascular occlusive event	43/20094 (0.2) 33/10060 (0.3) 10/10034 (0.1)	59/20044 (0.3) 48/10067 (0.5) 11/9977 (0.1)	OR 0-73 (0-49, 1-09) OR 0-69 (0-44, 1-08) OR 0-90 (0-38, 2-12)	No significant difference p = 0.1204 No significant heterogeneity $l^2 = NR (p = 0.5956)$
Trauma setting	7								
Al-Jeabory 2021 - SR High quality	N = 29 115 (3 RCTs and 11 Coh) CRASH-2 2010 (RCT) Guyette 2020 (RCT) Kakaei 2017 (RCT) EI-Menyar 2020 Howard 2017 Lipsky 2014	Mix of combat and civil trauma patients Civil Civil Civil Civil Combat Combat	Multiple countries Multi-country USA Iran Qatar USA Israel Afghanistan	TXA vs no TXA	In-hospital mortality	2099/13559 (15.5) 1463/10060 (14.5) 37/447 (8.3) 3/30 (10) 25/102 (24.5) 82/849 (9.7) 6/26 (23.1)	2547/15556 (16.4) 1613/10067 (16.0) 43/453 (9.5) 4/30 (13.3) 30/102 (29.4) 271/2924 (9.3) 0/10	OR 0.81 (0.62, 1.06) OR 0.89 (0.83, 0.96) OR 0.86 (0.54, 1.36) OR 0.72 (0.15, 3.54) OR 0.78 (0.42, 1.45) OR 1.05 (0.81, 1.36) OR 6.66 (0.34, 129.92)	No significant difference p = 0.12 Significant heterogeneity $I^2 = 83\%$ ($p < 0.00001$)
	Morrison 2012 Myers 2019 Neeki 2017 Neeki 2018 Rivas 2021 Swendsen 2013 Valle 2014 Wafaisade 2016	Combat Civil Civil Civil Civil Civil Civil Civil	USA USA USA USA USA USA Germany			148/293 (50.5) 136/189 (72.0) 8/128 (6.3) 13/362 (3.6) 106/654 (16.2) 9/52 (17.3) 25/109 (22.9) 38/258 (14.7)	218/603 (36.2) 161/189 (85.2) 13/125 (10.4) 30/362 (8.3) 91/254 (35.8) 17/74 (23.0) 14/105 (13.3) 42/258 (16.3)	OR 1.80 (1.36, 2.39) OR 0.45 (0.27, 0.74) OR 0.57 (0.23, 1.44) OR 0.41 (0.21, 0.80) OR 0.35 (0.25, 0.48) OR 0.70 (0.29, 1.73) OR 1.93 (0.94, 3.97) OR 0.89 (0.55, 1.43)	
Almuwallad 2021 SR High quality	N = 2140 (1 RCT and 2 Coh) Guyette 2020 (RCT) Wafaisade 2016 Neeki 2018	Civilian trauma patients	Trauma, MC USA Germany	TXA vs no TXA	Mortality, 24 hour	38/1067 (3.6) 16/447 (3.6) 15/258 (5.8) 7/362 (19)	62/1073 (5.8) 17/453 (3.8) 32/258 (12.4) 13/362 (3.6)	OR 0.60 (0.37, 0.99) OR 0.95 (0.47, 1.91) OR 0.44 (0.23, 0.83) OR 0.53 (0.21, 1.34)	No significant difference p = 0.05 Minimal heterogeneity $l^2 = 27\% (p = 0.26)$

Study ID	udy ID Sample size Patient Setting				Outcome	Results				
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
	N = 2143 (1 RCT and 2 Coh) Guyette 2020 (RCT) Wafaisade 2016 Neeki 2018		Multiple countries USA Germany USA		Mortality, 28 to 30 days	85/1062 (8.0) 36/442 (8.1) 36/258 (14.0) 13/362 (4.0)	117/1072 (10.9) 45/452 (10) 42/258 (16.3) 30/362 (8.3)	OR 0.69 (0.47, 1.02) OR 0.80 (0.51, 1.27) OR 0.83 (0.51, 1.35) OR 0.41 (0.21, 0.80)	No significant difference p = 0.06 Minimal heterogeneity $l^2 = 38\% (p = 0.20)$	
El-Menyar 2018 SR <i>Moderate</i> <i>quality</i>	N = 769 (2 Coh) Wafaisade 2016 Neeki 2017	Adult patients with traumatic injury presenting to the emergency department requiring blood	Prehospital (air rescue)	TXA vs placebo	Mortality, 24 hour	20/386 (5.2)	41/383 (10.7)	OR 0.49 (0.27, 0.84) OR 0.47 (0.25, 0.89) OR 0.54 (0.18, 1.66)	Favours TXA p = NR No significant heterogeneity $l^2 = 0\% (p = 0.82)$	
	N = 769 (2 Coh) Wafaisade 2016 Neeki 2017	transfusion			Mortality, 30-day	44/386 (11.4)	55/383 (14.4)	OR 0.86 (0.56, 1.32) OR 0.86 (0.53, 1.38) OR 0.87 (0.32, 2.32)	No significant difference p = NR No significant heterogeneity $l^2 = 0\% (p = 0.98)$	
Cannon 2017 SR <i>Moderate</i> <i>quality</i>	N = 21 666 (1 RCT, 2 Coh) CRASH-2 2010 (RCT) Cole 2015 Morrison 2013	Patients with severe trauma at risk of death from haemorrhage* *patients requiring blood transfusion and/or with an injury score greater than 25	Civilian and military trauma	TXA vs no TXA	Mortality, timing not specified	1550/10616 (14.6) 1463/10050 (14.6) 30/160 (18.8) 57/406 (14.0)	1828/11050 (16.5) 1613/10067 (16.0) 36/225 (16) 179/758 (23.6)	RR 0.70 (0.54, 1.20) OR 0.81 (0.54, 1.20) OR 0.89 (0.83, 0.96) OR 1.21 (0.71, 2.07) OR 0.53 (0.38, 0.73)	No significant difference p = 0.29 Substantial heterogeneity $l^2 = 82\% (p < 0.04)$ p = 0.004 p = 0.48 p = 0.0001	
Myers 2019 Retrospective Coh Serious risk of bias	N = 378	Patients presenting to a level 1 trauma centre	Level 1 trauma centre (NR)	TXA within 3 hours of presentation vs No TXA	Survival	136/189 (72)	161/189 (85)	aOR 0.86 (0.23, 3.25)	No significant difference p = 0.83	
Huebner 2017 SR <i>Critically low</i> <i>quality</i>	N = 6797 (2 Coh) Wafaisade 2016 Harvin 2015	Patients with trauma at risk of death from haemorrhage	Trauma Air Rescue (German) SC (US)	TXA vs placebo or no TXA	Mortality, all- cause within 24 hours	NR/NR (5.8) NR/98 (34)	NR/NR (12.8) NR/924 (10)	NR aOR 1.92 (1.05, 3.25)	Favours TXA, p = 0.01 Favours placebo, p = 0.035	
	N = 896 (1 Coh) MATTERs 2012		Military trauma (US)		Mortality, within 48 hours	NR/NR (NR)	NR/NR (NR)	RD 6.6 (NR)	Favours TXA, p = 0.004	
	N = 7693 (3 Coh)		Trauma		Mortality, In- hospital <i>MT subgroup</i>					

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
	MATTERs 2012 Wafaisade 2016 Harvin 2015		Military (US) Air Rescue (German) SC (US)	-		NR/NR (NR) NR/NR (14.4) NR/NR (14.7) NR/98 (40)	NR/NR (NR) NR/NR (28.1) NR/NR (16.3) NR/924 (17)	RD 6.5 (NR) RD 13.7 (NR) RR 0.49 (NR) NR aOR 0.74 (0.38, 1.40)	Favours TXA, p = 0.03 Favours TXA, p = 0.04 Favours TXA, p = NR Favours TXA, p = NR
	N = NR (4 Coh) Eckart 2014 Valle 2014 Swendsen 2013 Cole 2015		Trauma (Afghanistan) (UK) (US)	-	Mortality, timing not specified Swendsen 2013 re- analysis (N = NR)	NR/NR (15) NR/NR (27) NR/NR (5.8) NR/NR (4.3)	NR/NR (9) NR/NR (17) NR/NR (17.6) NR/NR (19.1)	aOR 0.27 (0.85, 0.89) NR NR NR	Favours TXA, $p = 0.03$ Favours placebo, $p = 0.024$ Favours TXA, $p = 0.05$ Favours TXA, $p = 0.03$
	N = 20 211 (1 RCT) CRASH-2 2010		Adult trauma MC (over 40		Mortality, within 4 weeks of injury	NR/NR (INR)	NR/NR (NR)	RR 0.91 (0.85, 0.97)	Favours TXA, p = 0.0035
	N = 20 211 (1 RCT) CRASH-2 2010		countries)		Mortality due to bleeding TXA within 1 hour 1 to 3 hours after 3 hours	NR/NR (5.3) NR/NR (4.8) NR/NR (4.4)	NR/NR (7.7) NR/NR (6.1) NR/NR (3.1)	RR 0.68 (0.57, 0.82) RR 0.79 (0.64, 0.97) RR 1.44 (1.12, 1.84)	Favours TXA, p < 0.0001 Favours TXA, p = 0.03 Favours placebo, p = NR
Ker 2015 SR High quality	N = 20 367 (2 RCTs) CRASH-2 2010 Yutthakasomsunt 2013	Adult trauma patients with, or at risk of, significant bleeding, including patients with moderate to severe TBI	MC (over 40 countries, Thailand)	IV TXA vs standard care	Mortality, all- cause <i>All trauma</i>	1475/10180 (14.5) 1463/10060 (14.5) 12/120 (10)	1631/10187 (16.0) 1613/10067 (16.0) 18/120 (15)	RR 0.90 (0.85, 0.97) RR 0.91 (0.85, 0.97) RR 0.67 (0.34, 1.32)	Favours TXA p = 0.003 No significant heterogeneity $l^2 = 0 (p = 0.38)$
Ausset 2015 SR <i>Critically low</i> <i>quality</i>	N = 20 896 (1 RCT, 2 Coh) CRASH-2 2010 Valle 2014 Cole 2014	Adult trauma patients with wide range of injury severities	Civilian trauma MC (over 40 countries, US)	TXA vs no TXA	Mortality, overall	NR NR (14.5) NR/150 NR/160 (8)	NR NR (16) NR/150 NR/225 (8)	NR aRR 0.015 NR NR	Meta-analysis not conducted $p = NR, Favours TXA \circ$ p = NR, Not significant f $p = NR, Not significant g$
						Cole 2014 noted a m patients with shock 0.16 (0.31, 0.86)	nultivariate analysis that revealed an ef	in the subgroup of fect favouring TXA OR	-
	N = 2228 (2 Coh) Morrison 2012 Morrison 2013		Combat trauma, SC (Afghanistan)		Mortality, overall	NR/293 (17.4) NR	NR/603 (23.9) NR	OR 0.61 (0.42, 0.89 NR	Favours TXA ^h p = NR, Not significant ⁱ
						Morrison 2012 noted patients requiring a	l an effect favouring massive transfusio	g TXA in a subgroup of n	
						NR (14.4)	NR (28.1)	OR 7.2 (3.0, 17.3)	p = NR, Favours TXA

Study ID	Sample size	Patient	Setting C	Comparison	Outcome	Results				
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Medical emerg	jency								·	
Bennett 2014 SR <i>Low</i>	N – 301 (3 RCT5)³ Bagnenko 2011 Bergqvist 1980 Engqvist 1979	Adult patients with severe or massive upper gastrointestinal bleeding	SC (Russia, Sweden, S tockholm)	IV and oral TXA vs placebo or no TXA	Mortality, all- cause	15/149 (10.1) 1/22 (4.5) 3/26 (12) 11/102 (10.8)	20/152 (13.2) 3/25 (12) 5/25 (20) 12/102 (11.8)	RR 0.77 (0.41, 1.46)* RR 0.38 (0.04, 3.38) RR 0.59 (0.16, 2.26) RR 0.92 (0.42, 1.98)	No significant difference p = 0.42.° No significant heterogeneity t ² = 0 (p = 0.69)	
HALT-IT 2020 RCT <i>Low risk of</i>	N = 11 937	Adult patients with severe or massive gastrointestinal	MC (UK, Pakistan, Nigeria, Egypt, Malaysia, Georgia,	IV TXA vs placebo	Mortality, all- cause	564/5956 (9.5)	548/5981 (9.2)	RR 1.03 (0.92, 1.16)	<i>No significant difference</i> <i>p</i> = NR	
bias		bleeding	Romania, Nepal, Sudan, Saudi Arabia, Spain, Ireland, Albania, Papua		Mortality due to bleeding, within 24 hours	124/5956 (2.1)	120/5981 (2.0)	RR 1.04 (0.81, 1.33)	<i>No significant difference</i> <i>p</i> = NR	
			Albania, Papua New Guinea, and Australia)		Mortality due to bleeding, within 28 days	253/5956 (4.2)	262/5981 (4.4)	RR 0.97 (0.82, 1.15)	<i>No significant difference</i> <i>p</i> = NR	
Obstetrics and	I maternity setting									
Della-Corte 2020 SR	N = 14 335 (2 RCTs) Ducloy-Bouthors 2011 WOMAN 2017	Patients with postpartum haemorrhage	Obstetrics France Multiple countries	TXA vs placebo or not treatment	Mortality (maternal) due to bleeding	110/7155 (1.5) 0/72 110/7083 (1.6)	135/7180 (1.9) 0/72 135/7108 (1.9)	RR 0.82 (0.64, 1.05)	NR	
quality	N = 14 335 (2 RCTs) Ducloy-Bouthors 2011 WOMAN 2017		Obstetrics France Multiple countries		Mortality (maternal), all- cause	148/7155 (2.1) 0/72 148/7083 (2.1)	172/7180 (2.4) 0/72 172/7108 (2.4)	RR 0.86 (0.69, 1.07)	NR	
Shakur 2018 SR	N = 20 172 (2 RCTs) WOMAN 2017	Women after birth following a		IV TXA vs placebo or	Mortality (maternal), all-	227/10036 (2.3)	256/9985 (2.6)	RR 0.88 (0.74, 1.05)	<i>No significant difference</i> <i>p</i> = 0.16	
High quality	Ducloy-Bouthers 2011	pregnancy of at least 24 weeks' gestation with PPH, regardless of mode of bitth or		standard care	cause	All estimable data ar were zero events in ((N = 151)	a are from one study (WOMAN 2017). There in either group in Ducloy-Bouthers 2011		Heterogeneity NA	
	N = 20 011 (1 RCT) WOMAN 2017	other aspects of third stage management			Timing from birth Less than 1 hr 1 to 3 hrs More than 3 hrs	80/4846 (1.7) 57/2674 (2.1) 90/2514 (3.6)	80/4726 (1.7) 83/2682 (3.1) 92/2569 (3.6)	RR 0.98 (0.72, 1.33) RR 0.69 (0.49, 0.96) RR 1.00 (0.75, 1.33)	p = 0.87 No difference p = 0.028 Favours TXA p = 1.0 No difference	
	N = 20 172 (2 RCTs)					155/10036 (1.5)	191/9985 (1.9)	RR 0.81 (0.65, 1.00)	Favours TXA p = 0.046	

Study ID	Sample size	Patient population *Estimated blood loss after vaginal birth > 500 mL, or > 1000 mL after caesarean section or estimated blood loss enough to compromise	Setting	Comparison	Outcome	Results				
Study designª	(no. of trials) included in analysis		(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
	WOMAN 2017 Ducloy-Bouthers 2011		Obstetrics (France, UK, Nigeria, Pakistan, Uganda,		Mortality (maternal), due to bleeding	All estimable data an were zero events in ((N = 151)	re from one study (either group in Duc	WOMAN 2017). There cloy-Bouthers 2011	Heterogeneity NA	
	N = 20 011 (1 RCT) WOMAN 2017	or estimated blood loss enough to compromise the haemodynamic status of the woman	n Pakistan, Uganda, ss Kenya, Cameroon, Sudan, Tanzania, Nepal, Zambia, Albania, Democratic Republic of Congo, Bangladesh, Ethiopia, Burkina Faso, Jamaica, Ghana, Papua New Guinea, Egypt, Colombia and		Timing from birth Less than 1 hr 1 to 3 hrs More than 3 hrs	49/4846 (1.0) 40/2674 (1.5) 66/2514 (2.6)	60/4726 (1.3) 67/2682 (2.5) 63/2569 (2.5)	RR 0.80 (0.55, 1.16) RR 0.60 (0.41, 0.88) RR 1.07 (0.76, 1.51)	<i>p</i> = 0.23 No difference <i>p</i> = 0.096 <i>Favours TXA</i> <i>p</i> = 0.70 No difference	
Paediatrics										
Huebner 2017 SR <i>Critically low</i> <i>quality</i>	N = 766 (1 Coh) Eckart 2014, adjusted for confounders	Patients with trauma at risk of death from haemorrhage Patients were predominantly male, mean age 11 years with penetrating trauma	Paediatric trauma (Afghanistan)	TXA vs no TXA	Mortality, timing not specified	NR/NR (15)	NR/NR (9)	OR 0.27 (0.85, 0.89)	Favours TXA, p = 0.03	

Studies with strikethrough do not meet the PICO criteria for this question.

aOR, adjusted odds ratio; aRR, adjusted risk ratio; CI, confidence interval; Coh, cohort study; hrs, hours; ICH, intracranial haemorrhage; IU, international units; IV, intravenous; MC, multicentre; M-H, Mantzel-Hentzel; NR, not reported; OR, odds ratio; RCT, randomised controlled trial; RD, risk difference; RR, relative risk; SC, single centre; TBI, traumatic brain injury; TXA, tranexamic acid; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic review studies with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. Due to missing data, the analysis excludes 4 patients in the intervention group and 4 patients in the placebo group in CRASH-2; a further 109 patients in the WOMAN trial were excluded (50 in the intervention group and 59 in the placebo group) as time to treatment was greater than 24 hours. All models were controlled for systolic blood pressure (5 mm Hg interval) and age (10-yr intervals), which are strong risk factors for death due to bleeding.

e. Ausset 2015 noted a post-hoc analysis of CRASH-2 had revealed when TXA was administered within 1 hour after trauma, mortality was reduced by one-third. Between hours 1–3, mortality was reduced by one-fifth. When given after the third hour, mortality due to bleeding appeared to increase.

- f. The study by Valle (2014) is confounded. Ausset 2015 noted that mortality was higher in the TXA group, but that the propensity score failed to account for important variables, resulting in the TXA group being more severely injured than the control group. No multivariate analysis was performed to account for these differences.
- g. The survival benefit of TXA in Cole 2014 is confounded. Patients who received TXA had higher ISS, incidence of shock (base deficit > 6 mEq/L) and transfusion requirements. A multivariate analysis in the subgroup of patients with shock revealed an effect favouring TXA OR 0.16 (0.31, 0.86).
- h. Ausset 2015 noted that the survival benefit of TXA in Morrison 2012 is confounded by the retrospective study design, with cryoprecipitate used more often in the TXA massive transfusion group. Factors significantly associated with death in the entire cohort included: Glasgow Coma Score of 8 or less, hypotension and coagulopathy.
- i. Propensity score adjusted for predictors of mortality, including RBC, FFP and plasma. After adjustment for platelet administration the OR was 0.62 (95% CI 0.43, 0.90). Ausset 2015 noted that the survival benefit of TXA in Morrison 2013 remained confounded by the heterogeneous use of rFVIIa.
- j. Bennett (2014) meta-analysed 8 RCTs involving patients with upper gastrointestinal bleeding. Only 3 RCTs were in patients with critical bleeding but the studies were confounded by the administration of oral TXA (in combination with IV TXA) and were not reflective of current standard of care.

4.8.3.2 Morbidity

A summary of the evidence relating to morbidity (e.g. vascular events, multiple organ failure, acute respiratory distress syndrome) associated with TXA in patients with critical bleeding is presented in Table 4.71.

Vascular events (any)

In a meta-analysis of data from studies included in this review (see Figure 4.47), the RCT evidence in critically bleeding trauma patients (CRASH-2) suggested there was little to no difference on the incidence of vascular events in trauma patients who received TXA (168/10 060, 1.67%) compared with those who did not receive TXA (201/ 10 067, 1.99%) (RR 0.84, 95% CI 0.68, 1.02; p = 0.08; random effect) (*GRADE: very low*).

Among the cohort studies conducted in critically bleeding trauma patients, the risk of vascular events was higher among those who received TXA (106/1801, 5.89%) compared with those who did not receive TXA (122/ 3157, 3.86%) (RR 1.63; 95%Cl 1.17, 2.29; p = 0.004, $l^2 = 23\%$) (*GRADE: Very low*). Noting there was a wide variety of injury severity and bleeding risk in the included studies, with the likelihood a missing data relating to inconsistencies in the measurement of the outcome.

In patients with acute gastrointestinal bleeding, the RCT evidence (HALT-IT) suggested that the risk of any thromboembolic event was similar among those who received TXA (86/5952, 1.4%) compare with those who did not receive TXA (72/5977, 1.2%) (RR 1.2, 95% CI 0.88, 1.64; p = 0.25, random effect) (see Figure 4.47). It was noted that the risk for venous thromboembolic events (DVT, PE) appeared to be higher among those who received TXA (48/5952, 0.8%) compared with those who did not receive TXA (26/5977, 0.4%) (RR 1.85; 95% CI 1.15, 2.98; p = 0.01, random effect) (*GRADE: Low*) (see Figure 4.48). The authors noted a similar risk was observed when patients who did not received the maintenance dose of TXA were excluded from the analysis (42 vs 20 events; RR 2.11; 95% CI 1.24, 3.59). The risk of arterial thromboembolic events (MI, stroke) was similar across groups (RR 0.7% vs 0.8%; RR 0.92, 95% CI 0.60, 1.39; (*GRADE: Low*).

In the obstetric setting, the RCT evidence (WOMAN) suggested there was little to no difference on the incidence of vascular events in women with major obstetric haemorrhage who received TXA (31/10 034, 0.31%) compared with those who did not receive TXA (34/ 9977, 0.34%) (RR 0.91, 95% CI 0.56, 1.47; p = 0.69; random effect) (*GRADE: very low*).

Organ failure

One RCT (WOMAN 2017) in the obstetric setting reported on other morbidity outcomes that were considered in this review. The data (see Table 4.49) suggested this is no differences between women with major obstetric haemorrhage who received TXA compared with those who did not for the outcomes of multiple organ failure (RR 0.94, 95% CI 0.71, 1.23; p = 0.65; random effect), respiratory failure (RR 0.87, 95% CI 0.67, 1.12; p = 0.27; random effect), or renal failure (RR 1.09; 95% CI 0.85, 1.39; p = 0.51; random effect) (*GRADE: very low*).

Figure 4.47 Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, vascular events (any)

	Tranexami	c acid	Cont	rol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% CI
1.3.1 Trauma (RCTs)							
CRASH-2 2010 (Civilian)	168	10060	201	10067	100.0%	0.84 [0.68, 1.02]	
Subtotal (95% CI)		10060		10067	100.0%	0.84 [0.68, 1.02]	\bullet
Total events	168		201				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.72 (P = 0.08	3)						
1.3.2 Trauma (Coh)							
Swendsen 2013 (adult trauma)	6	52	0	74	1.3%	18.40 [1.06, 319.58]	
Neeki 2017 (prehospital)	2	362	2	362	2.8%	1.00 [0.14, 7.06]	
Morrison 2012 (MATTERS)	8	293	2	603	4.3%	8.23 [1.76, 38.52]	
Wafaisade 2016 (prehospital, civilian)	4	71	10	121	7.5%	0.68 [0.22, 2.09]	
El-Menyar 2020 (Civilian)	9	102	5	102	8.3%	1.80 [0.62, 5.19]	
Cole 2015 (Civilian, ISS 15)	8	160	9	225	10.2%	1.25 [0.49, 3.17]	
Harvin 2014 (adult trauma)	6	98	41	934	12.1%	1.39 [0.61, 3.20]	
Guyette 2020 (STAAMP) (Civilian)	25	447	14	453	17.3%	1.81 [0.95, 3.44]	
Haren 2014 (adult trauma)	9	27	25	94	17.6%	1.25 [0.67, 2.35]	
Myers 2019 (Civilian)	29	189	14	189	18.5%	2.07 [1.13, 3.79]	
Subtotal (95% CI)		1801		3157	100.0%	1.63 [1.17, 2.29]	◆
Total events	106		122				
Heterogeneity: Tau ² = 0.06; Chi ² = 11.65	5, df = 9 (P =	0.23); l ² =	= 23%				
Test for overall effect: Z = 2.86 (P = 0.00	04)						
1.3.3 Medical emergency (RCT)							
Roberts 2020 (HALT-IT)	86	5952	72	5977	100.0%	1.20 [0.88, 1.64]	
Subtotal (95% CI)		5952		5977	100.0%	1.20 [0.88, 1.64]	◆
Total events	86		72				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.15 (P = 0.25	5)						
1.3.5 Obstetrics and maternity							
Ducloy-Bouthers 2011	0	72	0	72		Not estimable	
WOMAN 2017 (PPH)	31	10034	34	9977	100.0%	0.91 [0.56, 1.47]	
Subtotal (95% CI)		10106		10049	100.0%	0.91 [0.56, 1.47]	
Total events	31		34				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.40 (P = 0.69	9)						
						-	
							Favours TXA Favours placebo or no T

Test for subgroup differences: $Chi^2 = 12.43$, df = 3 (P = 0.006), l² = 75.9%

Figure 4.48 Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, venous and arterial events (GI bleeding)

	Tranexamic	acid	Contr	ol		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	1	M-H, Random, 95% Cl
1.4.1 Venous events (PE	and DVT)							
Roberts 2020 (HALT-IT) Subtotal (95% CI)	48	5952 5952	26	5977 5977	100.0% 1 00.0%	1.85 [1.15, 2.98] 1.85 [1.15, 2.98]		-
Total events Heterogeneity: Not applica	48 ble		26					
Test for overall effect: Z = 2	2.54 (P = 0.01))						
1.4.2 Arterial events (MI,	stroke)							
Roberts 2020 (HALT-IT) Subtotal (95% CI)	42	5952 5952	46	5977 5977	100.0% 1 00.0%	0.92 [0.60, 1.39] 0.92 [0.60, 1.39]		↓
Total events Heterogeneity: Not applica Test for overall effect: Z = 0	42 ble 0.41 (P = 0.68))	46					
							0.02	0.1 1 10 50 Favours TXA Favours placebo or no TXA

Figure 4.49 Forest plot of comparison: TXA vs no TXA, outcome: Morbidity, other (obstetrics)

	Tranexami	c acid	Cont	rol	Risk Ratio			Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C		M-H, Random, 95% Cl	
1.4.2 multiple organ fail	ure								
Ducloy-Bouthers 2011	0	77	0	74		Not estimable			
WOMAN 2017 (PPH)	99	10032	105	9985	100.0%	0.94 [0.71, 1.23]			
Subtotal (95% CI)		10109		10059	100.0%	0.94 [0.71, 1.23]		\bullet	
Total events	99		105						
Heterogeneity: Not applic	able								
Test for overall effect: Z =	= 0.46 (P = 0	.65)							
1.4.3 respiratory failure									
WOMAN 2017 (PPH)	108	10033	124	9985	100.0%	0.87 [0.67, 1.12]			
Subtotal (95% CI)		10033		9985	100.0%	0.87 [0.67, 1.12]		\bullet	
Total events	108		124						
Heterogeneity: Not applic	able								
Test for overall effect: Z =	= 1.09 (P = 0	.27)							
1.4.5 renal failure									
Ducloy-Bouthers 2011	0	77	0	74		Not estimable			
WOMAN 2017 (PPH)	129	10033	118	9985	100.0%	1.09 [0.85, 1.39]			
Subtotal (95% CI)		10110		10059	100.0%	1.09 [0.85, 1.39]		•	
Total events	129		118						
Heterogeneity: Not applic	able								
Test for overall effect: Z =	= 0.67 (P = 0	.51)							
							+		
							0.050	.2 1 5	20
Test for subgroup differer	nces: Chi² =	1.61, df =	2 (P = 0	.45), l² =	0%		Favours	experimental] Favours [control]	

Table 4.71 Results for TXA versus no TXA: Patients with critical bleeding - Morbidity

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Any setting									
Ageron 2020 SR <i>Moderate</i> <i>quality</i>	N = 28 333 (2 RCTs) CRASH-2 WOMAN	CRASH-2 are trauma patients, the WOMAN trials are women with a postpartum haemorrhage	Trauma and Obstetrics	TXA vs placebo	Any vascular occlusive events	118/14270 (0.01) NR NR The authors stratifi risk of death due bl vascular occlusive e baseline risk catego	152/14063 (0.01) NR NR ed individual patie eeding and found events with TXA an ories (0 = 0.25)	NR NR nt data by baseline no increased risk of id it did not vary by	No significant difference p = 0.255
	N = 28 333 (2 RCTs) CRASH-2 WOMAN				Myocardial infarction	24/14270 (0.00) NR NR	46/14063 (0.00) NR NR	NR NR NR	No significant difference <i>p</i> = 0.909
	N = 28 333 (2 RCTs) CRASH-2 WOMAN				Stroke	32/14270 (0.00) NR NR	42/14063 (0.00) NR NR	NR NR NR	No significant difference <i>p</i> = 0.152
	N = 28 333 (2 RCTs) CRASH-2 WOMAN	-			Pulmonary embolism	54/14270 (0.00) NR NR	56/14063 (0.00) NR NR	NR NR NR	No significant difference p = 0.739
	N = 28 333 (2 RCTs) CRASH-2 WOMAN				Deep vein thrombosis	28/14270 (0.00) NR NR	30/14063 (0.00) NR NR	NR NR NR	No significant difference p = 0.214
Chornenki 2019 SR High quality	N = 42 808 (5 RCTs) CRASH-2 2010 Yutthakasomsunt 2013 Arulkumaran 2017 Sprigg 2018		Trauma, Obstetrics and Medical (No countries listed)	TXA vs placebo or no TXA	Stroke	85/21424 (0.4) 6/50 (12.0) 55/10060 (0.5) 0/120 (0) 8/10033 (0.1) 16/1161 (1.4)	88/21384 (0.4) 2/50 (4.0) 66/10067 (0.7) 3/118 (2.5) 6/9985 (0.1) 11/1164 (0.9)	RR 1.10 (0.68, 1.78) RR 3.00 (0.64, 14.16) RR 0.83 (0.58, 1.19) RR 0.14 (0.01, 2.69) RR 1.33 (0.46, 3.82) RR 1.46 (0.68, 3.13)	No significant difference p = 0.71 Mild heterogeneity $l^2 = 31\%$ ($p = 0.21$)

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
	N = 42 470 (3 RCTs) CRASH-2 2010 Arulkumaran 2017 Sprigg 2018	Patients requiring treatment or prevention of postpartum haemorrhage, intracranial			Myocardial infarction	48/21254 (0.2) 35/10060 (0.3) 2/10033 (0.0) 11/1161 (0.9)	64/21216 (0.3) 55/10067 (0.5) 3/9985 (0.0) 6/1164 (0.5)	RR 0.88 (0.43, 1.84) RR 0.64 (0.42, 0.97) RR 0.66 (0.11, 3.97) RR 1.84 (0.68, 4.95)	No significant difference p = 0.74 Moderate heterogeneity $l^2 = 46\%$ ($p = 0.15$)
	N = 43 161 (6 RCTs) Chowdhary 1996 Tsementzis 1990 Ross 2000 CRASH-2 2010 Arulkumaran 2017 Sprigg 2018	haemorrhage, subarachnoid haemorrhage, traumatic brain injury or non- specific trauma injury			Pulmonary embolism	113/21598 (0.5) 1/65 (1.5) 2/50 (4.0) 1/229 (0.4) 72/10060 (0.7) 17/10033 (0.2) 20/1161 (1.7)	116/21563 (0.5) 1/64 (1.6) 1/50 (2.0) 0/233 (0) 71/10067 (0.7) 20/9985 (0.2) 23/1164 (2.0)	OR 0.97 (0.75, 1.26) OR 0.98 (0.06, 16.08) OR 2.04 (0.18, 23.27) OR 3.07 (0.12, 75.65) OR 1.01 (0.73, 1.41) OR 0.85 (0.44, 1.62) OR 0.87 (0.47, 1.59)	No significant difference p = 0.83 No significant heterogeneity $l^2 = 0 (p = 0.94)$
	N = 46 287 (6 RCTs) Tsementzis 1990 Shakur 2010 Sprigg 2014 Arulkumaran 2017 Sprigg 2018 Sentilhes 2010				Deep vein thrombosis	63/23164 (0.3) 0/50 (0) 40/10060 (0.4) 1/16 (6.25) 3/10033 (0.0) 19/1161 (1.6) 0/1844 (0)	66/23123 (0.3) 3/50 (6.0) 41/10067 (0.4) 0/8 (0) 7/9985 (0.1) 14/1164 (1.2) 1/1849 (0.1)	RR 0.14 (0.01, 2.70) RR 0.98 (0.63, 1.51) RR 1.59 (0.07, 35.15) RR 0.43 (0.11, 1.65) RR 1.36 (0.69, 2.70) RR 0.33 (0.01 (8.20)	<i>p</i> = NR
Gayet-Ageron 2018 SR High quality	N = 40 138 (2 RCTs) CRASH-2 2010 WOMAN 2017	Patients with acute severe bleeding	Trauma (MC, over 40 countries)	TXA vs placebo	Myocardial infarction (fatal and non-fatal)	37/20094 (0.2) 35/10060 (0.3) 2/10034 (0.0)	58/20044 (0.3) 55/10067 (0.5) 3/9977 (0.0)	OR 0-64 (0-43, 0-97) OR 0-64 (0-42, 0-98) OR 0-66 (0-11, 3-95)	Favours TXA p = 0.0371 No significant heterogeneity $l^2 = NR (p = 0.9788)$
	N = 40 138 (2 Patients with acute severe bleeding Obstetrics and maternity (MC, 21 countries) TXA v CRASH-2 2010 WOMAN 2017 V V V	TXA vs placebo	Deep vein thrombosis (fatal and non- fatal)	43/20094 (0.2) 40/10060 (0.4) 3/10034 (0.0)	48/20044 (0.2) 41/10067 (0.4) 7/9977 (0.1)	OR 0-90 (0-60, 1-36) OR 0-98 (0-63, 1-52) OR 0-42 (0-11, 1-64)	No significant difference p = NR No significant heterogeneity $I^2 = NR (p = 0.2483)$		
					Pulmonary embolism (fatal and non- fatal)	89/20094 (0.4) 72/10060 (0.7) 17/10034 (0.2) 65/20094 (0.3)	91/20044 (0.5) 71/10067 (0.7) 20/9977 (0.2) 72/20044 (0.4)	OR 0.98 (0.73, 1.32) OR 1.02 (0.74, 1.42) OR 0.84 (0.44, 1.61)	No significant difference p = NR No significant heterogeneity $l^2 = NR (p = 0.6025)$ No significant difference
					ou one	00/20004 (0.0)	12/20044 (0.4)	01(000,121)	no agrimeant amerence

Study ID	Study ID Sample size Patient Setting Comparison Outcome Results											
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b			
					(fatal and non- fatal)	57/10060 (0.6) 8/10034 (0.1)	66/10067 (0.7) 6/9977 (0.1)	OR 0-87 (0-61, 1-24) OR 1-32 (0-46, 3-81)	p = NR No significant heterogeneity $l^2 = NR (p = 0.4647)$			
Trauma settin	g											
Al-Jeabory 2021 SR High quality	N = 22 270 (5 studies)	Mix of combat and civilian trauma patients	Trauma (multiple countries)	TXA vs no TXA	Myocardial infarction	45/11288 (0.4)	64/10982 (0.6)	OR 0.66 (0.45, 0.97)	Favours TXA p = 0.03 No significant heterogeneity $l^2 = 0\%$			
	N = 22 270 (5 studies)				Stroke	73/11288 (0.6)	76/10982 (0.7)	OR 0.90 (0.65, 1.24)	No significant difference p = 0.50 Moderate heterogeneity $l^2 = 40\%$			
	N = 2271 (6 studies)				Thromboembolic events	67/1308 (5.1)	62/963 (6.4)	OR 0.89 (0.37, 2.11)	No significant difference p = 0.79 Moderate heterogeneity $l^2 = 60\%$			
	N = 25 912 (5 studies)				Pulmonary embolism	137/1211 (1.1)	117/1380 (0.8)	OR 1.57 (0.79, 3.13)	No significant difference p = 0.20 Significant heterogeneity $l^2 = 80\%$			
	N = 26 165 (6 studies)	-							Deep vein thrombosis	105/12240 (0.9)	105/13925 (0.8)	OR 1.13 (0.51, 2.51)
	N = 385 (1 study)				Coagulation failure	5/160 (3.1)	5/225 (2.2)	OR 1.42 (0.40, 4.99)	No significant difference p = 0.58 Heterogeneity NA			
	N = 1480 (3 studies)				Multiple organ failure	106/681 (15.6)	156/799 (19.5)	OR 0.87 (0.66, 1.16)	No significant difference p = 0.35 Moderate heterogeneity $l^2 = 39\%$			
	N = 1011 (2 studies)				Acute kidney failure	22/212 (10.4)	17/799 (2.1)	OR 1.97 (1.01, 3.86)	No significant difference p = 0.05			

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% Cl)	<i>Statistical significance p</i> -value Heterogeneity ^b
									No significant heterogeneity l² = 0%
	N = 385 (1 study)				Hepatic failure	5/160 (3.1)	2/225 (0.9)	OR 1.21 (0.81, 1.82)	No significant difference <i>p</i> = 0.35 Heterogeneity NA
	N = 186 (1 study)				Sepsis	4/67 (6.0)	8/119 (6.7)	OR 0.88 (0.26, 3.04)	No significant difference <i>p</i> = 0.84 Heterogeneity NA
	N = 385 (1 study)				Infection	89/160 (55.6)	113/225 (50.2)	OR 1.24 (0.83, 1.87)	No significant difference <i>p</i> = 0.30 Heterogeneity NA
Almuwallad 2021 SR <i>High quality</i> El-Menyar 2018 SR <i>Moderate</i> <i>quality</i>	N = 2020 (1 RCT, 3 Coh) N = 769 (2 Coh)	Civilian trauma patients Adult patients with traumatic injury presenting to the emergency department requiring blood	Trauma, MC Germany USA Qatar USA Prehospital (air rescue)	TXA vs no TXA TXA vs placebo	Thromboembolic events (Venous) Wafaisade 2016 Neeki 2018 EI-Menyar 2019 Guyette 2020 (RCT) Thromboembolic events (Venous) Wafaisade 2016 Neeki 2017	40/982 (4.0) 4/71 (5.6) 2/362 (0.6) 9/102 (8.8) 25/447 (5.6) 6/386 (1.55)	31/1038 (3.0) 10/121 (8.3) 2/362 0.6) 5/102 (4.9) 14/453 (3.1) 12/383 (3.1)	OR 1.49 (0.90, 2.46) OR 0.66 (0.20, 2.20) OR 1.00 (0.14, 7.14) OR 1.88 (0.61, 5.81) OR 1.86 (0.95, 3.62) OR 0.74 (0.27, 2.07) OR 0.67 (0.20, 2.22) OR 0.98 (0.14, 7.04)	No significant difference p = 0.12 No significant heterogeneity $l^2 = 0\% (p = 0.48)$ <i>No significant difference</i> p = NR No significant heterogeneity $l^2 = 0\% (p = 0.75)$
Nishida 2017 SR <i>Critically low</i> <i>quality</i>	N = 23 117 (2 RCTs, 6 Coh)	transfusion Patients with trauma-induced coagulopathy	MC (~40 countries)	TXA vs standard care	Thromboembolic events (Venous)	209/10881(1.9)	288/12236 (2.4)	RR 1.32 (0.80, 2.16)	No significant difference p = 0.28 Substantial heterogeneity $l^2 = 61\% (p = 0.02)$
					N = 20 365 (2 RCTs) Shakur 2010 Yutthakasemsunt 2013)	168/10180 (1.7) 168/10060 (1.7) 0/120	201/10185 (2.0) 201/10067 (2.0) 0/118	RR 0.84 (0.68, 1.02) RR 0.84 (0.68, 1.02) Not estimable	<i>No significant difference p</i> = 0.08 Heterogeneity NA
					N = 2752 (6 Coh)	41/701 (5.8)	87/2051 (4.2)	RR 1.61 (0.86, 3.01)	No significant difference p = 0.14 Substantial heterogeneity

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b
					Morrison 2012 Swendsen 2013 Haren 2014 Harvin 2014 Cole 2015 Wafaisade 2015	8/293 (2.7) 6/52 (11.5) 9/27 (3.3) 6/98 (6.1) 8/160 (5) 4/71 (5.6)	2/603 (0.3) 0/74 25/94 (26.6) 41/934 (4.4) 9/225 (4) 10/121 (8.2)	RR 8.23 (1.76, 38.52) RR 18.40 (1.06, 319.58) RR 1.25 (0.67, 2.35) RR 1.39 (0.61, 3.20) RR 1.25 (0.49, 3.17) RR 0.68 (0.22, 2.09)	I ² = 52% (<i>p</i> = 0.06)
Cannon 2017 SR <i>Moderate</i> quality	N = 21 408 (1 RCT, 2 Coh)	Patients with severe trauma at risk of death from haemorrhage* *patients requiring blood transfusion and/or with an injury score greater than 25	Civilian and military trauma	TXA vs no TXA	Thromboembolic events (Venous) CRASH-2 2010 Cole 2015 Morrison 2012	191/10513 (1.8) 168/10060 (1.7) 8/160 (5) 15/293 (5.1)	213/10895 (1.95) 201/10067 (2) 3/603 (0.5) 9/225 (4)	OR 2.00 (0.53, 7.50) RD 0.019 OR 0.83 (0.68, 1.03) OR 1.26 (0.48, 3.35) OR 10.79 (3.10, 37.58)	No significant difference p = 0.30 Substantial heterogeneity I ² = 88% (p = 0.0003)
Myers 2019 Retrospective Coh Serious risk of bias	N = 378	Patients presenting to a level 1 trauma centre	Level 1 trauma centre (NR)	TXA within 3 hours of presentation vs No TXA	Thromboembolic events (Venous)	29/189 (15.3)	14/189 (7.4)	OR 3.26 (1.3, -9.1)	Favours intervention p= 0.02
Huebner 2017 SR <i>Critically low</i>	N = 20 211 (1 RCT) CRASH-2	Patients with trauma at risk of death from	Civilian trauma	TXA vs placebo or no TXA	Vasco occlusive events	NR/NR (1.7)	NR/NR (2.0)	NR	No significant difference <i>p</i> = NR Heterogeneity NA
quality	N = NR (1 Coh) Cole 2015	naemorrnage			Thromboembolic events	NR/NR (8)	NR/NR (2)	NR	Favours placebo p = 0.01 Heterogeneity NA
	N = 126 (1 Coh) Swendsen 2013				DVT/PE Subgroup analysis	NR/NR (11.5) NR/NR (12)	NR/NR (0) NR/NR (0)	NR	Favours placebo, $p = 0.004$ p = 0.012 Heterogeneity NA
Ker 2015 SR High quality	N = 20 367 (2 RCTs) CRASH-2 2010 Yutthakasemsunt 2013	Adult trauma patients with, or at risk of, significant bleeding, including patients with moderate to severe TBI with wide range of injury severities	MC, 40 countries, Thailand	IV TXA vs standard care	Myocardial infarction	351/10180 (3.4) 35/10060 (0.3) 0/120	58/10187 (0.6) 55/10067 (0.5) 3/120 (2.5)	RR 0.61 (0.40, 0.92) RR 0.64 (0.42, 0.97) RR 0.14 (0.01, 2.74)	No significant difference p = 0.019 No significant heterogeneity $l^2 = 0 (p = 0.32)$
	N = 20 367 (2 RCTs)				Deep vein thrombosis	40/10180 (0.4) 40/10060 (0.4) 0/120	42/10187 (0.4) 41/10067 (0.4) 1/120 (0.8)	RR 0.95 (0.62, 1.47) RR 0.98 (0.63, 1.51) RR 0.33 (0.01, 8.10)	No significant difference p = 0.83 No significant heterogeneity $l^2 = 0 (p = 0.51)$

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis	рориацоп	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
	CRASH-2 2010 Yutthakasomsunt 2013									
	N = 20 367 (2 RCTs) CRASH-2 2010 Yutthakasomsunt 2013				Pulmonary embolism	72/10180 (0.7) 72/10060 (0.7) 0/120	71/10187 (0.7) 71/10067 (0.7) 0/120	RR 1.01 (0.73, 1.41) RR 1.01 (0.73, 1.41) Not estimable	<i>No significant difference</i> <i>p</i> = 0.93 Heterogeneity NA	
	N = 510 (2 RCTs) CRASH-2 2010 Yutthakasomsunt 2013				Stroke	0/253 0/133 0/120	1/257 (0.4) 1/137 (0.7) 0/120	RR 0.34 (0.01, 8.35) RR 0.34 (0.01, 8.35) Not estimable	<i>No significant difference</i> <i>p</i> = 0.51 Heterogeneity NA	
	N = 510 (2 RCTs) CRASH-2 2010 Yutthakasemsunt 2013				Deep vein thrombosis	0/253 0/133 0/120	3/257 (1.2) 2/137 (1.5) 1/120 (0.8)	RR 0.25 (0.03, 2.26) RR 0.21 (0.01, 4.25) RR 0.33 (0.01, 8.10)	No significant difference p = 0.22 No significant heterogeneity $l^2 = 0 (p = 0.83)$	
Ausset 2015 SR <i>Critically low</i>	N = 20 211 (1 RCT) CRASH-2 2010		Civilian trauma MC (over 40 countries,	TXA vs no TXA	Vaso-occlusive events, overall	NR (1.7)	NR (2.0)	NR	<i>No significant difference</i> <i>p</i> = NR	
quality			US)		Thromboembolic events (Venous)	NR	NR	NR	<i>No significant difference</i> <i>p</i> = NR	
					Pulmonary embolism	NR	NR	NR	<i>No significant difference</i> <i>p</i> = NR	
					Stroke	NR	NR	NR	<i>No significant difference</i> <i>p</i> = NR	
					Myocardial infarction	NR	NR	NR	Favours TXA p = NR	
	N = 385 (1 Coh) Cole 2014				Multiorgan failure	NR/160 (30)	NR/225 (37)	NR	<i>No significant difference</i> <i>p</i> = NR	
					Patients with shock	NR	NR	OR 0.27 (0.1, 0.73)	Favours TXA d	
	N = 896 (1 Coh) Morrison 2012		Combat trauma, SC (Afghanistan)	TXA vs no TXA	Thromboembolic events (Venous)	NR/293	NR/603	NR	<i>No significant difference</i> <i>p</i> = NR	
	N = 9127 (1 RCT) CRASH-3 2019				Haemorrhagic complications	16/4613 (0.3)	22/4514 (0.5)	RR 0.71 (0.37, 1.35)	No significant difference p = 0.30	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA n/N (%)	No TXA n/N (%)	Risk estimate (95% CI)	<i>Statistical significance p-value</i> Heterogeneity ^b
Medical emerg	gency								
Bennett 2014 SR High quality	N = 251 (2 RCT) ^e Bagnenko 2011 Engqvist 1979	Adult patients admitted with severe or major upper	SC (Russia, Stockholm)	IV and oral TXA vs placebo or no TXA	Any thromboembolic event	5/124 (4.0) 0/22 5/102 (4.9)	2/127 (1.6) 0/25 2/102 (2.0)	RR 2.50 (0.50, 12.59) Not estimable RR 2.50 (0.50, 12.59)	No significant difference p = 0.27 ^c
	N = 204 (1 RCT) ^e Engqvist 1979	gastrointestinal bleeding			DVT	4/102 (3.9)	2/102 (2.0)	RR 2.00 (0.37, 10.68)	No significant difference p = 0.42 °
	N = 204 (1 RCT) e Engqvist 1070 E 2020 N = 11.920 Adult patients with MC (UK, Dakis)			MI, PE and cerebral infarction	4/102 (3.9)	2/102 (2.0)	RR 2.00 (0.37, 10.68)	No significant difference p = 0.42 °	
HALT-IT 2020 RCT <i>Low risk of</i>	N = 11 929	Adult patients with acute gastrointestinal	MC (UK, Pakistan, Nigeria, Egypt, Malaysia, Georgia,	IV TXA vs placebo	Any thromboembolic event	86/5952 (1.4%)	72/5977 (1.2%)	RR 1.20 (0.88, 1.64)	No significant difference p = NR
bias		bleeding	Romania, Nepal, Sudan, Saudi Arabia, Spain,		Venous events (DVT, PE)	48/5952 (0.8)	26/5977 (0.4)	RR 1.85 (1.15, 2.98)	Favours no TXA p = NR
	Arabia, Spain, Ireland, Albania, Papu	Arabia, Spain, Ireland, Albania, Papua		Deep vein thrombosis	23/5952 (0.4)	12/5977 (0.2)	RR 1.92 (0.96, 3.86)	Favours no TXA p = NR	
			Albania, Papua New Guinea, and Australia)		Pulmonary embolism	28/5952 (0.5)	16/5977 (0.3)	RR 1.76 (0.95, 3.24)	Favours no TXA p = NR
					Arterial events (MI, stroke)	42/5952 (0.7)	46/5977 (0.8)	RR 0.92 (0.60, 1.39)	<i>No significant difference p</i> = NR
					myocardial infarction	24/5952 (0.4)	28/5977 (0.5)	RR 0.86 (0.50, 1.48)	<i>No significant difference p</i> = NR
					Stroke	19/5952 (0.3)	18/5977 (0.3)	RR 1.06 (0.56, 2.02)	<i>No significant difference p</i> = NR
Obstetric setti	ng								
Della-Corte 2021	N = 144 (1 RCT) Ducloy-Bouthors 2011	Patients with postpartum	Obstetrics France	TXA vs placebo or no treatment	Deep vein thrombosis	0/72	0/72	Not estimable	NR
SR <i>Moderate</i>		haemorrhage			Pulmonary embolism	0/72	0/72	Not estimable	NR
quality					Myocardial infarction	0/72	0/72	Not estimable	NR
					Stroke	0/72	0/72	Not estimable	NR
					Organ failure	0/72	0/72	Not estimable	NR
	N = 14 332 (2 RCTs)		Obstetrics		Surgical intervention	1379/7152 (19.3)	1453/7180 (20.2)	RR 0.95 (0.89, 1.02)	No significant difference p = NR

Study ID	Sample size	Patient	Setting Co	Comparison C	Outcome	Results				
Study design ^a	(no. of trials) included in	population	(Location)					Risk estimate	Statistical significance	
J	analysis					n/N (%)	n/N (%)	(95% CI)	<i>p</i> -value Heterogeneity ^b	
	Ducloy-Bouthors 2011 WOMAN 2017		France Multiple countries			4/72 (5.6) 1375/7080 (19.4)	5/72 (6.9) 1448/7108 (20.4)	NR NR	No significant heterogeneity l ² = 0%	
Shakur 2018 SR	N = 20 015 (1 RCT) WOMAN 2017	Women after birth following a	(France, UK, Nigeria, Pakistan,	IV TXA vs placebo or standard care	Serious maternal morbidity (any)	223/10030 (2.2)	224/9985 (2.2)	RR 0.99 (0.83, 1.19)	No significant difference p = 0.92	
High quality	N = 20 168 (2 RCTs) WOMAN 2017 Ducloy-Bouthers 2011	pregnancy of at least 24 weeks' gestation with PPH, regardless of mode	Uganda, Kenya, Cameroon, Sudan, Tanzania, Nepal, Zambia, Albania, Democratic Republic of Congo, Bangladesh, Ethiopia, Burkina Faso, Jamaica, Ghana, Papua New Guinea, Egypt, Colombia and Cote d'Ivoire)		Serious maternal morbidity, multiple organ failure	99/10109 (1.0) 99/10032 (1.0) 0/77	105/10059 (1/0) 105/9985 (1.1) 0/74	RR 0.94 (0.71, 1.23) RR 0.94 (0.71, 1.23) Not estimable	No significant difference <i>p</i> = 0.65	
	N = 20 169 (2 RCTs) WOMAN 2017 Ducloy-Bouthers 2011	aspects of third stage management			Serious maternal morbidity, renal failure	129/10220 (1.3) 129/10033 (1.3) 0/77	118/10059 (1.2) 118/9985 (1.2) 0/74	RR 1.09 (0.85, 1.39) RR 1.09 (0.85, 1.39) Not estimable	No significant difference <i>p</i> = 0.51	
	N = 20 169 (2 RCTs) WOMAN 2017 Ducloy-Bouthers 2011	*Estimated blood loss after vaginal birth > 500 mL, or > 1000 mL after caesarean section			Serious maternal morbidity, maternal seizure	33/10110 (0.3) 33/10033 (0.3) 0/77	43/10059 (0.4) 43/9985 (0.4) 0/74	RR 0.76 (0.49, 1.20) RR 0.76 (0.49, 1.20) Not estimable	No significant difference <i>p</i> = 0.24	
	N = 20 018 (1 RCT) WOMAN 2017	or estimated blood loss enough to compromise the haemodynamic status of the woman			Serious maternal morbidity, respiratory failure	108/10033 (1.1)	124/9985 (1.2)	RR 0.87 (0.67, 1.12)	No significant difference p = 0.27	
	N = 20 018 (1 RCT) WOMAN 2017				Serious maternal morbidity, cardiac arrest	110/10033 (1.1)	115/9985 (1.2)	RR 0.95 (0.73, 1.23)	No significant difference <i>p</i> = 0.71	
	N = 20 169 (1 RCT) WOMAN 2017				Serious maternal morbidity, hepatic failure	29/10033 (0.3)	30/9985 (0.3)	RR 0.96 (0.58, 1.60)	No significant difference <i>p</i> = 0.88	

Studies with strikethrough do not meet the PICO criteria for this question.

aOR, adjusted odds ratio; CI, confidence interval; Coh, cohort study; DVT, deep vein thrombosis; hrs, hours; IU, international units; IV, intravenous; MC, multicentre; M-H, Mantzel-Hentzel; NA, not applicable; NR, not reported; OR, odds ratio; PE, pulmonary embolism; PPH, postpartum haemorrhage; RCT, randomised controlled trials, RR, relative risk; SC, single centre; SD, standard deviation; TXA, tranexamic acid; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic review studies with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

d. The benefit of TXA in Cole 2014 is confounded. Patients who received TXA had higher ISS, incidence of shock (base deficit > 6 mEq/L) and transfusion requirements. A multivariate analysis in the subgroup of patients with shock revealed an effect favouring TXA OR 0.27 (0.1, 0.7).

e. The included studies were confounded by the administration of oral TXA (in combination with IV TXA) and were not reflective of current standard of care.

4.8.3.3 Blood loss

A summary of the evidence relating to blood loss associated with TXA in patients with critical bleeding is presented in Table 4.72.

None of the included RCTs or cohort studies were found to report reliable data relating to blood loss. The available evidence was therefore not further considered.

Table 4.72 Results for TXA versus no TXA: Patients with critical bleeding – Blood loss

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study designª	(no. of trials) included in analysis	population	(Location)			TXA Mean ± SD (n)	No TXA Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b
Obstetrics and	l maternity setting								
Shakur 2018 SR High quality	N = 151 (1 RCT) Ducloy-Bouthors 2011	Women after vaginal birth following a pregnancy of at	MC (France)	IV TXA vs placebo or standard care	Blood loss, 500 mL or more after randomisation	12/77 (15.6)	23/74 (31.1)	RR 0.50 (0.27, 0.93)	Favours TXA p = 0.029 Heterogeneity NA
	N = 151 (1 RCT) Ducloy-Bouthors 2011	least 24 weeks' gestation with PPH Estimated blood loss after vaginal birth > 800 mL			Blood loss, 1000 mL or more after randomisation	4/77 (5.2)	8/74 (10.8)	RR 0.48 (0.15, 1.53)	<i>No significant difference</i> <i>p</i> = 0.21 Heterogeneity NA
	N = 151 (1 RCT) Ducloy-Bouthors 2011				Mean blood loss, mL	280 ± 320 (n = 77)	387 ± 409 (n = 74)	MD -107.00 (-224.44, 10.44)	<i>No significant difference</i> <i>p</i> = 0.074 Heterogeneity NA

Cl, confidence interval; hrs, hours; IU, international units; MC, multicentre; MD, mean difference; M-H, Mantzel-Hentzel; NR, not reported; OR, odds ratio; RCT, randomised controlled trial; RR, relative risk; SD, standard deviation; TXA, tranexamic acid; UK, United Kingdom; US, United States; WMD, weighted mean difference

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic review studies with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

4.8.3.4 Transfusion volume

A summary of the evidence relating to transfusion volumes associated with TXA in patients with critical bleeding is presented in Table 4.73

Red blood cells

In a meta-analysis of data from studies included in this review (see Figure 4.50), the RCT evidence in critically bleeding trauma patients (CRASH-2) suggested there was little to no difference on the volume of RBC transfused in patients who received TXA (mean 6.06 units) compared with those who did not receive TXA (mean 6.29 units) (SMD –0.02, 95%CI –0.02, 0.02; p = 0.25; random effect) (*GRADE: Low*).

Among the cohort studies that reported data, the volume of RBC transfused was higher among patients who received TXA (range 4.42 units to 22 units) compared with those who did not receive TXA (range 2 to 16 units) (SMD 0.53; 95%Cl 0.22, 0.85; p = 0.001, $l^2 = 90\%$) (*GRADE: Very low*). Noting there was substantial heterogeneity with a wide variety of injury severity and bleeding risk in the included studies, with the results likely to differ after adjustments for confounders across all studies (e.g. patients who received TXA had higher incidence of shock, blood loss and transfusion needs).

In patients with acute gastrointestinal bleeding, the RCT evidence (HALT-IT) suggested there was little to no difference on the volume of RBC transfused in patients who received TXA (mean 2.8 units) compared with those who did not receive TXA (mean 2.9 units transfused) (MD –0.10, 95%CI –0.21, 0.01; p = 0.08; random effect) (*GRADE: Low*).

Other blood components

None of the included RCTs or cohort studies in the trauma setting reported sufficient data relating to transfusion volumes of other blood components. The available evidence was therefore not further considered.

In patients with acute gastrointestinal bleeding, the RCT evidence (HALT-IT) suggested there was little to no difference on the volume of FFP transfused in patients who received TXA (mean 0.9 units) compared with those who did not receive TXA (mean 1.0 units) (MD – 0.10, 95%CI –0.21, 0.01; p = 0.07; random effect) (*GRADE: Low*). Similar results were also observed for the volume of PLT transfused (mean 0.2 units) (MD 0.00, 95%CI –0.04, 0.04; p = 1.00; random effect) (*GRADE: Low*).

(trauma)									
	Trane	xamic a	acid	c	Control		;	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% Cl	IV, Random, 95% CI
1.5.1 Trauma (RCTs)									
CRASH-2 2010 (Civilian) Subtotal (95% CI)	6.06	9.98	5067 5067	6.29	10.31	5160 5160	20.8% 20.8%	-0.02 [-0.06, 0.02] -0.02 [-0.06, 0.02]	
Heterogeneity: Not applicable Test for overall effect: Z = 1.15 (P = 0.25)									
1.5.2 Trauma (Coh)									
Cole 2015 (Civilian, ISS 15)	7	7.4	160	2	5	225	19.6%	0.82 [0.61, 1.03]	
Myers 2019 (Civilian)	4.43	5.57	189	2.53	3.35	189	19.7%	0.41 [0.21, 0.62]	
Morrison 2013 (MATTERS II) (with cryo)	22	13.2	258	20.1	16	168	19.8%	0.13 [-0.06, 0.33]	+=
Morrison 2013 (MATTERS II) (without cryo) Subtotal (95% CI)	8	6.2	148 755	6	0.8	758 1340	20.0% 79.2%	0.77 [0.59, 0.95] 0.53 [0.22, 0.85]	•
Heterogeneity: Tau ² = 0.09; Chi ² = 31.13, df = Test for overall effect: Z = 3.29 (P = 0.0010)	3 (P < 0	.00001)	; l² = 90	1%					

6500 100.0%

0.42 [0.03, 0.80]

-2

-1

0

Favours TXA Favours no TXA

Figure 4.50 Forest plot of comparison: TXA vs no TXA, outcome: RBC transfusion volume (trauma)

Figure 4.51 Forest plot of comparison: TXA vs no TXA, outcome: RBC transfusion volume (trauma)

5822

Total (95% CI)

Test for overall effect: Z = 2.13 (P = 0.03)

Heterogeneity: Tau² = 0.18; Chi² = 138.30, df = 4 (P < 0.00001); l² = 97%

Test for subgroup differences: $Chi^2 = 11.60$, df = 1 (P = 0.0007), l² = 91.4%

	Tranex	amic a	acid	Control			Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.7.1 Red blood cells									
Roberts 2020 (HALT-IT) Subtotal (95% CI)	2.8	2.4	4076 4076	2.9	2.7	4129 4129	100.0% 100.0%	-0.10 [-0.21, 0.01] -0.10 [-0.21, 0.01]	
Heterogeneity: Not applica Test for overall effect: Z =	ble 1.77 (P =	0.08)							
1.7.2 Frozen plasma									
Roberts 2020 (HALT-IT) Subtotal (95% CI)	0.9	2.4	4076 4076	1	2.6	4129 4129	100.0% 100.0%	-0.10 [-0.21, 0.01] -0.10 [-0.21, 0.01]	
Heterogeneity: Not applica	ble								
Test for overall effect: Z =	1.81 (P =	0.07)							
1.7.3 Platelets									
Roberts 2020 (HALT-IT) Subtotal (95% CI)	0.2	0.9	4076 4076	0.2	1	4129 4129	100.0% 100.0%	0.00 [-0.04, 0.04] 0.00 [-0.04, 0.04]	↓
Heterogeneity: Not applica	ble	1 00)							
Test for overall effect. $Z = 0$	0.00 (F =	1.00)							
									+ + + + + + + + + + + + + + + + + + + +
									-1 -0.5 0 0.5 1
									Favours TXA Favours placebo or no TXA

Table 4.73 Results for TXA versus no TXA: Patients with critical bleeding – Transfusion volume

Study ID	Sample size	Patient :	Setting (Comparison	Outcome	Results				
Study designª	(no. of trials) included in analysis	population	(Location)			TXA Mean ± SD (n)	No TXA Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b	
Trauma setting	7	·								
Cannon 2017 SR <i>Moderate</i> <i>quality</i>	N = 11 944 (1 RCT, 2 Coh) CRASH-2 2010 Cole 2015 Morrison 2013 CRYO+ Morrison 2013 CRYO- Morrison 2013 total	Patients with severe trauma at risk of death from haemorrhage* *patients requiring blood transfusion and/or with an injury score greater than 25	Civilian and military trauma	TXA vs no TXA	RBC transfusion volume, units	NR (5633) 6.06 ± 9.98 (5067) 7 ± 7.4 (160) 22 ± 13.2 (258) 8 ± 6.2 (148)	NR (6311) 6.29 ± 10.31 (5160) 2 ± 5 (225) 20.1 ± 16 (168) 6 ± 0.8 (758)	MD 2.14 (-0.36, 4.63) MD -0.23 (-0.62, 0.16) MD 5.00 (3.68, 6.32) MD 1.90 (-1.01, 4.81) MD 2.00 (1.00, 3.00) MD 1.99 (1.04, 2.94)	No significant difference p = 0.09 Substantial heterogeneity l² = 96 (p < 0.00001)	
Huebner 2017 SR <i>Critically low</i> <i>quality</i>	N = 300 (1 Coh) Valle 2014	Patients with trauma at risk of death from haemorrhage	Civilian trauma SC (US)	TXA vs no TXA	Total volume of RBC required in operating room, mL	2250	1500	NR	Favours placebo p = 0.002	
					Total volume fluid received in ED, mL	2675	2250	NR	Favours placebo p = 0.025	
					Total volume FFP in operating room, mL	1750	1125	NR	Favours placebo p = 0.009	
Ker 2015 SR High quality	N = 20 127 (1 RCT) CRASH-2 2010	Adult trauma patients with, or at risk of, significant bleeding, including patients with moderate to severe TBI	MC, 40 countries, Thailand	IV TXA vs standard care	Volume of blood transfused, mean <i>All trauma</i>	3.05 ± 7.7 (10 060)	3.22 ± 8.02 (10 067)	MD -0.17 (-0.39, 0.05)	<i>No significant difference</i> <i>p</i> = 0.13 Heterogeneity NA	
Myers 2019 Retrospective	N = 378	Patients presenting to a level 1 trauma	Level 1 trauma centre (NR)	TXA within 3 hours of	Transfusion of platelets, units	1.18 ± 2.17 (NR)	0.43 ± 1.43 (NR)	NR	Favours intervention P < 0.001	
Coh Serious risk of		centre		presentation vs No TXA	Transfusion of RBC, units	4.43 ± 5.57 (NR)	2.53 ± 3.35 (NR)	NR	Favours intervention p < 0.001	
UIdS					Transfusion of FFP, units	2.77 ± 5.14	1.44 ± 3.37 (NR)	NR	Favours intervention p < 0.001	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study designª	(no. of trials) included in analysis	population	(Location)			TXA Mean ± SD (n)	No TXA Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance p</i> -value Heterogeneity ^b	
Medical emerg	ency									
HALT-IT RCT <i>Low risk of</i>	N = 11937	Adult patients with severe or massive gastrointestinal	nts with MC (UK, Pakistan, nassive Nigeria, Egypt, stinal Malaysia, Georgia, Romania, Nepal, Sudan, Saudi	IV TXA vs placebo	Whole blood or RBC transfusion volume, units	2·8 ± 2·4 (4076)	2·9 ± 2·7 (4129)	MD -0.06 (0.05, -0.18)	NR	
bias		bleeding			FFP transfusion volume, units	0.9 ± 2.4 (4076)	1.0 ± 2.6 (4129)	MD -0·05 (-0·01, - 0·23)	NR	
	Arabia, Spain, Ireland, Albania, Papua New Guinea, ar Australia)	Arabia, Spain, Ireland, Albania, Papua New Guinea, and Australia)		Platelet transfusion volume, units	0·2 ± 0·9 (4076)	0·2 ± 1·0 (4129)	MD -0·02 (0·02, - 0·06)	NR		

Cl, confidence interval; Coh, cohort study; ED, emergency department; FFP, fresh frozen plasma; hrs, hours; IQR, interquartile range; IU, international units; IV, intravenous; MD, mean difference; M-H, Mantzel-Hentzel; NA, not applicable; NR, not reported; OR, odds ratio; RBC, red blood cells; RCT, randomised controlled trial; RR, relative risk; SC, single centre; SD, standard deviation; TXA, tranexamic acid; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic review studies with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4

4.9 Viscoelastic haemostatic assays (Question 8)

Question 8 - (interventional)

In patients *with* critical bleeding, does the use of viscoelastic haemostatic assays change patient outcomes?

4.9.1 Methods

This question examined the effects of viscoelastic haemostatic assays (TEG and ROTEM) compared to the use of an MHP and/or standard laboratory tests in guiding the transfusion of blood components in patients with critical bleeding (i.e. major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion) as outlined in Figure 4.52.

Studies were eligible for inclusion from any setting (including trauma, obstetrics, and perioperative) if, at the time of study inclusion, patients had major bleeding that was likely to result in the need for transfusion. Studies where bleeding status was not assessed at the time of inclusion were excluded (such as those that randomised patients prior to elective cardiac surgery). Studies in neonates (newborns up to 28 days) and studies in individuals with hereditary bleeding disorders were also not eligible for inclusion.

Viscoelastic haemostatic assays other than TEG or ROTEM (i.e. Sonoclot) were not eligible for inclusion.





Abbreviations: ARDS, acute respiratory distress syndrome; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; INR, international normalised ratio; MTP, massive transfusion protocol; PLT, platelets; RBC, red blood cells; ROTEM, rotational thromboelastometry; TE, thromboembolic event; TEG, thromboelastography

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months). b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other.

c. Newborns up to 28 days following birth.

The selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 2000, noting studies published prior to 2000 and identified within a systematic review were also eligible for inclusion. No restrictions were applied to study design (or size), meaning nonrandomised studies (with concurrent or noncurrent controls) and observational cohort studies were eligible for inclusion.

Assuming all relevant primary studies had been identified in the included systematic reviews³⁶ (171-181); the screening of primary studies was limited to studies published from January 2015. This was based off the latest literature search date of the most comprehensive identified systematic review (Wikkelsø 2017).

An updated literature search was conducted in August 2019³⁷ and again in September 2021³⁸ to identify any new studies meeting the eligibility criteria. In these updated searches the focus was the identification of systematic reviews (of RCTs or cohort studies). With the latest search date of the best available systematic review used as a starting point for screening for additional RCTs.

4.9.2 Summary of evidence

4.9.2.1 Systematic reviews

Twelve systematic reviews (171-181) were included that that assessed the effects of TEG or ROTEM to guide blood component therapy in patients with critical bleeding.

Four reviews were focused on patients with acute need for transfusion due to bleeding in any clinical setting (Roullet 2018, Wikkelsø 2017, Fahrendorrf 2017, Haas 2014), 2 focused on adult trauma patients (Da Luz 2014, Bugaev 2020), one was in bleeding management in patients with end-stage liver disease (Saner 2016), 4 on patients with coagulopathic bleeding in cardiac surgery (Li 2019, Serraino 2017, Deppe 2016, Corredor 2015), and one in management of major obstetric haemorrhage (Amgalan 2020).

The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.74.

A matrix illustrating the overlap of RCTs identified in each review is provided in Table 4.75.

Among the 24 RCTs identified by the included systematic reviews, there were 16 RCTs that were not included in the evidence evaluation because they were conducted in patients who did not have major bleeding at study inclusion (182-197). Twelve of these RCTs were in the cardiac setting (Karkouti 2016, Agarwal 2015, Nakayama 2015, Cui 2010, Girdauskas 2010, Ak 2009, Westbrook 2009, Rauter 2007, Kultufan Turan 2006, Avidan 2004, Royston 2001, Shore-Lesserson 1999), one in liver transplant (Wang 2010), one in hepatic surgery (De Pietri 2016), one in surgical excision of burn wounds (Schaden 2012), and one in scoliosis surgery (Cao 2016).

³⁶ 10 systematic reviews found (Li 2019, Roullet 2018, Fahrendorrf 2017, Serraino 2017, Wikkelsø 2017, Deppe 2016, Saner 2016, Corredor 2015, Da Luz 2014, Haas 2014).

³⁷ One systematic review found (Drumheller 2019) did not provide any additional data than that already included, therefore was not considered further (duplicate data).

³⁸ Two systematic reviews found (Amgalan 2020, Bugaev 2020) and included in the review.

A further 2 RCTs (122, 124) were not included because the studies were not appropriately designed to answer our research question. The RCT by Rahe-Meyer 2013 was confounded by different transfusion protocols as it assessed the administration of fibrinogen concentrate guided by TEG compared with standard care (FFP and platelet therapy) in bleeding patients undergoing elective thoracic or thoracoabdominal aortic replacement surgery involving CPB. The RCT by Collins 2017 evaluated the effects of early fibrinogen replacement (with fibrinogen concentrate) compared with placebo in women with ongoing postpartum haemorrhage (1000 to 1500 mLs). Participants in both groups received ROTEM guided care.

A matrix illustrating the overlap of nonrandomised cohort studies identified in each reviews is provided in Table 4.76.

Among the 38 nonrandomised cohort studies identified by the included systematic reviews, 24 (141, 198-220) were not included in the evidence evaluation because they did not include a comparator group (Rourke 2012, Johannsen 2013), involved patients who were not critically bleeding (Spiess 1995, Anderson 2006, Spalding 2007, Rahe-Meyer 2009a; Rahe-Meyer 2009b, Görlinger 2010, Noval-Padillo 2010, Trzebicki 2010, Görlinger 2011, Romlin 2011, Görlinger 2012, Hvas 2012, Wang 2012, Xu 2014, Leon-Justel 2015, Roullet 2015, Bedreli 2016, De Pietri 2016, St-Onge 2018, Kuiper 2019) or were confounded by inclusion of the intervention in both treatment groups (Mallaiah 2015, Collins 2017).

Review ID <i>Review</i> quality	Study design	Population	Intervention	Comparator	Outcomes									
Any clinical se	Any clinical setting													
Roullet 2018 (172) <i>Critically low</i>	Narrative review (Position paper)	Adult and paediatric patients with trauma, cardiac surgery and liver transplant, women with PPH	TEG or ROTEM guided algorithm	Standard care	Mortality Transfusion volume									
Fahrendorrf 2017 (173) <i>Low</i>	SR / MA of RCTs	Patients in acute need for blood transfusions due to bleeding	TEG or ROTEM guided algorithm	Standard care (clinician's discretion and/or conventional coagulation tests)	Mortality Transfusion volume									
Wikkelsø 2017 (175, 176) <i>High</i>	SR / MA of RCTs	Adults and children with bleeding	TEG or ROTEM guided algorithm	Standard care (clinical judgement, usual care and standard laboratory tests)	Mortality Transfusion volume									
Haas 2014 (181) <i>Critically low</i>	Narrative review	Trauma patients, cardiac and aortic surgical patients, liver transplantation	TEG or ROTEM guided algorithm	Standard care	Mortality Transfusion needs									

Table 4.74	Characteristics and quality of systematic review evidence: TEG or ROTEM versus
	usual care

Review ID Review quality	Study design	Population	Intervention	Comparator	Outcomes
Trauma					
Da Luz 2014 (180) <i>Moderate</i>	SR of RCTs and observational studies	Adult trauma patients	TEG or ROTEM guided algorithm	Standard care	Mortality Transfusion needs
Bugaev 2020 (221) <i>Moderate</i>	SR / MA of RCTs and observational studies	Adult trauma patients ^a	TEG or ROTEM guided algorithm	Standard care	Mortality Transfusion needs
Surgical (liver	transplant)				
Saner 2016 (177) <i>Critically low</i>	Narrative review (primary studies retrieved)	Bleeding management in patients with end-stage liver disease	TEG or ROTEM guided transfusion	Standard care	Mortality Transfusion needs
Surgical (card	iac)				
Li 2019 (171) <i>Moderate</i>	SR / MA of RCTs and observational studies	Patients with coagulopathic bleeding in cardiac surgery	TEG or ROTEM guided algorithm	Standard care	Mortality Morbidity Transfusion needs
Serraino 2017 (174) High	SR / MA of RCTs	Patients with coagulopathic bleeding in cardiac surgery	TEG or ROTEM guided algorithm	Standard care	Mortality Morbidity Transfusion needs
Deppe 2016 (178) <i>Moderate</i>	SR / MA of RCTs and observational studies	Patients undergoing cardiac surgery	TEG or ROTEM guided algorithm	Standard care	Mortality Major morbidity Transfusion needs
Corredor 2015 (178) <i>Moderate</i>	SR / MA of RCTs and observational studies	Patients undergoing cardiac surgery b	TEG or ROTEM guided algorithm	Standard care	Mortality Morbidity Transfusion needs
Obstetrics and	l maternity				
Amgalan 2020 (222) <i>Critically low</i>	SR of RCTs and observational studies	Pregnant women at risk of thrombosis or haemorrhage during pregnancy or peripartum	TEG or ROTEM guided transfusion	Standard care	Morbidity Transfusion needs

Abbreviations: MA, meta-analysis; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; SR, systematic review; TEG, thromboelastography

a. Bugaev 2020 examined the effects of TEG or ROTEM to guide blood component therapy in 3 populations: trauma, surgical and critically ill. Only the trauma population has been considered in this review.

b. Corredor 2015 also provided a narrative summary of 30 observational studies predicting post-operative bleeding loss. As this outcome was not considered in this review, these studies were not included here.

		Trauma	Surç	gical (cardi	ac)															Surgical (liver)	Sur (oth	gical 1er)		Obstetrics
	Study ID	Gonzalez 2016	Karkouti 2016 ª	Agarwal 2015ª	Nakayama 2015 ª	Rahe-Meyer 2013 ^b	Weber 2012	Kempfert 2011	Paniagua 2011	Cui 2010 ª	Girdauskas 2010 ^ª	Ak 2009ª	Westbrook 2009 ª	Rauter 2007 ^ª	Kultufan Turan 2006 ª	Avidan 2004 ª	NCT00772239	Nuttall 2001	Royston 2001 ^ª	Shore-Lesserson 1999	Wang 2010ª	Schaden 2012ª	Cao 2016ª	De Pietri 2016 ^ª	Collins 2017b ^b
	Amgalan 2020 ^e																								
	Bugaev 2020	✓																				Х			
	Li 2019 ^d		X				×	×	~		X	X	X	Х	Х	X		×	Х	X					
	Roullet 2018	~	X		X																No stu	dies	d		
0	Fahrendorrf 2017	✓					~	~	~		Х	Х	Х			Х		~	Х	X	Х	X	X	X	
3	Wikkelsø 2017				X		~	~	~	Х	Х	Х	Х	Х	Х	Х		~	X	X	х	Х			
èvie	Serraino 2017 °		Х		X		~		~	Х	Х	Х	Х			Х		~	Х	Х					
ď	Deppe 2016 ^c						~				Х	Х	X		Х	Х		~	X	X					
	Saner 2016 d																				Х			Х	
	Corredor 2015 °			X			~				Х	Х	Х			Х		~	Х	Х					
	Da Luz 2014 ^f																								
	Haas 2014					Х	~				Х											Х			

Table 4.75 Overlap table of RCTs identified by included systematic reviews: TEG or ROTEM versus usual care

✓ = study included in this review; X = study did not meet the inclusion criteria for this review; -- = study identified by the systematic review authors but not contribute any data

a. Population out of scope. Subjects not critically bleeding.

b. Study is confounded (all patients are assessed using a viscoelastic haemostatic assay) and does not provide usable data.

c. The systematic reviews by Li 2019, Serraino 2017, Deppe 2016 and Corredor 2015 were focused on patients in the cardiac setting.

d. The systematic review by Saner 2016 was focused on patients with end-stage liver disease.

e. The systematic review by Amgalan 2020 was focused on obstetrics patients.

f. The systematic review by Da Luz 2014 was focused on patients admitted with trauma.

Review ID	Tra	Trauma									Sur	gica	l (ca	rdia	c)									Surgical (liver transplant)							Obstetrics and maternity							
Study ID	Johansson 2013 ^a	Tapia 2013	Görlinger 2012a	Kashuk 2012	Rourke 2012 ^ª	Nienaber 2011	Schöchl 2011	Schöchl 2010 ^a	Unruh 2019	Guth 2019	Prat 2017	Nardi 2015	Kuiper 2019 b	St-Onge 2018 ^b	Sun 2014 ^b	Fassl 2013	Hanke 2012	Hvas 2012 ^b	Romlin 2011 b	Görlinger 2011 ^b	Rahe-Meyer 2009a ^b	Rahe-Meyer 2009b ^b	Spalding 2007 ^b	Anderson 2006 ^b	Spiess 1995 ^b	Bedreli 2016 ^b	Leon-Justel 2015 b	Roullet 2015 ^b	Görlinger 2012b ⁵	Wang 2012 ^b	Görlinger 2010 b	Noval-Padillo 2010 b	Trzebicki 2010 ^b	Snegovskikh 2018	Collins 2017b [€]	Mallaiah 2015 °	Barinov 2015	McNamara 2019
Amgalan 2020																																						~
Bugaev 2020							×		×	~	~	~																										
Li 2019 d													х	X		~	~			X				X	X													
Roullet 2018																	N	o sti	udies	fou	nd							X		X				~		Х		
Fahrendorrf 2017 •																																					1	
Wikkelsø 2017 •																																						
Serraino 2017 º																																						
Deppe 2016 d															х	~	~			x	x		x	x	x													
Saner 2016 ^f																										Х	х											
Corredor 2015 d																																						
Da Luz 2014 🛚		1		~			~	X																														
Haas 2014			~		X	~	~	X								~	~	X	X	X	X	X	X	X					X		X	Х	X	N	o stu	idies	four	nd

Table 4.76 Overlap table of nonrandomised cohort studies identified by included systematic reviews: TEG or ROTEM versus usual care

study included in this review; X = study did not meet the inclusion criteria for this review; -- study identified by the systematic review authors but not included (no usable data)

a. Study design out of scope, not a comparative study.

b. Population out of scope. Not all participants are critically bleeding.

c. Study is confounded (all patients are assessed using a viscoelastic haemostatic assay).

d. The systematic reviews by Li 2019, Deppe 2016 and Corredor 2015 were focused on patients in the cardiac setting.

e. The systematic reviews by Fahrendorff 2017, Serraino 2017 and Wikkelsø 2017 did not include nonrandomised studies.

f. The systematic review by Saner 2016 was focused on patients with end-stage liver disease.

g. The systematic review by Da Luz 2014 was focused on patients admitted with trauma.

4.9.2.2 Randomised controlled trials

The main characteristics and quality of the included RCTs and relevant outcomes assessed are summarised in Table 4.77.

There were 6 RCTs (223-228) identified by the included systematic reviews that were considered relevant to this review because they examined the effect of TEG or ROTEM in patients *with* critical bleeding. Two of the included studies used a TEG guided transfusion algorithm/haemorrhage protocol (Gonzalez 2016, Nuttall 2001) and the other 4 studies (Weber 2012, Kempfert 2011, Paniagua 2011, NCT00772239) used a ROTEM guided transfusion algorithm/haemorrhage protocol.

One additional RCT (Baksaas-Aasen 2020) was identified in the systematic review and handsearching process examined the effect of VHAs in adult trauma patients *with* critical bleeding (229, 230).

Baksaas-Aasen 2020 (iTACTIC) was a multicentre RCT conducted in Trauma centres located in Denmark, The Netherlands, Norway, Germany and the UK. The study focused on trauma-induced coagulopathy comparing outcomes in 396 patients in whom a local MHP had been initiated, with the transfusion algorithm/haemorrhage protocol guided by VHAs or conventional coagulation tests. The MHPs included empiric delivery of tranexamic acid, blood components delivered in a 1:1:1 ratio of RBC, plasma and platelet transfusions and limited infusion of crystalloid fluids.

Gonzalez 2016 was a single centre RCT conducted in the US that enrolled adults patients (aged over 18 years) with blunt or penetrating trauma sustained less than 6 hours before admission. Patients had to have an injury severity score greater than 15 and were likely to require transfusion of RBC within 6 hours from admission as indicated by clinical assessment. Patients were predominantly male (70.3%0 with a median (IQR) age of 30 (24 to 43). The number of patients with blunt / penetrating trauma was not reported.

Five RCTs (Weber 2012, Paniagua 2011, Kempfert 2011, NCT00772239, Nuttall 2001) were conducted at single centres and involved adult patients scheduled for cardiothoracic surgery, with various definitions for enrolment relating to diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery. Three studies were stopped early. Paniagua 2011 was terminated early due to slow recruitment and included 8 of 52 patients that did not meet the inclusion criteria. Weber 2012 was stopped early at an interim analysis due to clear benefits, and another study (NCT00772239) was stopped early due to futility (no data available).

The overall risk of bias for included RCTs was judged to be high (173, 174, 176). Most concerns were related to little or no allocation concealment or blinding of clinical personnel, which contributed to the high procedural bias favouring the intervention. Reporting bias was also considered high for blood loss, FFP transfusion and PLT transfusion due to incomplete reporting of outcome data, with no explanations given for missing data.

Study ID <i>Risk of bias</i>	Study design	Population	Intervention	Comparator	Outcomes
Trauma					
Baksaas- Aasen 2020 (iTACTIC) (229, 230) High	RCT, MC	Adult trauma patients with clinical signs of bleeding Activation of local MHP and if RBC transfusion had been initiated, randomised within 3 hours of injury and maximum of 1 hour after admission into the emergency department.	MHP guided by VHAs (n=201)	MHP guided by conventional coagulation test (n=195)	Mortality Morbidity Transfusion needs
Gonzalez 2016 (223) High	RCT, SC	Adult patients with blunt or penetrating trauma Injury sustained < 6 hours before admission, with ISS >15 and likely to require transfusion of RBC within 6 hours as indicated by clinical assessment	MTP guided by TEG (n=56)	MTP guided by conventional coagulation tests (aPTT, INR, fibrinogen level, D-dimer) (n=55)	Mortality Morbidity Transfusion needs
Cardiothora	cic surgery				
Weber 2012 (228) <i>High</i>	RCT, SC *study stopped at interim analysis	Adult patients scheduled for complex cardiothoracic surgery with CPB enrolled after heparin reversal if: (1) diffuse bleeding from capillary beds and/or (2) intraoperative or post-operative blood loss exceeding 250 mL/hour or 50 mL/10 min.	Peri- and post- operative management guided by ROTEM (n=50)	Algorithm based on standard laboratory tests (ACT, INR, aPTT, platelet count and fibrinogen level) (n=50)	Mortality, 6-mth Morbidity Transfusion needs
Paniagua 2011 (227) <i>High</i>	RCT, SC *study stopped due to slow enrolment	Adult patients undergoing cardiac surgery randomised if: (1) diffuse bleeding after protamine and/or (2) excessive bleeding after surgery (≥ 300 mL in the first hour; ≥ 250 mL in the second hour; ≥ 150 mL thereafter)	MTP guided by ROTEM (n=24)	Routine transfusion therapy based on standard laboratory coagulation tests (n=28)	Mortality Transfusion needs
Kempfert 2011 (224) <i>High</i>	RCT, SC	Adult patients with significant post-operative bleeding* following standard elective isolated or combined cardiac surgical procedures *(> 200 mL/hour)	MTP guided by ROTEM (n=52)	Transfusion protocol based on standard coagulation testing (n=52)	Transfusion needs
NCT007722 39 (225) High	RCT, SC *study stopped due to futility	Adult patients with abnormal bleeding after cardiac surgery or heart transplantation	Therapeutic algorithm guided by ROTEM (n=50)	Coagulation management based on standard laboratory tests (n=50)	No usable data
Nuttall 2001 (226) <i>High</i>	RCT, SC	Adult patients with abnormal microvascular bleeding after CPB	Transfusion algorithm guided by TEG (n=41)	Algorithm based on clinical judgement with or without laboratory tests (n=51)	Transfusion needs

Table 4.77 Characteristics and quality of RCT evidence: TEG or ROTEM versus usual care

Abbreviations: aPTT, activated partial thromboplastin time; CPB, cardiopulmonary bypass; INR, international normalised ratio; ISS, injury severity score; MC, multicentre; MHP, major haemorrhage protocol; MTP, massive transfusion protocol, PPH, postpartum haemorrhage; RCT, randomised controlled trial; SC, single centre; TEG, thromboelastography; VHA, viscoelastic haemostatic assay

4.9.2.3 Observational and cohort studies

The main characteristics and quality of the included nonrandomised studies and relevant outcomes assessed are summarised in Table 4.78.

There were 14 nonrandomised cohort studies identified by the included systematic reviews that examined the effects of TEG or ROTEM in guiding blood component therapy in patients *with* critical bleeding and were considered relevant to this review (137, 138, 220, 231-241). One additional nonrandomised cohort study (Wang 2017) (242) was identified in the literature search that examined the effects of TEG in patients who sustained traumatic liver and/or spleen injuries receiving emergent blood component therapy.

Six of the included studies used a TEG guided transfusion algorithm/haemorrhage protocol (Guth 2019, Unruh 2019, Wang 2017, Barinov 2015, Tapia 2013, Kashuk 2012), and 9 studies (McNamara 2019, Snegovskikh 2018, Prat 2017, Nardi 2015, Fassl 2013, Görlinger 2012, Hanke 2012, Nienaber 2011, Schöchl 2011) used a ROTEM guided transfusion algorithm/haemorrhage protocol.

Overall, 10 studies were conducted in the trauma setting (Guth 2019, Unruh 2019, Prat 2017, Wang 2017, Nardi 2015, Tapia 2013, Görlinger 2012, Kashuk 2012, Nienaber 2011, Schöchl 2011), 2 in the cardiac setting (Fassl 2013, Hanke 2012), and 3 in the obstetrics setting (McNamara 2019, Snegovskikh 2018, Barinov 2015).

In the trauma setting, 5 studies (Guth 2019, Wang 2017, Tapia 2013, Görlinger 2012, Kashuk 2012) were conducted at single centres and involved adult trauma patients (blunt and/or penetrating) with various definitions for injury severity and the timing or need for blood components (i.e. within 6 or 24 hours of admission). Five studies (Unruh 2019, Prat 2017, Nardi 2015, Nienaber 2011, Schöchl 2011) involved the collection of data from trauma registries (civilian and/or combat), with patients being selected based on injury severity (e.g. ISS \geq 16, base deficit \geq 2.0 mmol/L) or the need for blood components (e.g. receiving at least 3 units of RBC within the first 24 hours).

In the surgical setting, both studies were conducted at singles centres and included adult patients undergoing elective and urgent proximal aortic surgery with hypothermic circulatory arrest with major bleeding (Fassl 2013) or adult patients with acute type A aortic dissection and aortic valve replacement (Hanke 2012). The studies with conducted in Switzerland and Germany.

In the obstetric setting, all 3 studies evaluated the effect of a viscoelastic haemostatic assay guided algorithm for treatment of coagulopathy to improve outcomes for women with major obstetric haemorrhage. Two studies included women with severe PPH (defined as an estimated blood volume loss of \geq 1500 mLs) who had received care either before or after the introduction of a MHP that included a point-of-care viscoelastic assay. The studies were conducted at single centres in either the US (Snegovskikh 2018) or the UK (McNamara 2019) and reported data covering a 4- to 4.5-year period.

One study (Barinov 2015) was conducted in Russia and prospectively included women with PPH managed using a combined strategy involving TEG assessment of coagulation, early surgical haemostasis (estimated blood volume loss of ≥ 1000 mLs) and mechanical compression of the uterine wall combined with uterine cavity draining, via intrauterine balloon tamponade. The comparator group received uterine massage, manual examination of the uterus, and transfusion of FFP, RBC, PLT and protease inhibitors, with

late surgical haemostasis (blood loss volume \geq 2000 mL). In the cases of severe obstetric bleeding, autologous red blood cell reinfusion was carried out (cell salvage).

Many of the included observational cohort studies were at serious risk of bias. This is because they were often conducted before and after the introduction of the intervention into clinical practice, introducing concerns with procedural bias that would favour the intervention. The use of historical controls introduced issues with changes in clinical practices that occur over time. The studies also had issues with incomplete report of outcome data, short follow-up and small sample size.

Study ID	Study design	Population	Intervention	Comparator	Outcomes			
RISK OF DIAS								
Guth 2019	Before/after	Adult trauma	TEG guided	CCT-guided	Mortality			
(239) Serious	analysis, SC *historical controls	patients (ISS > 8) *who received at least one blood component (RBC, FFP, PLT) or coagulation factor concentrates (fibrinogen or PCC) during the first 24 hours after admission N=380	management including FFP, PCC and PLT * Patients admitted from 1 January 2005 to 31 December 2008	transfusion protocol * Patients admitted from 1 January 2012 to 31 December 2015	Transfusion volume			
Unruh 2019 (241) <i>Serious</i>	Retrospective analysis, trauma registry ^a *historical controls	Adult trauma patients who underwent MTP activation N=67	MTP guided by TEG * patients who underwent MTP activation from 1 July 2015 to 30 June 2016	Non-TEG guided MTP *patients who underwent MTP activation from 1 January 2014 to 31 December 2014	Mortality Blood component utilisation			
Prat 2017 (237) <i>Serious</i>	Retrospective analysis, trauma registry ^b *historical controls	Civilian and combat trauma patients N=219	ROTEM guided transfusion * after ROTEM deployment to Bagram Airfield.	Non-ROTEM guided transfusion * before ROTEM deployment to Bagram Airfield	Mortality Transfusion volume Transfusion ratios			
Wang 2017 (242) <i>Critical</i>	Retrospective cohort, SC	Adult trauma patients with sustained liver and/or spleen injuries receiving emergent BCT N=166	Blood component therapy guided by TEG	Non-TEG guided blood component therapy	Mortality Transfusion volume			
Nardi 2015 (235) <i>Serious</i>	Before/after analysis, MC ^c *historical controls	Severely injured trauma patients (ISS > 15) *receiving at least 3 units of RBC within the first 24 hours) N=226	ECS protocol guided by ROTEM *post-ECS adoption period (1 January 2013 to 31 December 2013)	Non-ROTEM guided transfusion protocol *pre-ECS adoption period (1 January 2011 to 31 December 2011)	Mortality Transfusion volume			
Tapia 2013 (234) <i>Moderate</i>	Before/after cohort, SC	Adult trauma patients (blunt or penetrating) 'receiving ≥ 6 units of RBC in the first 24 hours N=289	MTP guided by TEG	Non-TEG guided MTP	Mortality Transfusion volume			
Görlinger 2012 (220) <i>High</i>	Retrospective analysis, 3 x SC ^d *historical controls	Patients admitted in different perioperative settings (trauma, visceral and transplant, cardiovascular	ROTEM guided management with FC and PCC	Non-ROTEM guided transfusion protocol	Transfusion volume			

Table 4.78 Characteristics and quality of Observational and cohort studies evidence: TEG or ROTEM versus usual care

Study ID <i>Risk of bias</i>	Study design	Population	Intervention	Comparator	Outcomes
		and general and surgical intensive care) N=5590			
Kashuk 2012 (232) <i>Moderate</i>	Before/after analysis, SC	Adult trauma patients *receiving ≥ 6 units of RBC in the first 6 hours N=64	MTP guided by TEG	Non-TEG guided MTP	Mortality Transfusion volume
Nienaber 2011 (137) <i>High</i>	Retrospective analysis, 2 trauma registries ^e	Adult trauma patients with severe blunt trauma (ISS ≥ 16, base excess ≤ -2.0 mmol/L) N=36	ROTEM guided management including factor concentrates	Non-ROTEM guided transfusion protocol including FFP (no factor concentrates)	Mortality Morbidity Transfusion volume
Schöchl 2011 (138) <i>High</i>	Retrospective analysis, 2 trauma registries ^f	Adult trauma patients with severe trauma (ISS \geq 16, base deficit \geq 2.0 mmol/L) N=36	ROTEM guided management including FC and PCC	Non-ROTEM guided transfusion protocol including FFP (no FC and PCC)	Mortality Morbidity Transfusion volume
Surgical settin	ng				
Fassl 2013 (233) High	Retrospective case-control, SC	Adult patients undergoing elective and urgent proximal aortic surgery with hypothermic circulatory arrest with major bleeding ⁹ N=194	Haemostatic management guided by ROTEM including FFP, PLT, FC, PCC and rFVIIa	Conventional haemostatic management *matched controls	Mortality Morbidity Transfusion volume
Hanke 2012 (231) <i>High</i>	Case-control (pilot), SC	Adult patients with acute type A aortic dissection and aortic valve replacement N=10	Haemostatic management guided by ROTEM	Conventional haemostatic management *matched controls	Mortality Transfusion volume
Obstetrics and	d maternity				
McNamara 2019 (240) <i>Serious</i>	Retrospective analysis, SC 'matched historical controls	Women with major obstetric haemorrhage N=255 "estimated >1500 mL	MHP guided by ROTEM	Non-ROTEM guided MHP	Mortality Morbidity Transfusion volume
Snegovskikh 2018 (238) <i>High</i>	Retrospective analysis, SC *historical controls	Women with severe PPH* N=86 *estimated >1500 mL	MHP guided by ROTEM	Non-ROTEM guided MHP	Morbidity
Barinov 2015 (236) <i>High</i>	Open, prospective controlled trial, SC	Women with PPH (gestational age 28 to 42 weeks) N=119 *estimated >1000 mL	Combined strategy ^h of haemorrhage management including TEG	Conventional management	Transfusion volume

Abbreviations: BCT, blood component therapy; CCT, conventional coagulation tests; ECS, early coagulation support; FC, fibrinogen concentrate; FFP, fresh frozen plasma; ICU, intensive care unit; ISS, injury severity score; MTP, massive transfusion protocol; RBC, red blood cells; PCC, prothrombin complex concentrate; PLT, platelets; RBC, red blood cell; ROTEM, rotational thromboelastometry; rFVIIa, recombinant activated factor VII; SC, single centre; TEG, thromboelastography

a. Trauma registry of the American College of Surgeons verified Level I trauma centre.

b. Data collected from the Department of Defence Trauma Registry.

c. Data collected from 2 hospitals: S Camillo Hospital in Rome and Bufalini Hospital in Cesena, Italy.

d. Each hospital reported separately: (a) Trauma Centre Salzburg (b). Intensive care, University Hospital Essen (c) General/Surgical Critical Care, Medical University Innsbruck. Data from (a) and (b)reported here.

e. Comparison between Innsbruck Trauma databank (intervention) and Trauma-Registry-DGU, Germany (control).

f. Comparison between Salzburg Trauma Centre and the Trauma-Registry-DGU, Germany (control).
- g. Major bleeding defined as need for thoracic re-exploration or drainage volumes exceeding 1000 mL in the first 24 hours.
- h. Combined strategy included *early* surgical haemostasis if blood volume loss exceeded 1000 mL, mechanical pressure and intrauterine balloon tamponade; the comparator included uterine massage, transfusion of blood components, and *late* surgical haemostasis (blood volume loss exceeded 2000 mL).

4.9.3 Results

4.9.3.1 Mortality

A summary of the evidence relating to the outcome of mortality in patients with critical bleeding in whom TEG or ROTEM were used as part of a major haemorrhage protocol is presented in Table 4.79.

All identified systematic reviews suggested that the use of viscoelastic haemostatic assays to guide blood component, product and antifibrinolytic therapy provides no significant survival benefit in patients with critical bleeding, regardless of clinical setting.

A meta-analysis of data including evidence from both RCTs and cohort studies (see Figure 4.53) showed the mortality rate (latest timepoint) among patients who are critically bleeding to be lower when a TEG or ROTEM-guided transfusion protocol was used compared with haemostatic management guided by an MHP, standard laboratory tests or clinical judgement with or without laboratory tests (14.8% vs 17.9%; RR 0.75; 95% CI 0.64, 0.88; p = 0.004; random effect, I² = 0%).

Data from the included RCTs suggested the mortality rate to be lower in the TEG or ROTEM groups (19.8%) when compared with an MHP or transfusion algorithm/haemorrhage protocol that was not guided by a VHA (28.1%) (RR 0.61; 95% CI 0.37, 1.02; p = 0.06; random effect, I² = 44%). The difference was considered clinically important, despite not reaching statistical significance. *(GRADE: very low)*.

Data from the included cohort studies, suggested that TEG or ROTEM guided transfusion protocols were associated with reduced mortality compared with haemostatic management guided by an MHP, algorithm or standard laboratory tests (RR 0.75; 95% CI 0.62, 0.94; p = 0.004; $l^2 = 0$ %) (GRADE: very low).

In trauma patients (see Figure 4.54), a total of 952 patients received a TEG or ROTEMguided transfusion protocol, compared with 1474 patients who received a transfusion protocol guided by standard laboratory tests or clinical judgement with or without laboratory test. Among patients enrolled in 2 RCTs, the mortality rate (latest timepoint) was lower when a TEG or ROTEM-guided MHP was used (23.7%) than when the MHP was guided by standard laboratory tests (30.1%). The difference was not statistically significant but was considered clinically important (RR 0.75; 95% CI 0.48, 1.17; p = 0.20; l²=44%) (*GRADE: very low*).

In trauma patients, evidence in the cohort studies suggests that TEG or ROTEM-guided transfusion protocols are associated with a significantly lower mortality rate than transfusion protocols that are guided standard laboratory tests (19.3% vs 17.3%; RR 0.75; 95% Cl 0.62, 0.92; p = 0.004; $l^2 = 0$ %) (*GRADE: very low*).

In patients with diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery (see Figure 4.54), a ROTEM-guided transfusion protocol had a mortality rate of 6.6% (5/76), which was lower than the mortality rate of 20.6% (14/68) observed among those whose management was not guided by ROTEM (RR 0.33; 95% CI 0.12, 0.91; p = 0.03; $l^2 = 0$ %) (*GRADE: very low*).

No deaths were observed in the observational studies that assessed the effects of a TEG or ROTEM-guided transfusion protocol among women with severe obstetric haemorrhage.

The sample size of included studies were small and not optimal for detecting the outcome of interest.

Figure 4.53 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: Mortality, latest timepoint

	TEG or R	OTEM	Conti	ol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl
1.1.2 RCTs: coagulopathy or severe	bleeding at	inclusio	n				
Weber 2012 (Cardiac)	2	50	10	50	1.2%	0.20 [0.05, 0.87]	
Paniagua 2011 (Cardiac)	3	26	4	18	1.4%	0.52 [0.13, 2.05]	
Gonzalez 2016 (Trauma)	11	56	20	55	6.4%	0.54 [0.29, 1.02]	
Baksaas-Aasen 2020 (Trauma)	50	201	55	194	23.8%	0.88 [0.63, 1.22]	
Subtotal (95% CI)		333		317	32.7%	0.61 [0.37, 1.02]	\bullet
Total events	66		89				
Heterogeneity: Tau ² = 0.11; Chi ² = 5.38	, df = 3 (P =	0.15); l²	= 44%				
Test for overall effect: Z = 1.88 (P = 0.0	6)						
1.1.3 NRSIs: patients with critical ble	eding						
Snegovskikh 2018 (Coh, obstetrics)	0	28	0	58		Not estimable	
Barinov 2015 (Coh, PPH)	0	90	0	29		Not estimable	
McNamara 2019 (Coh, obstetrics) (1)	0	203	0	52		Not estimable	
Prat 2017 (Coh, trauma)	4	85	7	134	1.8%	0.90 [0.27, 2.99]	
Schochl 2011 (Coh, trauma)	6	80	60	601	3.9%	0.75 [0.34, 1.68]	
Wang 2017 (Coh, trauma)	12	86	19	80	6.0%	0.59 [0.31, 1.13]	
Nardi 2015 (Coh, trauma)	13	96	26	130	6.9%	0.68 [0.37, 1.25]	
Kashuk 2012 (Coh, trauma)	10	34	20	34	7.3%	0.50 [0.28, 0.90]	
Unruh 2019 (Coh, trauma)	15	47	11	20	7.7%	0.58 [0.33, 1.03]	
Guth 2019 (Coh, trauma)	33	102	34	102	16.6%	0.97 [0.66, 1.44]	
Tapia 2013 (Coh, trauma)	41	165	35	124	17.2%	0.88 [0.60, 1.30]	
Subtotal (95% CI)		1016		1364	67.3%	0.75 [0.62, 0.92]	\bullet
Total events	134		212				
Heterogeneity: Tau ² = 0.00; Chi ² = 5.62	, df = 7 (P =	0.59); l²	= 0%				
Test for overall effect: Z = 2.85 (P = 0.0	04)						
Total (95% CI)		1349		1681	100.0%	0.75 [0.64, 0.88]	•
Total events	200		301				
Heterogeneity: Tau ² = 0.00; Chi ² = 10.9	4, df = 11 (P	= 0.45);	l ² = 0%				
Test for overall effect: Z = 3.54 (P = 0.0	004)						UUD U.2 I D 20 Eavours TEG or ROTEM Eavours Control
Test for subgroup differences: Chi ² = 0.	54, df = 1 (P	= 0.46),	l ² = 0%				
Footnotes							

(1) women with major obstetric haemorrhage (estimated blood loss > 1500 mL) and coagulopathy

Figure 4.54 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: Mortality, by setting

	TEG or R	OTEM	Contr	ol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% CI
.2.1 Trauma (RCTs)							
Gonzalez 2016 (Trauma)	11	56	20	55	6.4%	0.54 [0.29, 1.02]	
Baksaas-Aasen 2020 (Trauma) Subtotal (95% CI)	50	201 257	55	194 249	23.8% 30.1%	0.88 [0.63, 1.22] 0.75 [0.48 , 1.17]	•
otal events	61		75				
leterogeneity: Tau ² = 0.05; Chi ² = 1.77, Test for overall effect: Z = 1.28 (P = 0.20	, df = 1 (P = 0))	0.18); l²	= 44%				
.2.2 Trauma (Coh)							
Prat 2017 (Coh, trauma)	4	85	7	134	1.8%	0.90 [0.27, 2.99]	
Schochl 2011 (Coh, trauma)	6	80	60	601	3.9%	0.75 [0.34, 1.68]	
Vang 2017 (Coh, trauma)	12	86	19	80	6.0%	0.59 [0.31, 1.13]	
lardi 2015 (Coh, trauma)	13	96	26	130	6.9%	0.68 [0.37, 1.25]	
ashuk 2012 (Coh, trauma)	10	34	20	34	7.3%	0.50 [0.28, 0.90]	
Jnruh 2019 (Coh, trauma)	15	47	11	20	7.7%	0.58 [0.33, 1.03]	
Guth 2019 (Coh, trauma)	33	102	34	102	16.6%	0.97 [0.66, 1.44]	_
lapia 2013 (Coh, trauma)	41	165	35	124	17.2%	0.88 [0.60, 1.30]	
Subtotal (95% CI)		695		1225	67.3%	0.75 [0.62, 0.92]	\bullet
Fotal events	134		212				
Heterogeneity: Tau ² = 0.00; Chi ² = 5.62,	, df = 7 (P =	0.59); l²	= 0%				
Test for overall effect: Z = 2.85 (P = 0.00	04)						
.2.3 Cardiac							
Neber 2012 (Cardiac)	2	50	10	50	1.2%	0.20 [0.05, 0.87]	·
Paniagua 2011 (Cardiac)	3	26	4	18	1.4%	0.52 [0.13, 2.05]	
Subtotal (95% CI)		76		68	2.6%	0.33 [0.12, 0.91]	
Total events	5		14				
Heterogeneity: Tau ² = 0.00; Chi ² = 0.90, Test for overall effect: Z = 2.15 (P = 0.03)	, df = 1 (P = 0 3)	0.34); l²	= 0%				
1.2.4 Obstetrics and maternity							
Snegovskikh 2018 (Coh. obstatrice)	٥	28	٥	58		Not estimable	
Sarinov 2015 (Coh PPH)	0	20 Q()	0	20		Not estimable	
AcNamara 2019 (Coh. obstetrics) (1)	0	203	0	23 52		Not estimable	
Subtotal (95% CI)	0	321	0	139		Not estimable	
Total events	0		٥				
Heterogeneity: Not applicable	0		0				
Test for overall effect. Not applicable							
rest for overall eneor. Not appliedble							
Fotal (95% CI)		1349		1681	100.0%	0.75 [0.64, 0.88]	◆
Total events	200		301				
Heterogeneity: Tau ² = 0.00; Chi ² = 10.94	4, df = 11 (P	= 0.45);	l ² = 0%				
Test for overall effect: Z = 3.54 (P = 0.00)04)						Favours TEG or ROTEM Favours Control
Test for subgroup differences: Chi ² = 2.4	47, df = 2 (P	= 0.29),	l ² = 18.9°	%			
-							

Footnotes

(1) women with major obstetric haemorrhage (estimated blood loss > 1500 mL) and coagulopathy

Table 4.79	Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding – M	ortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Various settings		·							
Fahrendorff 2017 SR High risk of bias	N = 579 (6 RCTs) Ak 2009 Wang 2010 Girdauskas 2010 Weber 2012 Gonzalez 2016 De Pietri 2016	Patients with an acute need for blood components due to bleeding (includes trauma, liver transplant and cardiac surgery)	SC, Various	TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional coagulation tests	Mortality, all-cause	30/291 (10.3) 3/114 (2.6) 2/14 (14.3) 4/27 (14.8) 2/50 (4) 11/56 (19.6) 8/30 (26.7)	47/288 (16.3) 2/110 (1.8) 3/14 (21.4) 5/29 (17.2) 10/50 (20) 20/55 (36.3) 7/30 (23.3)	OR 0.60 (0.34, 1.07) OR 0.60 (0.34, 1.07) OR 0.61 (0.09, 4.37) OR 0.83 (0.20, 3.50) OR 0.17 (0.03, 0.81) OR 0.43 (0.18, 1.01) OR 1.19 (0.3.85)	No significant difference p = 0.08 Mild heterogeneity I ² = 11% (p = 0.35)
Trauma setting									
Bugaev 2020 SR High risk of bias	N = 1488 (1 RCT, 5 Coh) Schochl 2011 Nardi 2015 Gonzalez 2016 Prat 2017 Unruh 2019 Guth 2019	Severely injured trauma patients	Trauma (Germany, US, NR)	ROTEM or TEG guided algorithm vs no ROTEM or TEG	Mortality, all-cause	82/466 (17.6) 6/80 (7.5) 13/96 (13.5) 11/56 (19.6) 4/85 (4.7) 15/47 (32.0) 33/102 (32.4)	158/1042 (15.2) 60/601 (10) 26/130 (20) 20/55 (36.4) 7/134 (5.2) 11/20 (55) 34/102 (33.3)	RR 0.75 (0.59, 0.95) RR 0.75 (0.34, 1.68) RR 0.68 (0.37, 1.25) RR 0.54 (0.29, 1.02) RR 0.90 (0.27, 299) RR 0.58 (0.33, 103) RR 0.87 (0.66, 1.44)	Favours TEG/ROTEM p = 0.02 No significant heterogeneity $l^2 = 0 (p = 0.60)$
Baksaas-Aasen 2020 RCT High risk of bias	N = 396	Adult trauma patients with clinical signs of bleeding and activation of local MHP.	MC, Various (Denmark, Netherlands, Norway, Germany, UK)	TEG or ROTEM guided algorithm vs conventional coagulation test	Mortality, all-cause 6 hours 24 hours 28 days 90 days	22/201 (11) 29/201 (14) 50/201 (25) 53/179 (29)	22/195 (11) 33/195 (17) 55/194 (28) 56/177 (31)	OR 0.97 (0.52,1.80) OR 0.83 (0.48, 1.42) OR 0.84 (0.54, 1.31) OR 0.91 (0.58, 1.42)	No significant difference p = 0.915 p = 0.495 p = 0.435 p = 0.678
Roullet 2018 RCT High risk of bias	N = NR (1 RCT) Gonzalez 2016	Patients with severe trauma who are likely to require transfusion therapy	SC (US)	TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional coagulation tests	Mortality, 28 days < 6 hours 6-24 hours	11/56 (19.6) 4/56 (7.1) 7/56 (12.5)	20/55 (35.7) 12/55 (21.8) 8/55 (14.5)	RR 0.54 (0.29, 1.02) ^c RR 0.33 (0.11, 0.95) RR 0.86 (0.33, 2.21)	Favours TEG p = 0.049
Wang 2017 Coh <i>High risk of bias</i>	N = 166 (1 Coh)	Patients with traumatic liver and/or spleen injury	SC (US)	TEG guided algorithm vs clinical judgement or usual treatment	Mortality, in-hospital (total) N = 166	12/86 (14)	19/80 (15)	RR 0.59 (0.31, 1.13) ^c	No association p = 0.11

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Da Luz 2014 Coh Serious risk of bias	N = 289 (1 Coh) Tapia 2013	Adult patients with severe trauma	SC (NR)	TEG or ROTEM guided algorithm vs standard of care	Mortality, 30 days patients receiving ≥ 6 U RBC Blunt Penetrating	41/165 (25) 14/47 (30) 27/118 (23)	35/124 (28) 13/52 (25) 22/72 (31)	RR 0.88 (0.60, 1.30) ^c NR NR	No association p = 0.52 No association observed in multivariate analysis (blunt or penetrating trauma)
						Subgroup: patients RBC	ubgroup: patients with penetrating trauma receiving ≥ 10 U RBC		
					Penetrating	22/66	20/37	NR	<i>p</i> = 0.04
	N = 68 (1 Coh) Kashuk 2012				Mortality •not adjusted for confounders	10/34 (29)	20/34 (59)	RR 0.50 (0.28, 0.90) ^c	Favours <i>TEG</i> p = 0.02
	N = 681 (1 Coh) Schöchl 2011			FC & PCC-guided vs FFP d	Mortality	6/80 (7.5)	60/601 (10)	RR 0.75 (0.34, 1.68) ^c	No association p = 0.69
	N = 131 (1 Coh) Schöchl 2010	-		FP, PLT, PCC- guided	Mortality vs TRISS predicted vs RISC predicted	NR (24.4)	NR (33.7) NR (28.7)	NR NR	No association p = 0.032 p > 0.05
					vs TRISS predicted	Subgroup: excludir	ng patients with TBI		Favours ROTEM
					vs RISC predicted	NR (14)	NR (27.8) NR (24.3)	NR NR	<i>p</i> = NR <i>p</i> = NR
Haas 2014 Coh	N = 681 (1 Coh) Schöchl 2011	Adult patients with severe	SC (Austria,	ROTEM guided algorithm vs	Mortality	6/80 (7.5)	60/601 (10)	RR 0.75 (0.34, 1.68) ^c	No association p = NR
Serious risk of bias	N = 36 (1 Coh) Nienaber 2011	trauma	Germany)	standard of care ^{d, e}		5/18 (13.9)	2/18 (11.1)	RR 2.50 (0.56, 11.25) °	No association p = 0.500
	N = 131 (1 Coh) Schöchl 2010				vs TRISS predicted	NR (24.4)	NR (33.7)	NR	Favours ROTEM p = 0.032
						Subgroup: excludir	ng 17 patients with TBI		Favours ROTEM
						NR (14)	NR (27.8)	NR	<i>p</i> = 0.0018
Surgical setting									
Wikkelsø 2016 SR High risk of bias	N = 717 (8 RCTs)		SC		Mortality, latest follow-up * *7 out of 8 RCTs were at hospital discharge	14/364 (3.9)	26/353 (7.4)	M-H Random effects RR 0.57 (0.30, 1.07) Adj. 0.59 (0.23, 1.54) '	No significant difference p = 0.08 No heterogeneity $l^2 = 0 (p = 0.54)$

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
	Ak 2009 Girdauskas 2010 Royston 2001 Shore-Lesserson 1999 Nakayama 2015 Paniagua 2011	Adults with diffuse or excessive bleeding after cardiac surgery or orthotopic	(Australia, Austria, China, France, Germany, Japan, Spain, Taiwan	TEG or ROTEM guided transfusion vs clinical judgement or usual treatment with or without predefined	TEG (4 RCTs) ROTEM (4 RCTs)	14/364 (3.9) 5/211 (2.4) 9/153 (5.9)	26/353 (7.4) 7/206 (3.4) 19/147 (12.9)	M-H Fixed effect RR 0.52 (0.28, 0.95) Adj. 0.51 (0.21, 1.26) [†] RR 0.72 (0.25, 2.07) RR 0.44 (0.21, 0.93)	Favours TEG/ROTEM p = 0.033 No heterogeneity $l^2 = 0$ ($p = 0.54$)
	Wang 2010	liver transplant	Turkey, UK, US)	algorithm to guide		Subgroup analyses	s: By comparator		
	Weber 2012			SLTS	vs clinical judgement or usual treatment (4 RCTs)	7/224 (3.1)	9/221 (4.1)	RR 0.81 (0.32, 2.01)	No significant difference p = 0.65 No heterogeneity $l^2 = 0$ (p = 0.53)
					vs SLT-guided transfusion (4 RCTs)	7/140 (5)	9/132 (6.8)	RR 0.36 (0.16, 0.84)	Favours TEG or ROTEM p = 0.018 No heterogeneity $l^2 = 0$ ($p = 0.49$)
Li 2019 SR High risk of bias	N = NR (5 RCTs, 3 Coh) Ak 2009 Shore-Lesserson 1999 Girdauskas 2010 Paniagua 2011 Weber 2012 Görlinger 2011 Kuiper 2019 St-Onge 2018	Adult patients undergoing cardiac surgery with CPB	(Austria, Canada, Germany, Spain, The Netherlands Turkey, UK, US)	TEG or ROTEM guided transfusion vs standard of care	Mortality, latest follow-up) RCTs only	132/2680 (5) 12/270 (4.4)	124/2293 (5.4) 23/259 (8.9)	RR 0.83 (0.53, 1.30) RR 0.5 (0.26, 0.96)	No significant difference p = 0.4 Moderate heterogeneity $I^2 = 25 (p = NR)$
Serraino 2017 SR High risk of bias	N = 689 (7 RCTs) Ak 2009 Girdauskas 2010 Nakayama 2015 Paniagua 2011 Royston 2001 Shore-Lesserson 1999 Weber 2012	Adult and paediatric patients undergoing cardiac surgery with bleeding	SC, cardiac (NR)	TEG or ROTEM guided transfusion (with or without other point-of-care platelet function tests) vs standard of care	Mortality, latest follow-up	12/350 (3.4)	23/339 (6.8)	RR 0.55 (0.28, 1.10)	No significant difference p = 0.09 No significant heterogeneity $l^2 = 1 (p = 0.40)$
Deppe 2016 SR High risk of bias	N = 5899 (6 RCTs, 5 Coh)	Patients with excessive bleeding after cardiac surgery	SC, cardiac (NR)	TEG or ROTEM guided transfusion vs standard of care	Mortality, all-cause	163/NR (5.4)	156/NR (5.7)	OR 0.92 (0.74, 1.16)	No significant difference p = 0.5193 Mild heterogeneity $l^2 = 14 (p = 0.4520)$

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
Corredor 2015 SR High risk of bias	N = 749 (6 RCTs) Shore-Lesserson 1999 Ak 2009 Girdauskas 2010 Royston 2001 Weber 2012 Agarwal 2015	Patients undergoing cardiac surgery	Cardiac surgery (NR)	TEG or ROTEM guided transfusion (with or without other point-of-care platelet function tests) vs standard of care	Mortality, latest follow-up	NR	NR	RR 0.66 (0.31, 1.39)	No significant difference p = 0.27 Heterogeneity NR		
Haas 2014 RCT High risk of bias	N = 100 (1 RCT) Weber 2012	Adult patients undergoing cardiac or aortic surgery	SC (NR)	ROTEM guided algorithm vs standard of care	Mortality, 6 months	NR/NR (4)	NR/NR (20)	NR	Favours ROTEM p = 0.013		
Obstetrics and ma	aternity										
Amgalan 2020 Coh High risk of bias	N = 100 (1 Coh) McNamara 2019	Women with MOH associated with coagulopathy (estimated blood loss >1500mL)	SC (NR)	ROTEM guided algorithm vs standard of care	Mortality, latest follow-up	0/203	0/203	Not estimable	No significant difference		

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: CI, confidence interval; Coh, cohort study; CPB, cardiopulmonary bypass; FC, fibrinogen concentrate; FFP, fresh frozen plasma; M-H, Mantzel-Hentzel; MHP, major haemorrhage protocol; MOH, major obstetric haemorrhage; NR, not reported; OR, odds ratio; PCC, prothrombin complex concentrate; PLT, platelets; RBC, red blood cells; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; RR, relative risk; SC, single centre; SLTs, standard laboratory tests; TEG, thromboelastography; UK, United Kingdom; US, United States; vs, versus

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25; (ii) mild heterogeneity if I² <25; moderate heterogeneity if I² >50.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

d. Schöchl 2011 compared ROTEM guided administration of FC and PCC vs standard care guided transfusion (receiving >2 units FFP [no FC or PCC]). Patients in intervention group received median 6 g FC (range 0—15) and 1200 IU PCC (range 0—6600) and those in the comparator group received median 6 Units FFP (range 2—51)

e. Nienaber 2011 compared ROTEM guided administration of FC and PCC with standard care guided transfusion of 1:1 FFP:RBC ratio.

f. Trial sequential analysis showed only 54 of required information size (717/1325) had been reached. Not statistically significant with control event proportion of 7.4.

4.9.3.2 Morbidity

A summary of the evidence relating to the incidence of thromboembolic events in patients with critical bleeding in whom TEG or ROTEM were used as part of a major haemorrhage protocol is presented in Table 4.80.

A summary of the evidence relating to major morbidities (multiple organ failure, renal failure, or need for postpartum hysterectomy) in patients with critical bleeding in whom TEG or ROTEM were used as part of a major haemorrhage protocol is presented in Table 4.81.

Thromboembolic events

In a meta-analysis of data from the included RCTs (see Figure 4.55), the rate of thromboembolic events in patients with critical bleeding who received a TEG or ROTEM-guided transfusion protocol was 7.2% (24/333) compared with 9.4% (30/318) among those whose received an MHP or transfusion protocol guided by standard laboratory tests. The difference between treatment groups was not significant (RR 0.83; 95% CI 0.41, 1.66; p = 0.60, I²= 26%) (*GRADE: very low*).

Among trauma patients, the rate of thromboembolic events reported in those who received a TEG or ROTEM-guided transfusion protocol was 9.3% (24/257), which was comparable with those whose MHP was guided by standard laboratory tests (11.2%; 28/250). The difference was not statistically significant (RR 0.90; 95% CI 0.42, 1.95; p = 0.80, $I^2 = 46\%$) (*GRADE: very low*).

In patients with diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery, the rate of thromboembolic events among those who received a ROTEM-guided transfusion protocol was 0% (0/76) compared with 2.9% (2/68) in the comparator group. The difference was not statistically significant (RR 0.20; 95% CI 0.01, 4.06; p = 0.29) (*GRADE: very low*). Only one study contributed data.

Multiple organ failure

Pooled data from the included RCTs (see Figure 4.56) suggested no difference in the incidence of multiple organ failure (4.3%, 11/257) among trauma patients who received a TEG or ROTEM-guided MHP compared with those whose MHP was guided by standard laboratory tests (3.2%, 8/250) (RR 1.33; 95% CI 0.53, 3.34; p = 0.54, I²=0%) (*GRADE: very low*).

Postpartum hysterectomy

Pooled data from the included cohort studies (see Figure 4.56) among women with severe PPH, suggested that the use of TEG or ROTEM is associated with a lower incidence of postpartum hysterectomy (8.4%) compared with treatment guided by standard care (33.8%) (RR 0.37; 95% CI 0.18, 0.77; p = 0.008; I²=54% (*GRADE: very low*).

Figure 4.55 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: thromboembolic events

	Favours TEG or RO	DTEM	Contr	ol		Risk Ratio	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl		
1.3.1 Thromboembolic events (tra	auma)								
Gonzalez 2016 (Trauma) (1)	9	56	6	55	35.7%	1.47 [0.56, 3.86]			
Baksaas-Aasen 2020 (Trauma) Subtotal (95% CI)	15	201 257	22	195 250	59.2% 94.9%	0.66 [0.35, 1.24] 0.90 [0.42 , 1. 95]	→		
Total events	24		28						
Heterogeneity: Tau ² = 0.15; Chi ² =	1.87, df = 1 (P = 0.17	'); l² = 46	5%						
Test for overall effect: Z = 0.26 (P =	0.80)								
1.3.2 Thromboembolic events (ca	irdiac)								
Paniagua 2011 (Cardiac)	0	26	0	18		Not estimable			
Weber 2012 (Cardiac)	0	50	2	50	5.1%	0.20 [0.01, 4.06]			
Subtotal (95% CI)		76		68	5.1%	0.20 [0.01, 4.06]			
Total events	0		2						
Heterogeneity: Not applicable									
Test for overall effect: Z = 1.05 (P =	0.29)								
Total (95% CI)		333		318	100.0%	0.83 [0.41, 1.66]	•		
Total events	24		30						
Heterogeneity: Tau ² = 0.11; Chi ² = 2	2.72, df = 2 (P = 0.26	i); l² = 26	5%						
Test for overall effect: Z = 0.53 (P = 0.60)							0.001 0.1 1 10 1000 Eavours TEG or ROTEM Eavours control		
Test for subgroup differences: $Chi^2 = 0.91$, $df = 1$ (P = 0.34), $l^2 = 0\%$									
Footnotes									

(1) Total number: deep vein thrombosis and pulmonary embolism

Figure 4.56 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: morbidity (multiorgan failure, need for hysterectomy)

	TEG or R	OTEM	Contr	ol	Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
1.4.3 Multiple organ failure							
Gonzalez 2016 (Trauma)	2	56	3	55	27.4%	0.65 [0.11, 3.77]	
Baksaas-Aasen 2020 (Trauma) Subtotal (95% CI)	9	201 257	5	195 250	72.6% 1 00.0%	1.75 [0.60, 5.12] 1.33 [0.53, 3.34]	
Total events	11		8				
Heterogeneity: Tau ² = 0.00; Chi ² = 0.8	88, df = 1 (P	= 0.35);	² = 0%				
Test for overall effect: $Z = 0.62$ (P = 0	.54)						
1.4.4 Postpartum hysterectomy							
Barinov 2015 (Coh, PPH)	4	90	9	29	25.7%	0.14 [0.05, 0.43]	
McNamara 2019 (Coh, obstetrics)	16	203	7	52	34.2%	0.59 [0.25, 1.35]	
Snegovskikh 2018 (Coh, obstetrics) Subtotal (95% CI)	7	28 321	31	58 1 39	40.1% 1 00.0%	0.47 [0.24, 0.93] 0.37 [0.18, 0.77]	•
Total events	27		47				
Heterogeneity: Tau ² = 0.22; Chi ² = 4.3	35, df = 2 (P	= 0.11);	² = 54%				
Test for overall effect: Z = 2.65 (P = 0	.008)	,					
							· · · · · · · · · · · · · · · · · · ·
							0.002 0.1 1 10 500
							Favours TEG or ROTEM Favours control

Table 4.80 Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding – Morbidity (thromboembolic events)

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting										
Baksaas-Aasen 2020 RCT High risk of bias	N = 396	Adult trauma patients with clinical signs of bleeding activating the local MHP * within 3 hours of injury and maximum of 1 hour after admission into the emergency department	MC, Various (Denmark, Netherlands, Norway, Germany, UK)	Viscoelastic haemostatic assays vs conventional coagulation test	Thromboembolic events *Inclusive of myocardial infarction and embolic stroke	15/201 (7.5)	22/195 (11.3)	NR	NR	
Gonzalez 2016	N = 111	Patients with severe	SC	TEG or ROTEM guided	Deep vein	8/56 (14.3)	6/55 (10.9)	NR	No significant difference	
High risk of bias		likely to require transfusion therapy	(03)	clinician's discretion and/or based on conventional coagulation tests	Pulmonary embolism	1/56 (1.8)	0/55 (0)	NR	p = 0.399 No significant difference p = 1.01	
Surgical setting										
Wikkelsø 2016 SR High quality	N = 305 (4 RCTs) Girdauskas 2010 Paniagua 2011 Shore-Lesserson 1999 Weber 2012	Adults with diffuse or excessive bleeding after cardiac surgery	SC (Germany, Spain, USA)	TEG or ROTEM guided transfusion vs clinical judgement or usual treatment with or without predefined algorithm to guide SLTs	Thromboembolic events	5/156 (3.2)	5/149 (3.4)	RR 1.04 (0.35, 3.07)	No significant difference p = 0.94 No heterogeneity $l^2 = 0 (p = 0.41)$	
Serraino 2017 SR High quality	N = 163 (2 RCTs) Girdauskas 2010 Shore-Lesserson 1999	Adult patients undergoing cardiac surgery with bleeding	Cardiac surgery (NR)	TEG or ROTEM guided transfusion (with or without other point- of-care platelet function tests) vs standard of care	Cerebrovascular accident (stroke)	5/80 (6.3)	3/81 (3.7)	RR 1.73 (0.41, 7.23)	No significant difference p = 0.45 No heterogeneity $l^2 = 0 (p = 0.68)$	
Deppe 2016 SR	N = 3975 (2 RCTs, 3 Coh)		SC, cardiac (NR)	TEG or ROTEM guided transfusion vs standard of care	Thromboembolic events	28/NR (1.3)	51/NR (2.9)	OR 0.44 (0.28, 0.70)	Favours ROTEM p = 0.0006 No heterogeneity	

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM	Risk estimate (95% Cl)	Statistical significance		
	anarysis						n/N (%)		<i>p</i> -value		
									Heterogeneity ^b		
Moderate		Patients with							l ² = 0 (<i>p</i> = 0.0005)		
quality	N = 4054	excessive bleeding			Cerebrovascular	12/NR (0.5)	18/NR (1.0)	OR 0.64 (0.31, 1.30)	No significant difference		
	(2 RCTs, 3 Coh)	after cardiac			accident (stroke)				<i>p</i> = 0.2841		
		surgery							No heterogeneity		
									$I^2 = 0 (p = 0.1345)$		
Haas 2014	N = 3865 (1 Coh)	Adult patients	SC	TEG or ROTEM guided	Composite	NR/2147 (1.77)	NR/1718 (3.19)	NR	Favours ROTEM		
Coh	Görlinger 2011	undergoing cardiac	(NR)	transfusion vs	thromboembolic				<i>p</i> = 0.011		
High risk of bias		or aortic surgery		standard of care	events						
Obstetrics and n	Obstetrics and maternity setting - no studies reported this outcome										

Paediatrics - no studies found

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: CI, confidence interval; Coh, cohort study; MC, multicentre; MHP, major haemorrhage protocol; NR, not reported; OR, odds ratio; RBC, red blood cells; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; RR, relative risk; SC, single centre; TE, thromboembolic event; TEG, thromboelastography; UK, United Kingdom; USA, United States of America

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25; (ii) mild heterogeneity if I² <25; moderate heterogeneity if I² <25; woderate heterogeneity if I² <50.

c. Calculated post-hoc using RevMan 5.4. M-H random effects

Table 4.81 Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding - Major morbidities

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results				
Study design ^a Risk of bias	(no. of trials) included in analysis		(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b	
Trauma setting										
Baksaas-Aasen 2021 RCT <i>High risk of bias</i>	N = 396	Adult trauma patients with clinical signs of bleeding activating the local MHP *randomised within 3 hours of injury and maximum of 1	MC, Various (Denmark, Netherlands, Norway, Germany, UK)	TEG or ROTEM guided transfusion vs conventional coagulation test	Multiple organ dysfunction syndrome* *Sequential Organ Failure Assessment score of 6 or more	141/164 (86)	134/159 (84)	OR 1.14 (0.62, 2.10)	No significant difference p = 0.668	
		hour after admission into			Organ failure	9/201 (4.5)	5/195 (2.6)	RR 1.75 (0.60, 5.12) °	NR	
		the entergency department			Acute kidney injury	6/201 (3.0)	6/195 (3.1)	RR 0.97 (0.32, 2.96) ^c	NR	
Haas 2014 Coh <i>Serious risk of</i> <i>bias</i>	N = 36 (1 Coh) Nienaber 2011	Adult patients with severe trauma	Trauma registry (German, Austria)	ROTEM guided administration of FC and PCC vs standard care guided transfusion of 1:1 FFP:RBC ratio	Multiple organ failure	3/18 (16.7)	11/18 (61.1)	RR 0.27 (0.09, 0.82) c	Favours ROTEM p = 0.015	
Surgical setting	(cardiac)									
Serraino 2017 SR High risk of bias	N = 424 (4 RCTs) Ak 2009 Girdauskas 2010 Paniagua 2011 Weber 2012	Adult patients undergoing cardiac surgery with bleeding	Cardiac surgery (NR)	TEG or ROTEM guided transfusion (with or without other point-of-care platelet function tests) vs standard of care	Acute kidney injury	23/217 (10.6)	39/207 (18.8)	RR 0.42 (0.20, 0.86)	Favours TEG/ROTEM p = 0.02 Mild heterogeneity $l^2 = 26 (p = 0.25)$	
Wikkelsø 2016 SR High risk of bias	N = 200 (3 RCTs) Girdauskas 2010 Paniagua 2011 Weber 2012	Adults with diffuse or excessive bleeding after cardiac surgery	SC (Germany, Spain)	TEG or ROTEM guided transfusion vs clinical judgement or usual treatment with or without predefined algorithm to guide standard laboratory tests	Dialysis dependent renal failure	16/103 (15.5)	30/97 (30.9)	RR 0.46 (0.28, 0.76)	Favours TEG/ROTEM p = 0.0028 No heterogeneity $l^2 = 0 (p = 0.48)$	
Deppe 2016 SR <i>High risk of bias</i>	N = 4263 (3 RCTs, 2 Coh) NR	Patients with excessive bleeding after cardiac surgery	SC, cardiac (NR)	TEG or ROTEM guided transfusion vs standard of care	Acute kidney injury	142/NR (6.0)	150/NR (7.8)	OR 0.77 (0.61, 0.98)	Favours TEG/ROTEM p = 0.0403 No heterogeneity	

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results					
Study design ^a Risk of bias	(no. of trials) included in analysis		(Location)			TEG/ROTEM n/N (%)	No TEG/ROTEM n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b		
									l ² = 0 (<i>p</i> = 0.0278)		
					RCTs only	NR	NR	OR 0.54 (0.27, 1.06)	No significant difference p = 0.1001 Heterogeneity NR		
Haas 2014 RCT High risk of bias	N = 100 (1 RCT) Weber 2012	Adult patients undergoing cardiac or aortic surgery	SC (NR)	TEG or ROTEM guided transfusion vs standard of care	Composite AEs (Acute renal failure, sepsis, TE, allergic reaction)	NR/NR (8)	NR/NR (38)	RR 0.30 (0.09, 1.03) ^c	Favours ROTEM ρ < 0.001		
Surgical setting	(Liver)										
Saner 2016 Coh Serious risk of bias	N = 200 (1 Coh) Leon Justel 2015	Patients with end-stage liver disease undergoing invasive procedure	SC (Spain)	TEG or ROTEM guided transfusion vs standard of care	Acute kidney injury	2/100 (2)	17/100 (17)	RR 0.12 (0.03, 0.50) ^c	Favours ROTEM p = 0.001		
Obstetrics and n	naternity										
Roullet 2018 Coh Serious risk of bias	N = 179 (2 Coh) Mallaiah 2015 Snegovskikh 2018	Patients with severe PPH	SC (UK, US)	ROTEM guided algorithm vs standard of care	Postpartum hysterectomy	10/79 (12.7)	37/100 (37)	RR 0.45 (0.25, 0.83) ^c	Favours ROTEM p = 0.048 No heterogeneity $l^2 = 0\% (p = 0.55)$		
Amgalan 2020 Coh Serious risk of bias	N = 100 (1 Coh) McNamara 2019	Women with MOH (estimated blood loss >1500mL) associated with coagulopathy	SC (UK)	ROTEM guided algorithm vs standard of care	Postpartum hysterectomy	16/203	7/52	RR 0.59 (0.25, 1.35) ^c	No significant difference ρ = 0.21		

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: AE, adverse event; CI, confidence interval; Coh, cohort study; FC fibrinogen concentrate; FFP, fresh frozen plasma; MC, multicentre; MHP, major haemorrhage protocol; NR, not reported; OR, odds ratio; PCC, prothrombin complex concentrate; PPH, postpartum haemorrhage; RBC, red blood cells; ROTEM, rotational thromboelastometry; RR, relative risk; SC, single centre; TE, thromboembolic event; TEG thromboelastography; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

4.9.3.3 Transfusion volume

A summary of the evidence relating to the volume of blood components transfused in patients with critical bleeding in whom TEG or ROTEM were used as part of a major haemorrhage protocol is presented in Table 4.82.

Red blood cells

A meta-analysis of data from the RCT and cohort studies included in this review (see Figure 4.57) showed a significant reduction in the volume of RBC transfused in patients with critical bleeding (any setting) who received a TEG or ROTEM-guided MHP (n=669) compared with those whose MHP was guided by standard laboratory tests (n=1089). The difference corresponded to around 2 units of RBC saved (SMD –0.38; 95% CI –0.61, –0.15; p = 0.001, I²= 75%).

Available data from 2 RCTs suggested that the volume of RBC transfused was not different between groups (SMD –0.06; 95% CI –0.38, 0.26; p = 0.73, I²= 0%), but data were not reported in 3 studies and 2 studies suggested an effect favouring TEG or ROTEM but did not provide suitable data for analysis.

Among the included observational cohort studies, a statistically significant reduction in the volume of RBC transfused was observed among patients who received a TEG or ROTEM-guided transfusion protocol (n=588) compared with those who received haemostatic management guided by a transfusion algorithm/haemorrhage protocol or standard laboratory tests (n=1017) (SMD -0.46; 95% CI -0.92, -0.28; p = 0.0005; I²= 78%).

In the trauma setting (see Figure 4.58), data from one RCT suggested that the use of a ROTEM-guided MHP did not reduce the volume of RBC transfused when compared to an MHP guided by standard laboratory tests (SMD –0.13; 95% CI –0.50, 0.25; p = 0.51). Among the cohort studies a significant association was observed (SMD –0.41; 95% CI –0.68, –0.14; p = 0.03; $I^2 = 78\%$).

In patients with diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery (see Figure 4.58), data from one small RCT suggested that there was no difference in volume of RBC transfused comparing a ROTEM-guided MHP with routine transfusion therapy based on standard laboratory tests (SMD 0.12; 95% CI –0.48, 0.72; p = 0.69). Data were not reported in 2 studies and 2 studies suggested an effect favouring TEG or ROTEM but did not provide suitable data for analysis.

Among women with severe PPH (see Figure 4.58), data from one observational study suggested that the use of ROTEM is associated with a statistically significant reduction in the volume of RBC transfused (around one unit saved) compared with management of coagulopathy guided by standard laboratory tests (SMD of –0.82; 95% CI –1.25, –0.39; p = 0.0002). One study suggested there was no reduction the median volume of RBC transfused. One study did not report this outcome.

Figure 4.57 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: RBC transfusion volume (units), by study design.

	EG or ROTEM Co			Control		Std. Mean Difference		Std. Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.5.1 RCTs									
Weber 2012 (Cardiac) (1)	0	0	50	0	0	50		Not estimable	
Nuttall 2001 (Cardiac) (2)	0	0	41	0	0	51		Not estimable	
Paniagua 2011 (Cardiac) (3)	7.096	5.576	26	6.416	5.464	18	7.5%	0.12 [-0.48, 0.72]	
Gonzalez 2016 (Trauma) (4) Subtotal (95% CI)	13.96	12.68	55 81	15.65	13.85	54 72	11.0% 18.6%	-0.13 [-0.50, 0.25] -0.06 [-0.38, 0.26]	
Heterogeneity: Tau ² = 0.00; Chi ² = 0.47	, df = 1 (P = 0.4	9); l² =	0%					
Test for overall effect: Z = 0.35 (P = 0.7	3)								
155 Cohe									
Neinaber 2011 (Cob. trauma) (5)	٥	٥	10	٥	٥	18		Not estimable	
Görlinger 2012a (Cob. trauma) (6)	0	0	0	0	0	0		Not estimable	
McNamara 2019 (Coh, obstetrics) (7)	0	0	203	0	0	52		Not estimable	
Unruh 2019 (Coh. trauma)	6	5.2	47	11	3.7	20	8.2%	-1.03 [-1.58, -0.47]	_ _
Barinov 2015 (Coh. PPH)	4.813	1.255	92	6.102	2.28	29	10.1%	-0.82 [-1.25, -0.39]	
Prat 2017 (Coh. trauma)	2	2.2	85	2	1.5	55	11.7%	0.00 [-0.34, 0.34]	+
Wang 2017 (Coh. trauma)	4	7	86	9	10	80	12.2%	-0.58 [-0.89, -0.27]	
Guth 2019 (Coh, trauma)	2	3	102	6	7.4	102	12.7%	-0.71 [-0.99, -0.42]	
Nardi 2015 (Coh, trauma)	6.5	4.8	96	8.1	6.7	130	13.0%	-0.27 [-0.53, -0.00]	
Schochl 2011 (Coh, trauma)	5.5	7	80	6	5.2	601	13.5%	-0.09 [-0.33, 0.14]	-
Subtotal (95% CI)			588			1017	81.4%	-0.46 [-0.72, -0.20]	\bullet
Heterogeneity: Tau ² = 0.09; Chi ² = 27.3	7, df = 6	(P = 0.	0001);	² = 78%					
Test for overall effect: Z = 3.48 (P = 0.0	005)								
Total (95% CI)			669			1089	100.0%	-0.38 [-0.61, -0.15]	•
Heterogeneity: Tau ² = 0.09; Chi ² = 31.4	4, df = 8	(P = 0.	0001);	² = 75%					
Test for overall effect: Z = 3.26 (P = 0.0	01)	,	,,						-4 -2 0 2 4
Test for subgroup differences: Chi ² = 3.	, 72, df =	1 (P = 0).05), l²	= 73.1%	6				Favours LEG OF KUTEM Favours control
Footnotes		,	,.						

(1) Favours TEG or ROTEM (p<0.001). Data reported as median (IQR): 3 (2, 6) vs 5 (4, 9)

(2) Favours TEG or ROTEM (p=0.039). Data reported as median (range): 2 (0, 9) vs 3 (0, 70)

(3) converted from mLs (250mL/U)

(4) As reported by Fahrendorff 2017 (total cumulative RBC transfusion needs)

(5) Favours TEG or ROTEM (p<0.005). Data presented as median (IQR): 3 (0, 5) vs 12.5 (8, 20).

(6) An estimated a 42% and 33% reduction in the total volume of RBCs transfused per year after the implementation of goal-directed therapy

(7) No difference between groups (p=0.158). Data reported as median (IQR) 3 (2, 5) vs 3 (2, 4).

Figure 4.58 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: RBC transfusion volume (units), by setting.

	TEG	or ROT	EM	(Control			Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.8.1 Trauma (RCTs)									
Baksaas-Aasen 2020 (Trauma) (1)	0	0	201	0	0	195		Not estimable	
Gonzalez 2016 (Trauma)	13.96	12.68	55	15.65	13.85	54	11.0%	-0.13 [-0.50, 0.25]	
Subtotal (95% CI)			55			54	11 .0 %	-0.13 [-0.50, 0.25]	\bullet
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.66 (P = 0.4	51)								
1.8.2 Trauma (Coh)									
Görlinger 2012a (Coh. trauma) (2)	0	0	0	0	0	0		Not estimable	
Neinaber 2011 (Coh, trauma) (3)	0	0	18	0	0	18		Not estimable	
Unruh 2019 (Coh. trauma)	6	5.2	47	11	3.7	20	8.2%	-1.03 [-1.58, -0.47]	
Prat 2017 (Coh. trauma)	2	2.2	85	2	1.5	55	11.7%	0.00 [-0.34, 0.34]	<u> </u>
Wang 2017 (Coh, trauma)	4	7	86	9	10	80	12.2%	-0.58 [-0.89, -0.27]	
Guth 2019 (Coh. trauma)	2	3	102	6	7.4	102	12.7%	-0.71 [-0.99, -0.42]	_ _
Nardi 2015 (Coh, trauma)	6.5	4.8	96	8.1	6.7	130	13.0%	-0.27 [-0.53, -0.00]	_
Schochl 2011 (Coh. trauma)	5.5	7	80	6	5.2	601	13.5%	-0.09 [-0.33, 0.14]	
Subtotal (95% CI)			496			988	71.3%	-0.41 [-0.68, -0.14]	\bullet
Heterogeneity: Tau ² = 0.09; Chi ² = 23.	11, df = 5	(P = 0.0	0003);	l² = 78%	, D				
Test for overall effect: Z = 2.95 (P = 0.	003)								
1.8.3 Surgical									
Weber 2012 (Cardiac) (4)	0	0	50	0	0	50		Not estimable	
Nuttall 2001 (Cardiac) (5)	0	0	41	0	0	51		Not estimable	
Paniagua 2011 (Cardiac) (6)	7.096	5.576	26	6.416	5.464	18	7.5%	0.12 [-0.48, 0.72]	
Subtotal (95% CI)			26			18	7.5%	0.12 [-0.48, 0.72]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.39 (P = 0.	69)								
184 Obstetrics and maternity									
McNamara 2019 (Cob. obstatrics) (7)	٥	٥	203	0	0	52		Not estimable	
Barinov 2015 (Coh. PPH)	4 813	1 255	200 92	6 102	2 28	29	10.1%	-0.82 [-1.25 -0.39]	
Subtotal (95% CI)	4.010	1.200	92	0.102	2.20	29	10.1%	-0.82 [-1.25, -0.39]	\bullet
Heterogeneity: Not applicable									
Test for overall effect: Z = 3.74 (P = 0.1	0002)								
Total (95% CI)			669			1089	100 0%	-0.38 [-0.61 -0.15]	
Hotorogonoity: $T_{21}^2 = 0.00$; $Ch^2 = 21$	11 df - 0	(P _ 0 (000	12 - 750	<u>c</u>		10010/0	0.00 [0.01, 0.10]	▼ + + + + + + + + + + + + + + + + + +
Test for overall effect: $7 = 3.26 (P = 0.0)$	+++, ui = 8 101)	יר = 0.0	JUU I);	1 = 70%	D				-2 -1 0 1 2
Test for subgroup differences: $Chi^2 = 9$	18 df -	3 (P - 0	0/1) 12	- 64 60	/_				Favours TEG or ROTEM Favours Control
Footnotes	.+0, ui =	J (F = 0	.04), F	- 04.03	ru				
(4) Outload and a stars set this subserve									

(1) Study did not report this outcome.

(2) An estimated 42% and 33% reduction in the total volume of RBCs transfused per year after the implementation of goal-directed therapy.

(3) Favours TEG or ROTEM (p<0.005). Data presented as median (IQR): 3 (0, 5) vs 12.5 (8, 20).

(4) Favours TEG or ROTEM (p<0.001). Data reported as median (IQR): 3 (2, 6) vs 5 (4, 9)

(5) Favours TEG or ROTEM (p=0.039). Data reported as median (range): 2 (0, 9) vs 3 (0, 70)

(6) Data converted from mLs (250mL/U)

(7) No difference between groups (p=0.158). Data reported as median (IQR) 3 (2, 5) vs 3 (2, 4).

Table 4.82 Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding - RBC transfusion volume

Study ID	Sample size	Patient population	Setting	Comparison	Outcome	Results					
Study design ^a Risk of bias	(no. of trials) included in analysis		(Location)			TEG/ROTEM Mean ± SD (n)	No TEG/ROTEM Mean ± SD (n)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
Various settings											
Fahrendorff 2017 SR High risk of bias	N = 453 (6 studies) Shore-Lesserson 1999 Wang 2010 Schaden 2012 Barinov 2015 Gonzalez 2016 Cao 2016	Patients with an acute need for blood components due to bleeding (includes trauma, burns excision, liver transplant, PPH and cardiac surgery)	Various	TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional coagulation tests	RBC transfusion volume, Units	Mean ± SD (260) 1.416 ± 1.948 (53) 14.2 ± 7.1 (14) 3.1 ± 2.1 (14) 4.813 ± 1.255 (92) 13.96 ± 12.68 (55) 4.5 ± 1.5 (32)	Mean ± SD (193) 1.9 ± 2.372 16.7 ± 12.8 (14) 4.8 ± 3 (16) 6.102 ± 2.28 (29) 15.65 ± 13.85 (54) 7.1 ± 1.2 (28)	SMD -0.64 (-1.12, -0.15) -0.22 (-0.61, 0.16) -0.23 (-0.98, 0.51) -0.63 (-1.37, 0.11) -0.82 (-1.25, -0.39) -0.13 (-0.59, 0.25) -1.88 (-2.49, -1.26)	Favours TEG/ROTEM p = 0.01 Substantial heterogeneity l ² = 82 (p = 0.001)		
Trauma setting											
Bugaev 2020 SR Serious risk of bias	N = 1459 (1 RCT, 6 Coh) Schaden 2012 Unruh 2019 Gonzalez 2016 Prat 2017 Guth 2019 Nardi 2015 Schochl 2011	Severely injured trauma patients	SC (not reported)	ROTEM or TEG vs no ROTEM or TEG	RBC transfusion volume, Units	N=480 3.1±1.6 (14) 6±5.2 (47) 9.5±8.1 (56) 2±2.2 (85) 2±3 (102) 6.5±4.8 (96) 5.5±7 (80)	N=979 4.3±2.2 (16) 11±3.7 (20) 11±8.1 (55) 2±1.5 (55) 6±7.4 (102) 8.1±6.7 (130) 6±5.2 (601)	SMD -0.38 (-0.64, -0.12) SMD -0.85 (-1.60, -0.10) SMD -1.03 (-1.58, -0.47) SMD 0.18 (-0.56, 0.19) SMD 0.00 (-0.34, 0.34) SMD -0.71 (-0.99, -0.42) SMD -0.72 (-0.53, -0.00) SMD -0.09 (-0.33, 0.14)	Favours TEG/ROTEM p = 0.004 Significant heterogeneity I ² = 74% (p = 0.0008)		
Wang 2017 Coh <i>Serious risk of bias</i>	N = 166 (1 Coh)	Patients with traumatic liver and/or spleen injury	SC (US)	TEG guided algorithm vs clinical judgement or usual treatment	RBC transfusion volume, Units	4 ± 7 (86)	9 ±10 (80)	NR	<i>Favours TEG</i> <i>p</i> < 0.01		
Haas 2014 SR High risk of bias	N = 5590 (1 Coh) Görlinger 2012a Görlinger 2012b	Adult patients with severe trauma	SC (Germany, Austria)	ROTEM guided algorithm vs standard of care	RBC transfusion volume, Units	Total per year 1282 888	Total per year 2215 1332	42% reduction 33% reduction	<i>Favours ROTEM</i> <i>p</i> = NR		
	N = 36 (1 Coh) Nienaber 2011			Algorithm with FC and PCC vs 1:1 FFP:RBC ratio	hrs after admission 0–6 hrs >24 hrs	Median (IQR) 1 (0, 3) 3 (0, 5)	Median (IQR) 7.5 (4, 12) 12.5 (8, 20)	NR NR	p < 0.005 p < 0.005		
Surgical setting											
Wikkelsø 2016 SR High risk of bias	N = NR (2 RCTs) Weber 2012 Nuttall 2001		SC (Germany, Spain, US)		RBC transfusion volume, Units	Median (IQR) 3 (2, 6) Median (range)	Median (IQR) 5 (4, 9) Median (range)		Favours TEG/ROTEM p < 0.001		
						2 (0, 9)	3 (0, 70)		<i>p</i> = 0.039		

Study ID	Sample size	Patient population S	Setting	Comparison	Outcome	Results					
Study design ^a Risk of bias	(no. of trials) included in analysis		(Location)			TEG/ROTEM Mean ± SD (n)	No TEG/ROTEM Mean ± SD (n)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
	N = 44 (1 RCT) Paniagua 2011	Adults or children with diffuse or excessive bleeding after cardiac surgery, orthotopic liver transplant or burns excision		TEG or ROTEM guided transfusion vs clinical judgement or usual treatment with or without predefined algorithm to guide SLTs	RBC transfusion volume, mL	Mean ± SD (n) 1774 ± 1394 (26)	Mean ± SD (n) 1604 ± 1366 (18)	SMD ^c 0.12 (-0.48, 0.72)	No significant difference <i>p</i> = NR		
Haas 2014 SR Serious risk of bias	N = 5338 (1 Coh) Görlinger 2012b	Adult patients undergoing visceral surgery or liver transplant	SC (Germany, Austria)	ROTEM guided algorithm vs standard of care	RBC transfusion volume, Units	Units per year 1319	Units per year 3454	62% reduction	Favours ROTEM p = NR		
	N = 100 (1 RCT) Weber 2012	Adult patients undergoing cardiac or aortic surgery	SC (Germany)	ROTEM guided algorithm vs standard of care	RBC transfusion volume, Units	Median (IQR) 3 (2, 6)	Median (IQR) 5 (4, 9)	NR	Favours ROTEM p < 0.001		
Obstetrics settin	ng										
Amgalan 2020 Coh Serious risk of bias	N = 100 (1 Coh) McNamara 2019	Women with MOH (estimated blood loss >1500mL) associated with coagulopathy	SC (UK)	ROTEM guided algorithm vs standard of care	RBC transfusion volume, Units	Median (IQR) 3 (2, 5) (n=203)	Median (IQR) 3 (2, 4) (n=52)	NR	No significant difference p = 0.158		

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: CI, confidence interval; Coh, cohort study; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; hrs, hours; IQR, interquartile range; MC, multicentre; MD, mean difference; n, number; NR, not reported; PCC, prothrombin complex concentrate; PLT, platelets; PPH, postpartum haemorrhage; RBC, red blood cells; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; SC, single centre; SD, standard deviation; SMD, standard mean difference; SoC, standard of care; TEG, thromboelastography; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

Fresh frozen plasma

A summary of the evidence relating to the volume of FFP transfused in patients with critical bleeding is presented in Table 4.83

A meta-analysis of available data from the RCT and cohort studies included in this review (see Figure 4.59) showed a significant reduction in the volume of FFP transfused in patients with critical bleeding (any setting) who received a TEG or ROTEM-guided MHP (n=594) compared with those who received a transfusion protocol guided by standard laboratory tests (n=572). The difference corresponded to around 2.4 units of FFP saved (SMD –0.62; 95% CI –1.19, –0.05; p = 0.03, I²= 95%).

Available data from the RCTs suggested that the volume of FFP transfused was not different between groups (SMD 0.02; 95% CI –0.30, 0.33; p = 0.93; I²= 0%) but data were not able to be included for 2 studies that suggested an effect favouring TEG or ROTEM. Among the included observational cohort studies, a statistically significant reduction in the volume of FFP transfused was observed among patients who a TEG or ROTEM-guided transfusion protocol (n=513) compared with those who received haemostatic management guided by a transfusion algorithm/haemorrhage protocol or standard laboratory tests (n=500) (SMD –0.82; 95% CI –1.51, –0.12; p = 0.02; I²= 96%).

In the trauma setting (see Figure 4.60), data from one RCT suggested that the use of TEGguided MHP did not reduce the volume of FFP transfused when compared with an MHP guided by standard laboratory tests (SMD –0.01; 95% CI –0.39, 0.37; p = 0.96). Among the cohort studies no significant association was observed (SMD –0.39; 95% CI –1.01, 0.23; p =0.22; $I^2 = 95\%$), noting FFP transfusion volumes were not reported for all studies, possibly due to the direction of effect being unfavourable for the intervention. Taken together the pooled data from the RCT and cohort studies suggests that the use of a TEG or ROTEMguided transfusion protocol does not reduce the volume of FFP transfused when compared to a transfusion protocol not guided by TEG or ROTEM (SMD –0.32; 95% CI – 0.86, 0.21; p = 0.23; $I^2 = 94\%$).

In patients with diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery (see Figure 4.60), data from one small RCT and one small cohort study suggested that there was no difference in volume of FFP transfused comparing a ROTEM-guided transfusion protocol with routine transfusion therapy based on standard laboratory tests (SMD –0.50; 95% CI –1.91, 0.91; p = 0.49; $l^2 = 70\%$). Data were not reported in 2 studies and 2 studies suggested an effect favouring TEG or ROTEM but did not provide suitable data for analysis.

Among women with severe PPH (see Figure 4.60), data from one observational study suggested that blood component therapy guided by TEG or ROTEM is associated with a large reduction in the volume of FFP transfused (around 4.4 units saved) compared with management of coagulopathy guided by standard laboratory tests (SMD of –2.73; 95% Cl – 3.28, –2.19; p < 0.0001). The other 2 studies did not report this outcome.

Figure 4.59 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: FFP transfusion volume (units), by study design.

	TEG	or RO	ГЕМ	(Control		;	Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
1.9.1 RCTs									
Nuttall 2001 (Cardiac) (1)	0	0	41	0	0	51		Not estimable	
Weber 2012 (Cardiac) (2)	0	0	50	0	0	50		Not estimable	
Baksaas-Aasen 2020 (Trauma) (3)	0	0	201	0	0	195		Not estimable	
Paniagua 2011 (Cardiac) (4)	3.196	4.752	26	2.828	3.988	18	10.9%	0.08 [-0.52, 0.68]	+
Gonzalez 2016 (Trauma)	7.49	7.37	55	7.57	7.86	54	11.8%	-0.01 [-0.39, 0.37]	+
Subtotal (95% CI)			81			72	22.7%	0.02 [-0.30, 0.33]	•
Heterogeneity: Tau ² = 0.00; Chi ² = 0	.06, df =	1 (P = ().80); l²	= 0%					
Test for overall effect: Z = 0.09 (P =	0.93)								
1.9.2 Cohs									
Görlinger 2012a (Coh, trauma) (5)	0	0	0	0	0	0		Not estimable	
Hanke 2012 (Coh, Cardiac)	1.6	2.2	5	9.2	6.6	5	6.8%	-1.40 [-2.86, 0.07]	
Barinov 2015 (Coh, PPH)	4.8	1.537	92	9.25	1.862	29	11.2%	-2.73 [-3.28, -2.19]	
Unruh 2019 (Coh, trauma)	4.5	4.1	47	4	4.1	20	11.2%	0.12 [-0.40, 0.64]	+-
Wang 2017 (Coh, trauma)	1	5	86	5	6	80	12.0%	-0.72 [-1.04, -0.41]	+
Guth 2019 (Coh, trauma)	0.5	1.5	102	5	5.2	102	12.0%	-1.17 [-1.47, -0.87]	-
Prat 2017 (Coh, trauma)	2	2.6	85	1	1.5	134	12.1%	0.50 [0.22, 0.77]	+
Nardi 2015 (Coh, trauma)	4.2	4.6	96	9	9.5	130	12.1%	-0.61 [-0.88, -0.34]	*
Subtotal (95% CI)			513			500	77.3%	-0.82 [-1.51, -0.12]	\bullet
Heterogeneity: Tau ² = 0.80; Chi ² = 1	42.73, df	= 6 (P	< 0.000	001); l² =	= 96%				
Test for overall effect: Z = 2.30 (P =	0.02)								
Total (95% CI)			594			572	100.0%	-0.62 [-1.19, -0.05]	•
Heterogeneity: Tau ² = 0.69; Chi ² = 1	53.85, df	= 8 (P	< 0.000	001); l² =	= 95%			-	
Test for overall effect: Z = 2.12 (P =	0.03)								-4 -2 U 2 4
Test for subgroup differences: Chi2 =	- 4.53, df	= 1 (P	= 0.03)	, l² = 77	.9%				Favouis IEG/RUTEIVI Favouis Cutilioi
Footnotes			,						
(1) Equation TEC or POTEM (p. 0.00)5) Doto	roporto	d ac m	odion (r		0 10	wc 1 (0 7	5)	

(1) Favours TEG or ROTEM (p=0.005). Data reported as median (range): 2 (0, 10) vs 4 (0, 75).

(2) Favours TEG or ROTEM (p<0.001). Data reported as median (IQR): 0 (0, 3) vs 5 (3, 8).

(3) Study did not report this outcome.

(4) Converted from mLs (250 mL/U)

(5) An estimated 79% and 94% reduction in the total volume of FFP transfused per year after the implementation of goal-directed therapy (two trauma centres).

Figure 4.60 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: FFP transfusion volume (units), by setting.

	TEG	or ROT	EM	(Control			Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% Cl
1.10.1 Trauma (RCTs)									
Baksaas-Aasen 2020 (Trauma) (1)	0	0	201	0	0	195		Not estimable	
Gonzalez 2016 (Trauma) Subtotal (95% CI)	7.49	7.37	55 55	7.57	7.86	54 54	11.8% 11.8%	-0.01 [-0.39, 0.37] -0.01 [-0.39, 0.37]	↓
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.05 (P =	0.96)								
1.10.2 Trauma (Coh)									
Görlinger 2012a (Coh, trauma) (2)	0	0	0	0	0	0		Not estimable	
Unruh 2019 (Coh, trauma)	4.5	4.1	47	4	4.1	20	11.2%	0.12 [-0.40, 0.64]	
Wang 2017 (Coh, trauma)	1	5	86	5	6	80	12.0%	-0.72 [-1.04, -0.41]	
Guth 2019 (Coh, trauma)	0.5	1.5	102	5	5.2	102	12.0%	-1.17 [-1.47, -0.87]	
Prat 2017 (Coh, trauma)	2	2.6	85	1	1.5	134	12.1%	0.50 [0.22, 0.77]	
Nardi 2015 (Coh, trauma)	4.2	4.6	96	9	9.5	130	12.1%	-0.61 [-0.88, -0.34]	
Subtotal (95% CI)			416			466	59.3%	-0.39 [-1.01, 0.23]	
Heterogeneity: Tau ² = 0.47; Chi ² = 7 Test for overall effect: Z = 1.22 (P =	6.74, df = 0.22)	= 4 (P <	0.0000)1); l² =	95%				
1.10.3 Surgical (RCTs)									
Nuttall 2001 (Cardiac) (3)	0	0	41	0	0	51		Not estimable	
Weber 2012 (Cardiac) (4)	0	0	50	0	0	50		Not estimable	
Paniagua 2011 (Cardiac) (5) Subtotal (95% CI)	3.196	4.752	26 26	2.828	3.988	18 18	10.9% 10.9%	0.08 [-0.52, 0.68] 0.08 [-0.52, 0.68]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.26 (P =	0.79)								
1.10.4 Surgical (Coh)									
Hanke 2012 (Coh, Cardiac) Subtotal (95% CI)	1.6	2.2	5 5	9.2	6.6	5 5	6.8% 6.8%	-1.40 [-2.86, 0.07] -1.40 [-2.86, 0.07]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 1.86 (P =	0.06)								
1.10.5 Obstetrics									
Barinov 2015 (Coh, PPH) Subtotal (95% CI)	4.8	1.537	92 92	9.25	1.862	29 29	11.2% 11.2%	-2.73 [-3.28, -2.19] - 2.73 [-3.28, -2.19]	→
Heterogeneity: Not applicable									
Test for overall effect: Z = 9.83 (P <	0.00001)								
Total (95% CI)			594			572	100.0%	-0.62 [-1.19, -0.05]	•
Heterogeneity: Tau ² = 0.69; Chi ² = 1	53.85, df	= 8 (P -	< 0.000	001); l² =	95%			-	
Test for overall effect: Z = 2.12 (P =	0.03)								-2 -1 U I Z
Test for subgroup differences: Chi2 =	= 74.59, 0	lf = 4 (P	< 0.00	001), l²	= 94.6%	6			

Footnotes

(1) Study did not report this outcome.

(2) An estimated 79% and 94% reduction in the total volume of FFP transfused per year after the implementation of goal-directed therapy (two trauma centres).

(3) Favours TEG or ROTEM (p=0.005). Data reported as median (range): 2 (0, 10) vs 4 (0, 75)

(4) Favours TEG or ROTEM (p<0.001). Data reported as median (IQR): 0 (0, 3) vs 5 (3, 8).

(5) converted from mLs (250 mL/U)

Table 4.83 Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding - FFP transfusion volume

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ª	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM Mean ± SD (n)	No TEG/ROTEM Mean ± SD (n)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
Various setting	gs								
Fahrendorff 2017 SR High risk of bias	N = 423 (5 studies) Shore-Lesserson 1999 Wang 2010 Barinov 2015 Gonzalez 2016 Cao 2016	Patients with an acute need for blood components due to bleeding (includes trauma, burns excision, liver transplant, PPH and cardiac surgery)	Various	TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional coagulation tests	FFP transfusion volume, Units	Mean ± SD (246) 0.133 ± 0.526 (53) 12.8 ± 7 (14) 4.8 ± 1.537 (92) 7.49 ± 7.37 (55) 0.867 ± 0.17 (32)	Mean ± SD (177) 0.804 ± 1.715 21.5 ± 12.7 (14) 9.25 ± 1.862 (29) 7.57 ± 7.86 (54) 1.904 ± 0.152 (28)	SMD -1.98 (-3.41, -0.54) -0.53 (-0.92, -0.14) -0.82 (-1.60, -0.05) -2.73 (-3.28, -2.19) -0.01 (-0.39, 0.37) -6.32 (-7.60, -5.05)	Favours TEG/ROTEM p = 0.007 Substantial heterogeneity $I^2 = 97 (p = 0.00001)$
Trauma settin	g								
Bugaev 2020 SR High risk of bias Roullet 2018 RCT High risk of bias	N = 827 (5 studies) Unruh 2019 Gonzalez 2016 Guth 2019 Prat 2017 Nardi 2015 N = NR (1 RCT) Gonzalez 2016	Severely injured trauma patients Patients with severe trauma	Trauma (NR) Trauma (NR)	ROTEM or TEG vs no ROTEM or TEG TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional	FFP transfusion volume, Units FFP transfusion volume, Units	N=386 4.5±4.1 (n=47) 5±4.4 (n=56) 0.5±1.5 (n=102) 2±2.6 (n=85) 4.2±4.6 (n=96) Authors note that the more FFP early comp *See Fahrendorff 2017 for	N=441 4±4.1 (n=20) 6±3.7 (n=55) 5±5.2 (n=102) 1±1.5 (n=134) 9±9.5 (n=130) a group receiving the r pared to the TEG group values	SMD -0.29 (-0.91, 0.34) SMD 0.12 (-0.40, 0.64) SMD -0.24 (-0.62, 0.13) SMD -1.17 (-1.47, -0.87) SMD 0.50 (0.22, 0.77) SMD -0.61 (-0.88, -0.34) outine tests received	No significant difference p = 0.36 Significant heterogeneity $l^2 = 94\%$ ($p < 0.00001$) No significant difference p = NR
Wang 2017 Coh Serious risk of bias	N = 166 (1 Coh)	Patients with traumatic liver and/or spleen injury	SC (US)	TEG guided algorithm vs clinical judgement or usual treatment	FFP transfusion volume, Units	1 ± 5 (86)	5 ± 6 (80)	NR	Favours TEG p < 0.01
Haas 2014 Coh Serious risk of bias	N = 5590 (1 Coh) Görlinger 2012a Görlinger 2012a	Adult patients with severe trauma	SC (Austria, Germany)	ROTEM guided algorithm vs standard of care	FFP transfusion volume, Units	Total per year (n) 48 (4) 261 (NR)	Total per year (n) 756 (63) 1221 (NR)	94% reduction 79% reduction	<i>p</i> = NR
Surgical settin	ŋg								-
Wikkelsø 2016	N = NR (2 RCTs)					Median (IQR)	Median (IQR)		Favours TEG/ROTEM

Study ID	dy ID Sample size Patient Setting Comparise				Outcome	Results					
Study design ^a	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM Mean ± SD (n)	No TEG/ROTEM Mean ± SD (n)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b		
SR High risk of	Weber 2012	Adults or children with diffuse or	SC (Germany, Spain, US)	TEG or ROTEM guided transfusion	FFP transfusion	0 (0, 3) Median (range)	5 (3, 8) Median (range)		<i>p</i> < 0.001		
DIAS	N = 44 (1 RCT) Paniangua 2011	excessive bleeding after cardiac surgery, orthotopic liver transplant or burns excision		vs clinical judgement or usual treatment with or without predefined algorithm to guide SLTs	FFP transfusion volume, mL	Mean ± SD (n) 799 ± 1188 (26)	4 (0, 75) Mean ± SD (n) 707 ± 997 (18)	0.08 (-0.52, 0.68)	<i>p</i> = 0.005 <i>No significant difference</i> <i>p</i> = NR		
Haas 2014 Coh	N = 5338 (1 Coh) Görlinger 2012b	Adult patients undergoing visceral surgery or liver transplant	SC (Germany)	ROTEM guided algorithm vs standard of care	FFP transfusion volume, Units	Units per year 223	Units per year 4465	95% reduction	Favours ROTEM p = NR		
Serious risk of bias	N = 10 (1 Coh) Hanke 2012	Adult patients undergoing cardiac	SC (NR)	ROTEM guided algorithm vs	FFP transfusion	Mean ± SD (n) 1.6 ± 2.2 (5)	Mean ± SD (n) 9.2 ± 6.6 (5)	MD -7.60 (-13.7, -1.5) ^c	Favours ROTEM p = 0.038		
	N = 100 (1 RCT) Weber 2012	or aortic surgery		standard of care	volume, Units	Median (IQR) 0 (0, 3)	Median (IQR) 5 (3, 8)	NR	<i>p</i> < 0.001		

Obstetrics setting – no studies reporting data

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: CI, confidence interval; Coh, cohort study; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; hrs, hours; IQR, interquartile range; MC, multicentre; MD, mean difference; n, number; NR, not reported; PCC, prothrombin complex concentrate; PLT, platelets; PPH, postpartum haemorrhage; RBC, red blood cells; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; SC, single centre; SD, standard deviation; SMD, standard mean difference; SoC, standard of care; SLTs, standard laboratory tests; TEG, thromboelastography; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

Platelets

A summary of the evidence relating to the volume of platelets (PLT) and other blood components transfused in patients with critical bleeding is presented in Table 4.84.

A meta-analysis of available data from the RCT and cohort studies included in this review (see Figure 4.61) showed no significant reduction in the volume of PLT transfused in patients with critical bleeding (any setting) who received a TEG or ROTEM-guided transfusion protocol (n=402) compared with haemostatic management guided by an MHP, standard laboratory tests or clinical judgement with or without laboratory tests (n=331) (SMD -0.21; 95% CI -0.51, 0.09; p = 0.17, $I^2 = 72\%$).

Available data from the RCTs suggested that the volume of PLT transfused was not different between groups (SMD 0.02; 95% CI –0.59, 0.64; p = 0.94; $l^2 = 65\%$) but data were not able to be included for 2 studies that suggested an effect favouring TEG or ROTEM. Among the observational cohort studies, the available data suggested there a non-significant reduction in the volume of PLT transfused (around one unit saved) among patients who received a TEG or ROTEM-guided transfusion protocol (n=284) compared with those who received haemostatic management guided by a transfusion algorithm/haemorrhage protocol or or standard laboratory tests (n=284) (SMD –0.31; 95% CI –0.64, 0.03; p = 0.07; $l^2 = 96\%$).

In the trauma setting (see Figure 4.62), data from one RCT suggested that the use of a TEG-guided MHP did not reduce the volume of PLT transfused when compared treatment not guided by TEG or ROTEM (SMD 0.30; 95% CI –0.12, 0.72; p = 0.16). Among the cohort studies a significant association was observed (SMD –0.43; 95% CI –0.78, –0.08; p = 0.02; $I^2 = 67\%$), noting PLT transfusion volumes were not reported for all studies, possibly due to the p-value or direction of effect being unfavourable to the intervention. Taken together the pooled data from the RCT and cohort studies suggests that the use of a TEG or ROTEM-guided transfusion protocol does not reduce the volume of FFP transfused when compared treatment not guided by TEG or ROTEM (SMD –0.25; 95% CI –0.66, 0.15; p = 0.22; $I^2 = 80\%$).

In patients with diffuse and/or abnormal bleeding from capillary beds and/or excessive blood loss after surgery (see Figure 4.62), data from one small RCT suggested that there was no difference in volume of PLT transfused comparing a ROTEM-guided transfusion protocol with with routine transfusion therapy based on standard laboratory tests (SMD – 0.33; 95% CI –0.94, 0.27; p = 0.28). Data were not reported in 2 studies and 2 studies suggested an effect favouring TEG or ROTEM but did not provide suitable data for analysis.

Among women with severe PPH (see Figure 4.62), data from one observational study suggested that the use TEG is not associated with any reduction in the volume of PLT transfused compared with management of coagulopathy guided by standard laboratory tests (SMD of 0.06; 95% CI –0.32, 0.43; p = 0.76). The other 2 studies did not report this outcome.

Fibrinogen replacement

There was little evidence reported relating to fibrinogen replacement therapy in patients with critical bleeding in whom TEG or ROTEM were used as part of an MHP (see Table 4.84). The evidence was therefore not considered further.

	ГЕМ	Control			:	Std. Mean Difference	Std. Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% Cl	IV, Random, 95% Cl
1.11.3 RCTs									
Weber 2012 (Cardiac) (1)	0	0	50	0	0	50		Not estimable	
Nuttall 2001 (Cardiac) (2)	0	0	41	0	0	51		Not estimable	
Paniagua 2011 (Cardiac) (3)	0.848	1.228	26	1.324	1.624	18	12.3%	-0.33 [-0.94, 0.27]	
Gonzalez 2016 (Trauma)	1.14	0.6	92	0.95	0.72	29	16.5%	0.30 [-0.12, 0.72]	±
Subtotal (95% CI)			118			47	28.8%	0.02 [-0.59, 0.64]	\bullet
Heterogeneity: Tau ² = 0.13; Chi ² = 2	2.84, df =	1 (P =	0.09); l ⁱ	² = 65%					
Test for overall effect: Z = 0.07 (P =	0.94)								
1.11.5 Cohs									
Görlinger 2012a (Coh, trauma) (4)	0	0	0	0	0	0		Not estimable	
Unruh 2019 (Coh, trauma)	1.5	1.5	47	2	0.7	20	13.9%	-0.38 [-0.90, 0.15]	
Barinov 2015 (Coh, PPH)	1.64	1.95	55	1.52	2.15	54	17.6%	0.06 [-0.32, 0.43]	- -
Wang 2017 (Coh, trauma)	0.4	1.5	86	2.9	4.8	80	19.2%	-0.71 [-1.02, -0.40]	
Nardi 2015 (Coh, trauma)	2.7	4.8	96	4.2	9	130	20.4%	-0.20 [-0.46, 0.07]	
Subtotal (95% CI)			284			284	71.2%	-0.31 [-0.64, 0.03]	\bullet
Heterogeneity: Tau ² = 0.08; Chi ² = 1	0.67, df	= 3 (P =	= 0.01);	l² = 72%	6				
Test for overall effect: Z = 1.81 (P =	0.07)								
Total (95% CI)			402			331	100.0%	-0.21 [-0.51, 0.09]	➡
Heterogeneity: Tau ² = 0.10: Chi ² = 1	7.84. df	= 5 (P =	= 0.003): ² = 72	2%			-	
Test for overall effect: Z = 1.37 (P =	0.17)	`		,,					-2 -1 U 1 2
Test for subgroup differences: Chi2:	, = 0.85, d	f = 1 (P	= 0.36), l ² = 0%	6				Favours leg of ROTEINE Favours Control
Footnotes		,	- 1						
(1) Equates TEC or POTEM (p=0.0)	1) Doto i	onortor	l oc mo	dian (IC	ND\. 2 (0	1 2) 10	2 (0 5)		

Figure 4.61 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: PLT transfusion volume (units), by study design.

(1) Favours TEG or ROTEM (p=0.01). Data reported as median (IQR): 2 (0, 2) vs 2 (0, 5).

(2) Favours TEG or ROTEM (p=0.0001). Data reported as median (range): 6 (0, 18) vs 6 (0, 144).

(3) converted from mLs (250 mL/U)

(4) An estimated 72% and 65% reduction in the total volume of PLTs transfused per year after the implementation of goal-directed therapy (two trauma centres).

Figure 4.62 Forest plot of comparison: TEG or ROTEM vs MHP or standard laboratory tests, outcome: PLT transfusion volume (units), by setting.

	TEG	or RO	ЕМ	(Control			Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% Cl
1.12.1 Trauma (RCTs)									
Baksaas-Aasen 2020 (Trauma) (1)	0	0	201	0	0	195		Not estimable	
Gonzalez 2016 (Trauma) Subtotal (95% CI)	1.14	0.6	92 92	0.95	0.72	29 29	16.5% 16.5%	0.30 [-0.12, 0.72] 0.30 [-0.12, 0.72]	
Heterogeneity: Not applicable Test for overall effect: Z = 1.40 (P = 0	0.16)								
1.12.2 Trauma (Coh)									
Görlinger 2012a (Coh, trauma) (2)	0	0	0	0	0	0		Not estimable	
Unruh 2019 (Coh, trauma)	1.5	1.5	47	2	0.7	20	13.9%	-0.38 [-0.90, 0.15]	-
Wang 2017 (Coh, trauma)	0.4	1.5	86	2.9	4.8	80	19.2%	-0.71 [-1.02, -0.40]	_
Nardi 2015 (Coh, trauma) Subtotal (95% CI)	2.7	4.8	96 229	4.2	9	130 230	20.4% 53.6%	-0.20 [-0.46, 0.07] -0.43 [-0.78, -0.08]	•
Heterogeneity: $Tau^2 = 0.06$; $Chi^2 = 5$. Test for overall effect: Z = 2.39 (P = 0	.98, df = 0.02)	2 (P = 0).05); l²	= 67%					
1.12.3 Surgical									
Weber 2012 (Cardiac) (3)	0	0	50	0	0	50		Not estimable	
Nuttall 2001 (Cardiac) (4)	0	0	41	0	0	51		Not estimable	
Paniagua 2011 (Cardiac) (5) Subtotal (95% CI)	0.848	1.228	26 26	1.324	1.624	18 18	12.3% 12.3%	-0.33 [-0.94, 0.27] - 0.33 [-0.94, 0.27]	
Heterogeneity: Not applicable Test for overall effect: Z = 1.08 (P = 0	0.28)								
1.12.5 Obstetrics									
Barinov 2015 (Coh, PPH) Subtotal (95% CI)	1.64	1.95	55 55	1.52	2.15	54 54	17.6% 17.6%	0.06 [-0.32, 0.43] 0.06 [-0.32, 0.43]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.30 (P = 0	0.76)								
Total (95% CI)			402			331	100.0%	-0.21 [-0.51, 0.09]	-
Heterogeneity: Tau ² = 0.10; Chi ² = 1	7.84, df =	= 5 (P =	0.003)	; l² = 72	%			-	
Test for overall effect: Z = 1.37 (P = 0	0.17)								Favours TEG or ROTEM Favours Control
Test for subgroup differences: Chi ² =	8.08, df	= 3 (P :	= 0.04)	, l² = 62	.9%				

Footnotes

(1) Study did not report this outcome.

(2) An estimated 72% and 65% reduction in the total volume of PLTs transfused per year after the implementation of goal-directed therapy (two trauma centres).

(3) Favours TEG or ROTEM (p=0.01). Data reported as median (IQR): 2 (0, 2) vs 2 (0, 5).

(4) Favours TEG or ROTEM (p=0.0001). Data reported as median (range): 6 (0, 18) vs 6 (0, 144).

(5) converted from mLs (250 mL/U)

Table 4.84 Results for TEG or ROTEM to guide BCT versus no TEG or ROTEM to guide BCT: Patients with critical bleeding – PLT, CRYO, PCC transfusion volume

Study ID	Sample size	Patient	Setting	Comparison	Outcome	ne Results			
Study design ^a <i>Risk of bia</i> s	(no. of trials) included in analysis	population	(Location)			TEG/ROTEM Mean ± SD (n)	No TEG/ROTEM Mean ± SD (n)	Risk estimate (95% CI)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Various setting	<u>gs</u>								
Fahrendorff 2017 SR High risk of bias	N = 423 (5 studies) Shore-Lesserson 1999 Wang 2010 Barinov 2015 Gonzalez 2016 Cao 2016	Patients with an acute need for blood components due to bleeding (includes trauma, burns excision, liver transplant, PPH and cardiac surgery)	Various	TEG or ROTEM guided algorithm vs the clinician's discretion and/or based on conventional coagulation tests	PLT transfusion volume, Units	Mean ± SD (246) 0.1 ± 0.276 (53) 27.3 ± 13.9 (14) 1.64 ± 1.95 (55) 1.14 ± 0.6 (92) 2.5 ± 1.3 (32)	Mean ± SD (177) 0.244 ± 0.471 30.1 ± 18.5 (14) 1.52 ± 2.15 (54) 0.95 ± 0.72 (29) 4.2 ± 0.6 (28)	SMD -0.34 (-0.92, 0.24) -0.37 (-0.76, 0.01) -0.17 (-0.91, 0.58) 0.06 (-0.32, 0.43) 0.30 (-0.12, 0.72) -1.62 (-2.21, -1.03)	No significant difference p = 0.25 Substantial heterogeneity l² = 87 (p = 0.00001)
Trauma setting	g								
Bugaev 2020 Coh Serious risk of bias	N = 404 (3 studies) Nardi 2015 Gonzalez 2016 Unruh 2019	Severely injured trauma patients	Trauma (NR)	ROTEM or TEG vs no ROTEM or TEG	PLT transfusion volume, Units	N=199 2.7±4.8 (n=96) 1±1.5 (n=56) 1.5±1.5 (n=47)	N=205 4.2±5.9 (n=130) 1±1.5 (n=55) 2±0.7 (n=20)	MD -0.44 (-1.05, 0.17) MD -1.50 (-2.90, -0.10) MD 0.00 (-0.56, 0.56) MD -0.50 (-1.03, 0.03)	No significant difference p = 0.16 Moderate heterogeneity $l^2 = 55\% (p = 0.11)$
Roullet 2018 RCT High risk of	N = NR (1 RCT) Gonzalez 2016	Patients with severe trauma	Trauma (NR)	TEG or ROTEM guided algorithm vs the clinician's	PLT transfusion volume, Units	Authors note that t more platelets early *See Fahrendorff 2017	he group receiving t y compared to the T for values	he routine tests received EG group.	No significant difference p = NR
bias				discretion and/or based on conventional coagulation tests	Fibrinogen	At 24 hrs, only the a different, being hig	amount of fibrinoger her in the group ma	n administered was naged with routine tests.	No significant difference p = NR
Wang 2017 Coh	N = 166 (1 Coh)	Patients with traumatic liver	SC (US)	TEG guided algorithm vs	PLT transfusion volume, Units	0.4 ± 1.5 (86)	2.9 ± 4.8 (80)	NR	Favours TEG p < 0.01
Serious risk of bias		and/or spleen injury		clinical judgement or usual treatment	CRYO transfusion volume, Units	0.1 ± 0.5 (86)	0.3 ± 1.2 (80)	NR	No significant difference p = NR
Haas 2014 Serious risk of bias	N = 5590 (1 Coh) Görlinger 2012a Görlinger 2012a	Adult patients with severe trauma	SC (Austria, Germany)	ROTEM guided algorithm vs standard of care	PLT transfusion volume, Units	Total per year 25 29	Total per year 90 82	72% reduction 65% reduction	Favours ROTEM p = NR
Surgical settin	g								
Wikkelsø 2016 SR	N = NR (2 RCTs) Weber 2012		SC (Germany, Spain, US)		PLT transfusion volume, Units	Median (IQR) 2 (0, 2) Median (range)	Median (IQR) 2 (0, 5) Median (range)		Favours TEG or ROTEM p = 0.010

Study ID	Sample size (no. of trials)	Patient population	Setting (Location)	Comparison	Outcome	Results			
Study						TEG/ROTEM	No TEG/ROTEM	Risk estimate	Statistical significance
design "	analysis					Mean ± SD (n)	Mean ± SD (n)	(95% CI)	<i>p</i> -value
RISK OF DIAS	,								Heterogeneity ^b
High risk of	Nuttall 2001	Adults or children		TEG or ROTEM		6 (0, 18)	6 (0, 144)		<i>p</i> = 0.0001
bias	N = NR1RCTs)	with diffuse or		guided	PLT transfusion	Mean ± SD	Mean ± SD	SMD ^c	
		excessive bleeding		transfusion vs	volume, mL				
	Paniangua 2011	after cardiac		clinical		212 ± 307 (26)	331 ± 406 (18)	-0.33 (-0.94, 0.27)	p = NR
		surgery, orthotopic		judgement or					
		liver transplant or		usual treatment					
		Durns excision		with of without					
				algorithm to					
				guide SLTs					
Haas 2014	N = 100 (1 RCT)	Adult patients	SC	ROTEM guided	PLT transfusion	Median (IQR)	Median (IQR)		Favours ROTEM
RCT	Weber 2012	undergoing cardiac	(NR)	algorithm vs	volume, units	2 (0, 2)	2 (0, 5)	ND	
High risk of		or aortic surgery		standard of care					<i>p</i> = 0.01
bias	N = 5338 (1 Coh)	Adult patients	SC	ROTEM guided	PLT transfusion	Total per year	Total per year		
	Görlinger 2012b	undergoing visceral (Germany) surgery or liver transplant	(Germany)	Germany) algorithm vs standard of care	volume, mL	149	433	66% reduction	p = NR
					FC required, g	per year	per year		Favours SoC
						745	68	9.9-fold increase	p = NR
					PCC required, IU	per year	per year		Favours SoC
						238 500	65 500	2.6-fold increase	<i>p</i> = NR

Studies with strikethrough do not meet the PICO criteria for this question.

Abbreviations: CI, confidence interval; Coh, cohort study; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; hrs, hours; IQR, interquartile range; MC, multicentre; MD, mean difference; n, number; NR, not reported; PCC, prothrombin complex concentrate; PLT, platelets; PPH, postpartum haemorrhage; RBC, red blood cells; RCT, randomised controlled trial; ROTEM, rotational thromboelastometry; SC, single centre; SD, standard deviation; SMD, standard mean difference; SoC, standard of care; TEG, thromboelastography; UK, United Kingdom; US, United States

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H random effects.

4.9.3.4 Time to transfusion

No evidence found.

4.9.3.5 Dose/type of transfusion

There was little evidence reported relating to the dose or type of transfusion (e.g. more than 5 units transfused) in patients with critical bleeding in whom TEG or ROTEM were used as part of an MHP. The evidence was therefore not considered further. The available data is provided in volume 3 of the Technical report.

4.10 Cell salvage (Question 9)

Question 9 - (interventional)

In patients with critical bleeding what is the effect of cell salvage on patient outcomes?

4.10.1 Methods

This question examined the effect and cost of cell salvage compared with no cell salvage in patients *with* critical bleeding (i.e. major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion) as outlined in Figure 4.63.

The question focused on individuals in an urgent or emergency setting, including those admitted to a trauma, maternity or perioperative setting. No age limits were applied, however studies in neonates (newborns up to 28 days) and studies in individuals with hereditary bleeding disorders were not eligible for inclusion.

Figure 4.63 PICO criteria: Question 9 - cell salvage



Abbreviations: ARDS, acute respiratory distress syndrome; CRYO, cryoprecipitate; FC, fibrinogen concentrate; FFP, fresh frozen plasma; INR, international normalised ratio; PLT, platelets; RBC, red blood cells; ROTEM, thromboelastometry; TE, thromboembolic event; TEG, thromboelastography

a. Adult (aged over 18 years), child (aged 2 to 12 years), adolescent (aged 13 to 18 years), infants (aged 1 to 23 months).

b. e.g. trauma, obstetric, perioperative (cardiothoracic, general surgery, gastrointestinal, liver transplant), paediatric, other. c. Newborns up to 28 days following birth.

Selection of studies was conducted according to the screening criteria described in Section 3.3.

The initial 2018 search was limited to studies published after 1990, noting primary studies published prior to 1990 that had been included within a systematic review were also eligible for inclusion. There were no restrictions applied to sample size.

Assuming relevant primary studies had been identified in the included systematic review and meta-analyses, the systematic screening of RCTs and nonrandomised studies was limited to studies published after 2015, based on the literature search date of the most recent and comprehensive identified systematic review (Meybohm 2016).

The literature search was updated in August 2019 and again in September 2021 to identify any new studies meeting the eligibility criteria. In this updated search, systematic reviews were first screened, then RCTs were screened. No new systematic reviews or RCTs were found.

Studies conducted in patients receiving intraoperative cell salvage in the elective setting, such as those scheduled for radical prostatectomy or other cancer-related surgery, total hip or knee arthroplasty, scoliosis surgery, minimally invasive cardiothoracic surgery, infrarenal abdominal aortic aneurysm surgery, craniosynostosis and caesarean section, were excluded. These studies were deemed more appropriate for assessment in the perioperative module.

4.10.2 Summary of evidence

4.10.2.1 Systematic reviews

Three systematic reviews (243-245) were identified that assessed the effects of cell salvage compared with no cell salvage in patients with critical bleeding (Nayar 2017, Meybohm 2016, Shantikumar 2011). The main characteristics and quality of these reviews and relevant outcomes assessed are summarised in Table 4.85. An overlap table listing the potentially relevant primary studies included in the reviews is provided in Table 4.86.

One other systematic review (Li 2015) (246) was identified in the literature search that did not provide any additional data than that of the Meybohm 2016, thus was not considered further in this review (duplicate data). A list of studies that met the PICO criteria for this question but were later excluded is provided in **Appendix B** (technical report, volume 2).

Nayar 2017 was a narrative review that assessed blood conservation strategies in the setting of acute orthopaedic trauma. The review authors noted 7 primary studies in their discussion of cell salvage that were retrieved for further assessment; but later deemed more appropriate for assessment in the perioperative module as patients were not critically bleeding.

Meybohm 2016 was a systematic review and meta-analysis of RCTs involving patients scheduled for all types of surgery randomised to washed cell salvage or no cell salvage. The primary outcome of interest was the number of patients exposed to RBC transfusion, with the volume of blood transfused, mortality and rates of infection also assessed. The authors identified 47 studies that met their inclusion criteria, one of which involved patients undergoing surgery for multiple trauma (penetrating abdominal trauma) and was considered relevant to this review (Bowley 2006).

Shantikumar 2011 was a systematic review of all available evidence relating to the use of cell salvage in abdominal aortic aneurysm (AAA) repair. Where possible, a meta-analysis of relevant data was performed, with a focus on the proportion of patients transfused, the volume of blood component used, complications and length of ICU and hospital stay. The author identified 23 studies that met their inclusion criteria, with 5 non-randomised studies involving ruptured AAA repair considered relevant to this review (Markovic 2009, Tawfick 2008, Serracino-Inglott 2005, Shuhaiber 2003, Posacioglu 2002).

Review ID Study design Review quality		Population	Intervention	Comparator	Outcomes
Surgical setting					
Nayar 2017 (243) Critically low	Narrative review	Acute orthopaedic trauma (61 studies) ^a	Cell salvage	Any	Transfusion volume Cost
Meybohm 2016 (245) <i>High</i>	SR / MA of RCTs	Patients scheduled for any type of surgery (47 studies) ^b	Washed cell salvage	No cell salvage	Mortality Morbidity Transfusion volume Costs
Shantikumar 2011 (244) <i>Critically low</i>	SR of RCTs and cohort studies	Abdominal aortic aneurysm surgery (23 studies) ^c	Washed cell salvage	No cell salvage	Mortality Morbidity Transfusion volume

Table 4.85 Characteristics and quality of SR and MA evidence: cell salvage versus no cell salvage

Abbreviations: MA, meta-analysis; RCT, randomised controlled trial; SR, systematic review

a. The authors found no studies considered relevant to this review. There were 6 studies in orthopaedic trauma; but none were in patients *with* critical bleeding.

b. The authors included one study in multiple trauma surgery (Bowley 2006) considered relevant to this review. A further 15 studies in orthopaedic surgery, 21 in cardiac surgery, 6 in vascular surgery, 2 in cancer surgery and 2 in paediatric surgery were not relevant to this review.

c. The authors included 5 studies in urgent rupture repair considered relevant to this review. A further 18 studies (9 uncontrolled studies, 5 non-randomised trials and 4 RCTs) involved elective surgery and therefore did not meet the PICO criteria for this review.

Table 4.86 Overlap table showing systematic reviews and included primary studies: cell salvage versus no cell salvage

Review ID		Nayar 2017	Meybohm 2016	Shantikumar 2011			
Trauma							
ۍ و	Bhangu 2013 ª						
Stu	Bowley 2006		*				
Orth	opaedic trauma						
	Firoozabadi 2015	Х					
	Bigsby 2013	Х					
₽	Canan 2013	Х					
ð	Odak 2013	Х					
Stu	Scannell 2009	Х					
	Cavalieri 1994	X					
	Schmidt 1998	Х					
Rupt	ured aneurysm repair						
	Markovic 2009			✓			
₽	Tawfick 2008			✓			
Study	Serracino-Inglott 2005			✓			
	Shuhaiber 2003			✓			
	Posacioglu 2002			✓			

a. Study identified and excluded by Li 2015 (not randomised).

X = study does not meet PICO criteria (participants not critically bleeding).

 \checkmark = study meets PICO criteria (participants with major haemorrhage that is life-threatening and is likely to result in the need for massive transfusion).

4.10.2.2 Randomised controlled trials

One small RCT (Bowley 2006) examining the effect of cell salvage in patients with critical bleeding was identified in the included systematic reviews. No additional RCTs were identified through the systematic review and handsearching process. A summary of the characteristics and quality of the identified RCT is provided in Table 4.87.

Bowley 2006 enrolled adult patients (aged over 18 years) presenting to emergency with penetrating torso injury requiring laparotomy and had exhibited hypotension (< 90 mm Hg) either prehospital or on arrival and in whom there was significant blood loss. The study was conducted in South Africa (within the Johannesburg Hospital Trauma Unit), and patients were predominantly male (40/44, 91%).

Table 4.87 Characteristics and quality of RCT evidence: cell salvage versus no cell salvage

Study ID <i>Risk of bias</i>	Study design	Population	Intervention	Comparator	Outcomes		
Trauma	Trauma						
Bowley 2006 (247) <i>High</i>	RCT	Adults with penetrating abdominal trauma N = 44	Cell salvage (n=21)	No cell salvage ª (n=21)	Mortality Transfusion volume		

Abbreviations: RCT, randomised controlled trial

a. Donor blood transfusion at the discretion of the attending medical staff

4.10.2.3 Observational and cohort studies

Five nonrandomised studies (Markovic 2009, Tawfick 2008, Serracino-Inglott 2005, Shuhaiber 2003, Posacioglu 2002) involving urgent AAA repair were identified in the included systematic review and were considered relevant to this review. One additional cohort study (Bhangu 2013) was identified through the systematic review and handsearching process that examined the effect of cell salvage in patients with critical bleeding. A summary of the characteristics and quality of the identified nonrandomised studies is provided in Table 4.88.

Trauma

Bhangu 2013 was a prospective cohort study conduct in Afghanistan among patients admitted to a combat support hospital with battle-related injury. Out of 130 patients (76% blast-injury, 22% gunshot, 2% road), 29 patients were judged by the attending military surgeon to likely require at least 10 units of RBC in the first 12 hours after injury. Eighteen cases were selected for intraoperative blood salvage, which was successfully completed in 17 patients (one died on operating table before cell salvage could occur). The control group included 11 patients who were admitted at the same time and received at least 10 units of RBC in the first 12 hours after injury but did not undergo cell salvage.

The study had important problems relating to insufficient information about potential confounders. It was also noted that blast injuries (predominantly from improvised explosive devices) drive environmental material deep into patients' wounds, leading to gross contamination. The study was therefore judged not applicable to the Australian context and was not included in the GRADE evidence summary tables or when developing recommendations.

Surgical setting

Markovic 2009 retrospectively reviewed clinical and financial outcomes relating to abdominal aortic surgery among 90 patients who received intraoperative cell salvage compared with 90 patients who did not receive intraoperative cell salvage at a single institution in Serbia. The patients were subdivided according to the type of operation, being aortoiliac occlusive disease (AOD), elective AAA repair or ruptured AAA repair. Only the ruptured AAA repair was relevant to this review.

Tawfick 2008 retrospectively reviewed ruptured AAA over a 9-year period (between June 1997 and June 2006) at a single hospital in Ireland. The study included both emergency open AAA repair and scheduled or elective AAA repair³⁹. The mean age for all patients who received cell salvage was 72 years, which was significantly higher (p = 0.01) than that of the control group (69 years). All other factors (preoperative cardiac, pulmonary and renal status, smoking, diabetes, mean preoperative haemoglobin) were comparable between groups.

Serracino-Inglott 2005 was a prospective cohort study that examined 154 ruptured AAA repairs reported to a regional vascular audit database in the UK over a 4-year period (January 2000 to June 2004). The 2 groups were matched for age, cardiac and respiratory symptoms, cardiac medication, incidence of myocardial infarction and diabetes.

Shuhaiber 2003 was a small retrospective cohort study conducted at a single centre in the UK among 128 patients who underwent AAA repair between 1992 and 1999 by a single vascular surgeon. Only 25 patients had emergency AAA repair (Group B), with the other 93 patients receiving elective AAA repair (Group A)³⁹. Among patients in Group B, the mean age was 74.3 years (range 58 to 84), all but 2 patients were male (23/25; 92%).

Posacioglu 2002 retrospectively reviewed mortality, post-operative morbidity and blood loss in 56 patients with suprarenal and infrarenal ruptured AAA repairs by a single surgeon in Turkey. There were no differences in baseline characteristics (98% [55/56] were male), with the mean age being 68 ± 8 years.

None of the above studies were randomised, due to the unpredictability and urgency of admissions and difficulties with ethical approval. All studies had important problems relating to patient selection bias, and outcome assessment and reporting bias.

Study ID Risk of bias	Study design	Population	Intervention	Comparator	Outcomes
Trauma					
Bhangu 2012 (248) <i>Serious</i>	Prospective cohort	Patients with sustained combat- related injury requiring massive transfusion ^a	Cell salvage (n=18)	No cell salvage (n=11)	Mortality Transfusion volume

Table 4.88 Characteristics and quality of observation and cohort studies: cell salvage versus no cell salvage

³⁹ Not relevant to this review.

Study ID Risk of bias	Study design	Population	Intervention	Comparator	Outcomes			
Surgical setting								
Markovic 2009 (249) <i>Serious</i>	Prospective cohort with historical controls	Ruptured AAA repair	Cell salvage (n=15)	No cell salvage (n=15)	Mortality Morbidity Transfusion volume			
Tawfick 2008 (250) <i>Serious</i>	Retrospective cohort	Emergency open AAA repair and scheduled or elective AAA repair	Cell salvage (n=27)	No cell salvage (n=28)	Mortality Morbidity Transfusion volume			
Serrancino- Inglott 2005 (251) <i>Serious</i>	Prospective cohort (regional audit database)	Ruptured AAA repairs over a 4- year period (January 2000 to June 2004)	Cell salvage (n=40)	No cell salvage (n=114)	Mortality Morbidity Transfusion volume			
Shuhaiber 2003 (252) <i>Serious</i>	Retrospective cohort	AAA surgery by a single surgeon	Cell salvage (n=4)	No cell salvage (n=21)	Mortality Morbidity Transfusion volume			
Posacioglu 2002 (253) <i>Serious</i>	Retrospective cohort	Suprarenal and infrarenal ruptured AAA repairs by a single surgeon	Cell salvage (n=40)	No cell salvage (n=16)	Mortality Morbidity Transfusion volume			

Abbreviations: AAA, abdominal aortic aneurysm a. Requiring at least 10 units of RBC in the first 12 hours after injury.
4.10.3 Results

4.10.3.1 Mortality

A summary of the evidence relating to mortality (at any timepoint up to 30 days) in patients with critical bleeding receiving intraoperative autologous transfusions (obtained by cell salvage) is presented in Table 4.89. None of the individual studies were powered to detect differences in mortality.

Overall, no difference in mortality comparing patients who received cell salvage with those who did not (regardless of clinical setting) was observed but the evidence is very uncertain. For most bleeding patients there is no substantial survival benefit associated with cell salvage.

Pooled data from the identified RCT and nonrandomised trials (see Figure 4.64) showed the mortality rate in patients with critical bleeding to be lower among those who received cell salvage (62/180, 34%) compared with those who did not (102/243, 42%). The difference was not significant (RR 0.83; 95% CI 0.65, 1.06; p = 0.14, $l^2 = 0$ %).

In trauma patients, there were 15 deaths among the 39 patients (38.5%) who received cell salvage compared with 15 deaths among the 34 (44%) patients who received standard care. The results suggest no difference between groups for the outcome of mortality (RR 1.03; 95% CI 0.68, 1.57; p = 0.88; I² = 0%). In considering the RCT evidence alone, the mortality rate was higher (66.7% vs 65.2%), with no difference between groups observed (RR 1.02; 95% CI 0.67, 1.56; p = 0.92)(*GRADE: very low*).

Among patients requiring urgent abdominal aortic aneurysm repair, there were fewer deaths among those who received cell salvage (47/141, 33%) compared with those who did not (87/209, 42%). An effect favouring cell salvage is suggested (RR 0.74; 95% CI 0.55, 1.01; p = 0.05; I² = 0%) (*GRADE: very low*). There were concerns of nonreporting bias for this outcome with some studies excluding patients who died in the theatre and other reporting combined mortality data (across treatment groups).

Figure 4.64 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Mortality, any timepoint up to 30 days

	Experim	ental	Contr	ntrol Risk Ratio			Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl		
1.1.1 Trauma									
Bhangu 2012 (coh, combat trauma) (1)	1	18	0	11	0.6%	1.89 [0.08, 42.82]	· · · · · · · · · · · · · · · · · · ·		
Bowley 2006 (RCT, penetrating trauma) (2) Subtotal (95% CI)	14	21 39	15	23 34	33.1% 33.7%	1.02 [0.67, 1.56] 1.03 [0.68, 1.57]			
Total events	15		15						
Heterogeneity: Tau ² = 0.00; Chi ² = 0.16, df = 1	(P = 0.69);	l ² = 0%							
Test for overall effect: Z = 0.15 (P = 0.88)									
1.1.2 Urgent abdominal aortic aneurysm rep	air								
Shuhaiber 2003 (Coh. urgent AAA) (3)	0	4	0	21		Not estimable			
Tawfick 2008 (Coh. urgent AAA) (4)	6	27	9	28	7.6%	0.69 [0.28, 1.68]			
Posacioglu 2002 (Coh. urgent AAA) (5)	16	40	8	16	15.6%	0.80 [0.43, 1.49]			
Markovic 2009 (Coh, ruptured AAA) (6)	12	30	14	30	17.7%	0.86 [0.48, 1.53]			
Serracino-Inglott 2005 (Coh. urgent AAA) (7)	13	40	56	114	25.5%	0.66 [0.41, 1.07]	_ _		
Subtotal (95% CI)		141		209	66.3%	0.74 [0.55, 1.01]	\bullet		
Total events	47		87						
Heterogeneity: Tau ² = 0.00; Chi ² = 0.54, df = 3	(P = 0.91);	l ² = 0%							
Test for overall effect: Z = 1.92 (P = 0.05)	(,								
Total (95% CI)		180		243	100.0%	0.83 [0.65, 1.06]	•		
Total events	62		102						
Heterogeneity: $Tau^2 = 0.00$: $Chi^2 = 2.35$ df = 5	(P = 0.80)	$ ^2 = 0\%$							
Test for overall effect: $7 = 1.48$ (P = 0.14)							0.2 0.5 1 2 5		
Test for subgroup differences: $Chi^2 = 1.54$ df =	1 (P - 0.2	1) 2 - 3	5.2%				Favours cell salvage Favours no cell salvage		
	(1 - 0.2	·/,· = 0	0.270						

Footnotes

(1) One patient in the intervention group died before cell salvage could occur.

(2) Cause of death: I = exsanguination (8/14) or MOF related to sepsis (6/14). C = exsanguination (10/15) and MOF related to sepsis (5/15).

(3) Ten out of 25 (40%) patients in the total study cohort died (intra- and post-operative). A further 5 patients in the control group died up to 30-days.

(4) Data retrieved from primary study. 30-day mortality

(5) Date retrieved from primary study. Includes post-opererative deaths only.

(6) Data retrieved from primary study. Includes intro-operative and post-operative deaths among patients with ruptured AAA.

(7) Data retrieved from primary study. Includes intro-operative and post-operative deaths.

Table 4.89 Results for cell salvage versus no cell salvage: Patients with critical bleeding – Mortality

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			Cell salvage n/N (%)	No cell salvage n/N (%)	Risk estimate (95% CI)	Statistical significance p-value Heterogeneity ^b
Trauma setting	1								
Meybohm 2016 RCT High risk of bias	N = 44 (1 RCT) Bowley 2006	Patients <i>with</i> penetrating torso injury requiring laparotomy and hypotension < 90 mm Hg	SC, trauma (South Africa)	Cell salvage vs no cell salvage	Mortality, timing not specified °	14/21 (66.7)	15/23 (65.2)	RR 1.02 (0.67, 1.56)	No significant difference p = 0.92 Heterogeneity NA
Bhangu 2012 Coh High risk of bias	N= 130 (1 Coh) Banghu 2012	Patients with sustained combat- related injury requiring massive transfusion ^d	SC, combat (Afghanistan)	Cell salvage vs no cell salvage	Mortality, timing not specified ^e	1/18	0/11	Not estimable	Not estimable
Surgical setting	1	-	-	-		-	-	-	-
Shantikumar 2011 Coh <i>Serious risk of</i> <i>bias</i>	N= 360 (5 Coh) Markovic 2009 Posacioglu 2002 Tawfick 2008 Serrancino-Inglott 2005 ^r Shuhaiber 2003 ^g	Patients undergoing emergency AAA repair	SC, vascular surgery (Ireland, Serbia, Turkey, UK)	Cell salvage vs no cell salvage	Mortality, any timepoint up to 30 days	47/137 (34.3) 12/30 (40) 16/40 (40) 6/27 (22) NR/40 (32) NR/4	87/188 (46.28) 14/30 (46.6) 8/16 (50) 9/28 (32) NR/114 (49) NR/21	RR 0.74 (0.55, 1.01) ^h	No significant difference p = 0.05 Moderate heterogeneity $I^2 = 0\%$ ($p = 0.91$)

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Cause of death was exsanguination (10/15) and MOF related to sepsis (5/15) in the control group and exsanguination (8/14) and MOF related to sepsis (6/14) in the intervention group.

d. Requiring at least 10 units of RBC in the first 12 hours after injury.

e. One patient in the intervention group died on operating table before cell salvage could occur.

f. Serracino-Inglott 2005 excluded patients who died in the theatre from the analysis and reported an effect favouring cell salvage (21% vs 44%; p = 0.01).

g. There were only 4 patients in the intervention group, therefore no meaningful difference in mortality between groups could be observed. Overall, 10/25 (40%) patients in the study cohort died (intraoperative and post-operative. A further 5 patients in the control group died within 30-days.

h. Calculated post-hoc using RevMan 5.4. M-H Random effects.

4.10.3.2 Morbidity

A summary of the evidence relating to morbidity in patients with critical bleeding receiving intraoperative autologous transfusions (obtained by cell salvage) is presented in Table 4.90.

Post-operative complications

The identified systematic reviews reported no significant difference in any post-operative complications between patients who received cell salvage compared with those who did not, regardless of clinical setting. For most bleeding patients there are no clear substantial harms associated with cell salvage, but the evidence is very uncertain (GRADE: very low).

Data from the identified RCT (see Figure 4.65), suggested that in patients with penetrating trauma, the risk of sepsis was comparable between those who received cell salvage and those who did not (RR 0.78; 95% CI 0.29, 2.09; p = 0.62).

In the surgical setting, patients requiring elective and urgent abdominal aortic aneurysm repair who had cell salvage were also no more likely to have respiratory complications, renal or gastrointestinal complications, than those who received standard care.

Not including the studies that reporting combined data for elective and urgent abdominal aortic aneurysm repair (see Figure 4.66), the risk of post-operative respiratory complications was higher among patients who received cell salvage (16/84, 19%) compared with those who did not (2/151, 1.3%); but the difference did not reach statistical significance (RR 3.20, 95% CI 0.83, 12.35; p = 0.09) (*GRADE: very low*).

Similar data were observed for post-operative renal complications (12% vs 1.3%; RR 2.00, 95% CI 00.49, 8.14; p = 0.33) (GRADE: very low) and post-operative gastrointestinal complications (4.8% vs 0.7%; RR 1.60, 95% CI 0.19, 13.24; p = 0.66) (GRADE: very low).

Need for re-operation

Among patients requiring elective and urgent abdominal aortic aneurysm repair, reoperation was needed in 8% (14/174) of patients who received cell salvage compared with 7% (17/241) in those who did not (RR 0.64, 95% CI 0.32, 1.30; p = 0.35; I²=0%).

Not including the studies that reporting combined data for elective and urgent abdominal aortic aneurysm repair (see Figure 4.66), the risk for re-operation was higher among patients who received cell salvage (6/84, 7%) compared with those who did not (2/151, 1.3%), but the difference was not significant (RR 1.20, 95% CI 0.27, 5.33; p = 0.81) (*GRADE: very low*).

Figure 4.65 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Morbidity - post-operative complications

	Cell salv	age	No cell sa	lvage		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
1.2.1 Sepsis							
Bowley 2006 (RCT, penetrating trauma) Subtotal (95% CI)	5	21 21	7	23 23	100.0% 1 00.0 %	0.78 [0.29, 2.09] 0.78 [0.29, 2.09]	
Total events	5		7				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.49$ (P = 0.62)							
1.2.2 Respiratory complications							
Shuhaiber 2003 (Coh, urgent AAA) (1)	0	4	0	21		Not estimable	
Serracino-Inglott 2005 (Con, urgent AAA)	0	40	0	114	00.00/	Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	16	40	2	16	26.8%	3.20 [0.83, 12.35]	
Markovic 2009 (Con, ruptured AAA) (2)	5	90	1	90	32.9%	0.71 [0.24, 2.17]	
Subtotal (95% CI)	8	101 275	11	86 327	40.3% 100.0%	0.62 [0.26, 1.47] 1.01 [0.40, 2.56]	
Total events	29		20				
Heterogeneity: Tau ² = 0.37; Chi ² = 4.36, df	= 2 (P = 0.1	1); l² =	54%				
Test for overall effect: Z = 0.02 (P = 0.99)							
1.2.3 Renal complications							
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Shuhaiber 2003 (Coh, urgent AAA) (4)	0	4	0	21		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	10	40	2	16	30.9%	2.00 [0.49, 8.14]	
Tawfick 2008 (Coh, urgent AAA) (5)	3	101	9	86	34.0%	0.28 [0.08, 1.02]	
Markovic 2009 (Coh, ruptured AAA) (6)	4	90	6	90	35.1%	0.67 [0.19, 2.28]	
Subtotal (95% CI)		275		327	100.0%	0.70 [0.24, 2.05]	\bullet
Total events	17		17				
Heterogeneity: Tau ² = 0.46; Chi ² = 4.08, df	= 2 (P = 0.1	3); l² =	51%				
Test for overall effect: Z = 0.65 (P = 0.51)							
1.2.4 Gastrointestinal complications							
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Shuhaiber 2003 (Coh, urgent AAA) (7)	0	4	0	21		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	4	40	1	16	36.8%	1.60 [0.19, 13.24]	
Markovic 2009 (Coh, ruptured AAA) (8)	5	90	2	90	63.2%	2.50 [0.50, 12.55]	
Subtotal (95% CI)		174		241	100.0%	2.12 [0.59, 7.65]	-
Total events	9		3				
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 0.11$, df Test for overall effect: Z = 1.15 (P = 0.25)	= 1 (P = 0.7	′4); l² =	0%				
1.2.5 Need for reoperation							
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Shuhaiber 2003 (Coh, urgent AAA) (9)	0	4	0	21		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	6	40	2	16	22.6%	1.20 [0.27, 5.33]	
Markovic 2009 (Coh, ruptured AAA) (10) Subtotal (95% CI)	8	90 174	15	90 241	77.4% 100.0%	0.53 [0.24, 1.20] 0.64 [0.32, 1.30]	
Total events	14		17				
Heterogeneity: Tau ² = 0.00; Chi ² = 0.88, df Test for overall effect: Z = 1.23 (P = 0.22)	= 1 (P = 0.3	85); l² =	0%				
							0.005 0.1 1 10 200
							Favours cell salvage Favours no cell salvage

Footnotes

(1) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

- (2) Data includes urgent and elective AAA and AOD. Separate data for ruptured AAA not available.
- (3) Data includes elective and emergent AAA repair. Seperate data for emergency AAA not available.
- (4) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(5) Need for dialysis. Data includes elective and emergent AAA repair. Seperate data for emergency AAA not available.

(6) Data includes urgent and elective AAA and AOD. Separate data for ruptured AAA not available.

(7) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(8) Data includes urgent and elective AAA and AOD. Separate data for ruptured AAA not available.

(9) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(10) Data includes urgent and elective AAA and AOD. Separate data for ruptured AAA not available.

Figure 4.66 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Morbidity - post-operative complications (urgent AAA repair)

	Cell salv	/age	No cell salvage		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% Cl
1.3.2 Respiratory complications							
Shuhaiber 2003 (Coh, urgent AAA) (1)	0	4	0	21		Not estimable	
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	16	40	2	16	100.0%	3.20 [0.83, 12.35]	+- -
Subtotal (95% CI)		84		151	100.0%	3.20 [0.83, 12.35]	
Total events	16		2				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.69 (P = 0.09)							
1.3.3 Renal complications							
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Shuhaiber 2003 (Coh, urgent AAA) (2)	0	4	0	21		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	10	40	2	16	100.0%	2.00 [0.49, 8.14]	
Subtotal (95% CI)		84		151	100.0%	2.00 [0.49, 8.14]	
Total events	10		2				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.97 (P = 0.33)							
1.3.4 Gastrointestinal complications							
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Shuhaiber 2003 (Coh, urgent AAA) (3)	0	4	0	21		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	4	40	1	16	100.0%	1.60 [0.19, 13.24]	
Subtotal (95% CI)		84		151	100.0%	1.60 [0.19, 13.24]	
Total events	4		1				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.44 (P = 0.66)							
1.3.5 Need for reoperation							
Shuhaiber 2003 (Coh, urgent AAA) (4)	0	4	0	21		Not estimable	
Serracino-Inglott 2005 (Coh, urgent AAA)	0	40	0	114		Not estimable	
Posacioglu 2002 (Coh, urgent AAA)	6	40	2	16	100.0%	1.20 [0.27, 5.33]	_
Subtotal (95% CI)		84		151	100.0%	1.20 [0.27, 5.33]	\bullet
Total events	6		2				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.24 (P = 0.81)							
							Favours cell salvage Favours no cell salvage

Footnotes

(1) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(2) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(3) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

(4) Small intervention group prevents meaningful comparison between groups. Overall, 14/25 (56%) patients had major complications.

Table 4.90 Results for cell salvage versus no cell salvage: Patients with critical bleeding – Morbidity: post-operative complications

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results				
Study design ^a	(no. of trials) included in analysis	population	(Location)			Cell salvage n/N (%)	No cell salvage n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b	
Trauma setting						- -		- -		
Meybohm 2016 RCT High risk of blas	N = 44 (1 trial) Bowley 2006	Patients with penetrating torso injury requiring laparotomy and hypotension < 90 mm Hg	SC, emergency (South Africa)	Cell salvage vs no cell salvage	Infection (sepsis)	5/21 (23.8)	7/23 (30.4)	RR 0.78 (0.29, 2.09)	No significant difference <i>p</i> = 0.62	
Surgical setting										
Shantikumar 2011 Coh	N = 360 (5 Coh) Markovic 2009		SC, vascular surgery	Cell salvage vs no cell salvage	Respiratory Posacioglu 2002	16/40 (40)	2/16 (12.5)	RR 3.20 (0.83, 12.35) ^c	No significant difference p = 0.09 ^d	
Serious risk of bias	Tawfick 2008 Serrancino-Inglott		(Ireland, Serbia, Turkey, UK)		Renal Posacioglu 2002	10/40 (25)	2/16 (12.5)	RR 2.00 (0.49, 8.14) ^c	<i>No significant difference</i> <i>p</i> = 0.475	
	2005 Shuhaiber 2003				Gastrointestinal Posacioglu 2002	4/40 (10)	1/16 (6.25)	RR 1.60 (0.19, 13.24) ^c	No significant difference p = 1.00	
					Need for re- operation Posacioglu 2002	6/40 (15)	2/16 (12.5)	RR 1.20 (0.27, 5.33) ^c	No significant difference p = 0.588	
					Any post-operative complication Serracino-Inglott 2005	The authors noted n groups for post-ope data provided.	o significant differe rative complication	ence between study s, but no further	No significant difference	
				Shuhaiber 2003	No meaningful diffe observed (only 4 pat Overall, 14/25 (56%) pati - haemorrhage and - infection, - non-graft thromb - arrythmia, cardiar respiratory failure	rence in complicati cients in the interve- ents had major compli d anastomotic leak, posis, embolism, myoca c failure, impaired rena	ons could be ntion group). cations including: ardial infarction, I function and	No significant difference		
					Marcovic 2009	Data were presenter AAA and AOD and a The authors noted no si - Transfusion-relate - Multiorgan failure failure - Stroke, Myocardia - Wound infection,	d for entire Coh tha re therefore not pre ignificant difference be ed complications e, Colon ischaemia, Res al infarction Bleeding or Re-operat	t includes elective esented here. etween study groups for: piratory failure, Renal tion	No significant difference	

Study ID	Sample size	e size Patient Setting	Setting (Comparison	Outcome	Results			
Study design ^a (n in ar	included in analysis	population	(Location)		Tawfick 2009	Cell salvage n/N (%)	No cell salvage n/N (%)	Risk estimate (95% Cl)	Statistical significance p-value Heterogeneity ^b
					Tawfick 2008	Data were presented elective and emerge presented here. The authors noted no si - Respiratory comp - Cardiac complica A significant effect favo - Need for renal dia	d for entire study of ency AAA and are the ignificant difference be ilications (ARDS, pneu- tions (arrythmias, ischauring no cell salvage o alysis ($p = 0.037$).	ohort that includes herefore not etween study groups for: monia, atelectasis) aemic cardiac event) bserved for:	
Obstetrics and m	aternity setting - I	no comparative evid	ence found						
Paediatric setting	g – no comparative	e evidence found							

Abbreviations: AAA, abdominal aortic aneurysm; CI, confidence interval; M-H, Mantzel-Hentzel; NR, not reported; RR, relative risk; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses Observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{het} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H. Random effects.

d. Posacioglu 2002 reported a statistically significant effect favouring no cell salvage p = 0.047

4.10.3.3 Transfusion volume

A summary of the evidence relating to transfusion volumes in patients with critical bleeding receiving intraoperative autologous transfusions (obtained by cell salvage) is presented in Table 4.91.

Overall, only limited conclusions can be drawn from the available evidence. For most bleeding patients there is a modest reduction in the volume of RBC transfused (between 2 and 5 red cell units saved), but the evidence is very uncertain (*GRADE: very low*).

Red blood cells

A meta-analysis of data from RCTs and cohort studies included in this review revealed a significant reduction in the volume of RBC transfused in patients with critical bleeding who received cell salvage (n=162) compared with those who did not (n=232), with an overall standardised mean difference (SMD) of –0.45 units (95% CI –0.87, –0.01; p = 0.05; random effects, I² = 71%).

In patients with penetrating trauma, evidence from the small RCT suggests a significant reduction in the volume of RBC transfused (around 4.7 red cell units saved) favouring cell salvage (SMD –0.82; 95% CI –1.44, –0.20; p = 0.009) (*GRADE: very low*).

Among patients requiring urgent AAA repair, the volume of RBC transfused was not significantly different between groups (SMD –0.36; 95% CI –0.87, –0.14; p = 0.16) (GRADE: very low).

Other blood components

In patients with penetrating trauma, evidence from the small RCT comparing cell salvage with standard care suggests no difference in the the volume of FFP (SMD 0.16; 95% CI – 0.44, 0.75; p = 0.61) or PLT transfused (SMD 0.26; 95% CI –0.33, 0.85; p = 0.39) (GRADE: very *low*).

Among patients requiring urgent AAA repair, there was no difference between groups in the the volume of FFP transfused (SMD 0.21; 95% CI –0.97, 1.40; p = 0.72) (GRADE: very low). There was no data relating to the volume of PLT transfused (if any).

Figure 4.67 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (RBC)

	Experimental Control			Control		:	Std. Mean Difference	Std. Mean Difference			
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
1.4.1 Trauma (units)											
Bowley 2006 (RCT, penetrating trauma) Subtotal (95% CI)	6.47	5.14	21 21	11.17	6.06	23 23	16.4% 16.4%	-0.82 [-1.44, -0.20] -0.82 [-1.44, -0.20]	•		
Heterogeneity: Not applicable											
Test for overall effect: Z = 2.59 (P = 0.009)											
1.4.2 Urgent abdominal aortic aneurysm rep	air (units))									
Posacioglu 2002 (Coh, urgent AAA)	5.8	3.84	40	3.63	2.87	16	16.9%	0.59 [0.00, 1.19]			
Serracino-Inglott 2005 (Coh, urgent AAA) (1)	4	6.2581	40	7	6.2581	114	21.0%	-0.48 [-0.84, -0.11]			
Tawfick 2008 (Coh, urgent AAA) (2) Subtotal (95% CI)	6	6.3853	27 107	12	6.3853	28 158	17.5% 55.5%	-0.93 [-1.48, -0.37] -0.28 [-1.06, 0.49]			
Heterogeneity: Tau ² = 0.40: Chi ² = 14.38. df = 2	2 (P = 0.00	08): ² =	86%								
Test for overall effect: Z = 0.72 (P = 0.47)	(//									
1.4.3 Urgent abdominal aortic aneurysm rep	air (mL)										
Markovic 2009 (Coh, ruptured AAA) (3)	1,890.1	1,186	30	2,755.9	1,265	30	18.2%	-0.70 [-1.22, -0.17]			
Shuhaiber 2003 (Coh, urgent AAA) Subtotal (95% CI)	2,800	857	4 34	3,161	2,155	21 51	9.9% 28.1%	-0.17 [-1.24, 0.90] -0.60 [-1.07, -0.13]	•		
Heterogeneity: Tau ² = 0.00; Chi ² = 0.75, df = 1 Test for overall effect: Z = 2.49 (P = 0.01)	(P = 0.39);	; l ² = 0%									
Total (95% CI)			162			232	100.0%	-0.44 [-0.87, -0.01]	•		
Heterogeneity: Tau ² = 0.20; Chi ² = 17.28, df = 5	5 (P = 0.00	4); l² = 7	'1%								
Test for overall effect: Z = 1.99 (P = 0.05)									-2 -1 U 1 Z Favours cell salvage Favours no cell salvage		
Test for subgroup differences: Chi ² = 1.13, df =	2 (P = 0.5	7), l² = 0	1%						i avours cen saivage i ravours no cen saivage		

Footnotes

SD estimated using calculations based on reported p-value and MD (as described in the Cochrane handbook).
 SD estimated using calculations based on reported p-value and MD (as described in the Cochrane handbook).
 Data sourced from primary study.

Figure 4.68 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (FFP)

	Experimental			Control			:	Std. Mean Difference	Std. Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% Cl		
1.5.1 Trauma (units)											
Bowley 2006 (RCT, penetrating trauma) (1) Subtotal (95% CI)	4.76	4.8	21 21	4.04	4.3	23 23	32.7% 32.7%	0.16 [-0.44, 0.75] 0.16 [-0.44, 0.75]			
Heterogeneity: Not applicable											
Test for overall effect: Z = 0.51 (P = 0.61)											
1.5.2 Urgent abdominal aortic aneurysm rep	air (units))									
Tawfick 2008 (Coh, urgent AAA) (2)	0	0	27	0	0	28		Not estimable			
Serracino-Inglott 2005 (Coh, urgent AAA) (3)	0	0	40	0	0	114		Not estimable			
Posacioglu 2002 (Coh, urgent AAA)	4.45	4.03	40	1.5	1.37	16	32.5%	0.83 [0.23, 1.43]	_ _		
Subtotal (95% CI)			107			158	32.5%	0.83 [0.23, 1.43]	-		
Heterogeneity: Not applicable											
Test for overall effect: Z = 2.71 (P = 0.007)											
1.5.3 Urgent abdominal aortic aneurysm rep	air (mL)										
Markovic 2009 (Coh, ruptured AAA) (4)	1,223.4	1,223	30	1,645.8	947	30	34.8%	-0.38 [-0.89, 0.13]			
Shuhaiber 2003 (Coh, urgent AAA) (5)	0	0	4	0	0	21		Not estimable			
Subtotal (95% CI)			34			51	34.8%	-0.38 [-0.89, 0.13]	\bullet		
Heterogeneity: Not applicable											
Test for overall effect: Z = 1.46 (P = 0.14)											
Total (95% CI)			162			232	100.0%	0.19 [-0.51, 0.89]			
Heterogeneity: Tau ² = 0.30; Chi ² = 9.07, df = 2	(P = 0.01)	; l ² = 78	%								
Test for overall effect: Z = 0.53 (P = 0.60)	· · ·										
Test for subgroup differences: Chi ² = 9.07, df =	2 (P = 0.0	1), l ² =	78.0%						Favours cell salvage Favours no cell salvage		
Footnotes											
(1) Data sourced from primary study.											
(2) Authors do not report separate data for eme	rgency AA	A repa	ir.								
(3) not reported.											

(4) Data retrieved from primary study.

(5) not reported.

Figure 4.69 Forest plot of comparison: cell salvage vs no cell salvage, outcome: Transfusion volume (PLT)

	Expe	rimer	ntal	С	ontrol			Std. Mean Difference	Std. Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% Cl
1.6.1 Trauma (units)									
Bowley 2006 (RCT, penetrating trauma) (1) Subtotal (95% CI)	1	2.2	21 21	0.56	0.94	23 23	100.0% 1 00.0%	0.26 [-0.33, 0.85] 0.26 [-0.33, 0.85]	
Heterogeneity: Not applicable Test for overall effect: Z = 0.86 (P = 0.39)									
1.6.2 Urgent abdominal aortic aneurysm re	pair (unite	s)							
Tawfick 2008 (Coh, urgent AAA) (2)	0	0	27	0	0	28		Not estimable	
Serracino-Inglott 2005 (Coh, urgent AAA) (3)	0	0	40	0	0	114		Not estimable	
Posacioglu 2002 (Coh, urgent AAA) (4) Subtotal (95% CI)	0	0	40 0	0	0	16 0		Not estimable Not estimable	
Heterogeneity: Not applicable Test for overall effect: Not applicable									
1.6.3 Urgent abdominal aortic aneurysm re	pair (mL)								
Shuhaiber 2003 (Coh, urgent AAA) (5)	0	0	4	0	0	21		Not estimable	
Markovic 2009 (Coh, ruptured AAA) (6) Subtotal (95% CI)	0	0	30 34	0	0	30 51		Not estimable Not estimable	
Heterogeneity: Not applicable Test for overall effect: Not applicable									
Total (95% CI)			55			74	100.0%	0.26 [-0.33, 0.85]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.86 (P = 0.39)									-1 -0.5 0 0.5 1 Favours cell salvage Favours no cell salvage
Test for subgroup differences: Not applicable									
Footnotes									
(1) Data sourced from primary study.									
(2) Authors do not report separate data for em	ergency A	AAA re	epair.						
(3) not reported.									
(4) not reported.									
(E) not reported									

(5) not reported.(6) not reported.

HTANALYSTS | National Blood Authority | Critical bleeding | Technical report Volume 1

Table 4.91 Results for cell salvage versus no cell salvage: Patients with critical bleeding - Transfusion volume

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			Cell salvage Mean ± SD (n)	No cell salvage Mean ± SD (n)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
Trauma setting									
Meybohm 2016 RCT High risk of bias	N = 44 (1 RCT) Bowley 2006	Patients with penetrating torso injury requiring laparotomy and	SC, emergency (South Africa)	Cell salvage vs no cell salvage	RBC transfusion volume, units (allogenic, first 24 hours)	6.47 ± 5.14 (21)	11.17 ± 6.06 (23)	MD -4.70 (-8.01, - 1.39)	Favours cell salvage p = 0.005 Heterogeneity NA
		hypotension < 90 mm Hg			FFP transfusion volume, units (allogenic, first 24 hours)	4.76 ± 4.8 (21)	4.04 ± 4.3 (23)	MD 0.72 (–1.98, 3.42) °	No significant difference <i>p</i> = 0.6 Heterogeneity NA
					PLT transfusion volume, units	1.0 ± 2.2 (21)	0.56 ± 0.94 (23)	MD 0.44 (-0.58, 1.46) ^c	No significant difference p = 0.40 Heterogeneity NA
Bhangu 2012 Coh Serious risk of bias	N = 130	Patients with combat-related injury (blast-injury, gunshot, road)	SC, combat support (Afghanistan)	Cell salvage vs no cell salvage	RBC transfusion volume, units	Total Units (n) 463 (130) Median (IQR) 14 (9.5–18.5) range 2–27	The authors estimated 7.6% total reduction w allogeneic transfusion patient cohort; and a p reduction per patient	d approximate then compared to s in the overall 130 potential median of 9.8%.	NR
					FFP transfusion volume, units Mechanism of injury (n) GSW (n=4)	Median (IQR)	NR		Test for subgroup difference p = 0.192
					Blast (n=13) Body area (n) Cavity (n=8) Extremity (n=9)	17 (10–22) 10 (4–13.5) 21 (15.5–24)			<i>p</i> = 0.004
					PLT transfusion volume, units	Median (IQR)	NR	NR	Test for subgroup difference
					(n) GSW (4)	2 (0.5–4.25)			<i>p</i> = 0.327
					BidSL (13) Body area (n) Cavity (8) Extremity (9)	3 (2-3) 2 (0.25–4.25) 3 (2.5–5.5)			<i>p</i> = 0.050

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials) included in analysis	population	(Location)			Cell salvage Mean ± SD (n)	No cell salvage Mean ± SD (n)	Risk estimate (95% Cl)	<i>Statistical significance</i> <i>p</i> -value Heterogeneity ^b
					CRYO transfusion volume, units Mechanism of injury (n) GSW (4) Blast (13)	Median (IQR) 1 (0.25-1.75) 2 (1-2)	NR	NR	Test for subgroup difference <i>p</i> = 0.335
					Body area (n) Cavity (8) Extremity (9)	1 (0–1.75) 2 (1–2)			<i>p</i> = 0.046
Surgical setting									
Shantikumar 2011 Coh Serious risk of	N= 265 (3 Coh) Posacioglu 2002 Tawfick 2008 Serrancino-Inglott 2005	Patients undergoing emergency AAA repair	SC, vascular surgery (Ireland, Serbia, Turkey,	Cell salvage vs no cell salvage	RBC transfusion volume, units (allogenic)	5.8 ± 3.84 (40) 6 (range 0–34) (27) 4 (range 0–24) (40)	3.63 ± 2.87 (16) 12 (range 3–38) (28) 7 (range 0–29) (114)	NR <i>p</i> = 0.026 NR NR <i>p</i> < 0.01	Favours no cell salvage p = NR Favours cell salvage
bias	N= 85 (2 Coh) Markovic 2009 Shuhaiber 2003		UK)		RBC transfusion volume, mL (allogenic)	1890.1 ± 1186 (30) 2800 ± 857 (4)	2755.9 ± 1265 (30) 3161 ± 2155 (21)	NR <i>p</i> = 0.0089 NR <i>p</i> = NR	<i>Favours cell salvage</i> No significant difference
	N = 56 (1 Coh) Posacioglu 2002				FFP transfusion volume, units (allogenic)	4.45 ± 4.03 (40)	1.5 ± 1.37 (16)	MD 2.95 (1.53, 4.37)	Favours no cell salvage p < 0.0001 Heterogeneity NA
	N= 60 (1 Coh) Markovic 2009				FFP transfusion volume, mL (allogenic)	1223.4 ± 1223 (30)	1645.8 ± 947 (30)	MD -422.40 ^c (-975.90, 131.10)	No significant difference p = 0.13 Heterogeneity NA
	N = 56 (1 Coh) Tawfick 2008				PLT transfusion volume, units	NR (27)	NR (28)	NR	No significant difference p = NR Heterogeneity NA

Abbreviations: AAA, abdominal aortic aneurysm; CI, confidence interval; MD; mean difference; NR, not reported; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to Systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Calculated post-hoc using RevMan 5.4. M-H. Random effects.

4.10.3.4 Financial cost

A summary of the evidence relating to transfusion volumes in patients with critical bleeding receiving intraoperative autologous transfusions (obtained by cell salvage) is presented in Table 4.92.

Overall, only limited conclusions can be drawn from the available evidence. Data appropriate to the Australian population and health care setting is needed.

In patients with penetrating trauma, there were no difference between study groups with regards to overall costs (MD –178.17, 95% CI –453.20 to 96.86) (2002 British Pound Sterling).

None of the included studies reported costs associated with cell salvage or allogenic transfusions specific to the emergency AAA patient population.

Table 4.92 Results for cell salvage versus no cell salvage: Patients with critical bleeding – Cost

Study ID	Sample size	Patient	Setting	Comparison	Outcome	Results			
Study design ^a	(no. of trials)	population	(Location)			Cell salvage	No cell salvage	Risk estimate	Statistical significance
Risk of bias	included in					Mean ± SD (n)	Mean ± SD (n)	(95% CI)	<i>p</i> -value
	analysis								Heterogeneity ^b
Trauma setting									
Meybohm 2016	N = 44 (1 RCT)	Patients with	SC, emergency	Cell salvage vs	Financial cost, £ •	812.23 ± 451.23	990.4 ± 479.48	MD -178.17	No significant difference
RCT	Bowley 2006	penetrating torso	(South Africa)	no cell salvage		(range 169.92,	(range 19.9, 1753.3)	(-453.20, 96.86) ^d	<i>p</i> = 0.2
High risk of bias		injury requiring				1747.5)			
		laparotomy and							
		nypotension < 90							
		mining							
Surgical setting									
Shantikumar	N= 350 (5 Coh)	Patients	SC, vascular	Cell salvage vs	Financial cost	None of the include	ed studies reported co	sts associated with	
2011	Markovic 2009	undergoing	surgery	no cell salvage		cell salvage or allog			
Coh	Posacioglu 2002	emergency AAA	(Germany,			emergency AAA pa	tient population.		
Serious risk of	Shuhaiber 2003	repair	Serbia, Turkey,						
bias	Serrancino-Inglott 2005		UK)						

Abbreviations: AAA, abdominal aortic aneurysm; CI, confidence interval; MD; mean difference; NR, not reported; UK, United Kingdom

a. Where only one RCT is identified in a systematic review, the evidence has been considered as RCT evidence. Where the systematic review assesses observational and cohort studies, the evidence has been considered as according to the study design features. The risk of bias of the included primary studies is based on the quality assessment reported by the systematic review authors.

b. Only applicable to systematic reviews with formal meta-analysis. Heterogeneity defined as follows: (i) no significant heterogeneity if P_{net} >0.1 and I² <25%; (ii) mild heterogeneity if I² <25%; moderate heterogeneity if I² between 25–50%; substantial heterogeneity I² >50%.

c. Data were in 2002 British Pound Sterling.

d. Calculated post-hoc using RevMan 5.4. M-H. Random effects.

5 References

1. NBA. Patient Blood Management Guidelines: Module 1 - Critical Bleeding/Massive Transfusion. <u>https://www.blood.gov.au/pbm-module-1</u>: National Blood Authority, Canberra; 2011 [cited 2011]. Available from: <u>https://www.blood.gov.au/pbm-module-1</u>.

 Higgins JPT, Lasserson T, Chandler J, Tovey D, Thomas J, Flemyng E, et al. Methodological Expectations of Cochrane Intervention Reviews. London: Cochrane; 2019 [cited 2019]. Available from: <u>https://community.cochrane.org/mecir-manual</u>.

3. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097. 10.1371/journal.pmed.1000097

4. Churchill R, Lasserson T, Chandler J, Tovey D, Thomas J, Flemyng E, et al. Standards for the reporting of new Cochrane Intervention Reviews. 2019 [cited 4 June 2020]. In: Methodological Expectations of Cochrane Intervention Reviews [Internet]. London: Cochrane, [cited 4 June 2020]. Available from: <u>https://community.cochrane.org/mecir-manual/standards-reporting-new-cochrane-intervention-reviews-r1-r109</u>.

5. Schünemann H, Brožek J, Guyatt G, Oxman A. GRADE Handbook. Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach [Internet]. 2013. [Accessed; (Updated October 2013). Available from: https://gdt.gradepro.org/app/handbook/handbook.html.

6. Mitra B, Cameron PA, Gruen RL, Mori A, Fitzgerald M, Street A. The definition of massive transfusion in trauma: a critical variable in examining evidence for resuscitation. *Eur J Emerg Med.* 2011;18(3):137-42. 10.1097/MEJ.0b013e328342310e

7. Zatta AJ, McQuilten ZK, Mitra B, Roxby DJ, Sinha R, Whitehead S, et al. Elucidating the clinical characteristics of patients captured using different definitions of massive transfusion. *Vox Sang.* 2014;107(1):60-70. 10.1111/vox.12121

8. Burtelow M, Riley E, Druzin M, Fontaine M, Viele M, Goodnough LT. How we treat: management of life-threatening primary postpartum hemorrhage with a standardized massive transfusion protocol. *Transfusion*. 2007;47(9):1564-72. 10.1111/j.1537-2995.2007.01404.x

9. Hewitt PE, Machin SJ. ABC of transfusion. Massive blood transfusion. *Bmj.* 1990;300(6717):107-9. 10.1136/bmj.300.6717.107

10. Stainsby D, MacLennan S, Hamilton PJ. Management of massive blood loss: a template guideline. *Br J Anaesth*. 2000;85(3):487-91. 10.1093/bja/85.3.487

11. Stainsby D, MacLennan S, Thomas D, Isaac J, Hamilton PJ. Guidelines on the management of massive blood loss. *British journal of haematology*. 2006;135(5):634-41. 10.1111/j.1365-2141.2006.06355.x

12. Sihler KC, Napolitano LM. Complications of massive transfusion. *Chest*. 2010;137(1):209-20. 10.1378/chest.09-0252

13. Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019): Cochrane; 2019 [cited 2019]. Available from:

www.training.cochrane.org/handbook.

14. Higgins J, Lasserson T, Chandler J, Tovey D, Thomas J, Flemyng E, et al. Methodological Expectations of Cochrane Intervention Reviews. London: Cochrane; 2019 [cited 2019]. Available from: <u>https://community.cochrane.org/mecir-manual</u>.

15. Aromataris E, Munn Z, editors. JBI Manual for Evidence Synthesis. <u>https://jbi-global-wiki.refined.site/space/MANUAL/4687404/About+this+Manual</u>: Joanna Briggs Institute; 2020.

16. Tufanaru C, Munn Z, Aromataris E, Campbell J, Hopp L. Chapter 3: Systematic reviews of effectiveness. In: Aromataris E, Munn Z, editors. *Joanna Briggs Institute Reviewer's Manual*: The Joanna Briggs Institute; 2017.

17. Gale AJ. Continuing education course #2: current understanding of hemostasis. *Toxicol Pathol.* 2011;39(1):273-80. 10.1177/0192623310389474

18. Merlin T, Weston A, Tooher R. Extending an evidence hierarchy to include topics other than treatment: revising the Australian 'levels of evidence'. *BMC medical research methodology*. 2009;9:34. 10.1186/1471-2288-9-34

19. NBA. Patient Blood Management Guidelines: Module 1 - Critical Bleeding/Massive Transfusion: Technical report volume 1 and volume 2

<u>https://www.blood.gov.au/system/files/documents/pbm-module1-tech-vol1_0.pdf</u> and <u>https://www.blood.gov.au/system/files/documents/pbm-module1-tech-vol2.pdf</u>: National Blood Authority, Canberra; 2011 [cited 2011]. Available from:

https://www.blood.gov.au/system/files/documents/pbm-module1-tech-vol1_0.pdf https://www.blood.gov.au/system/files/documents/pbm-module1-tech-vol2.pdf.

20. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008. 10.1136/bmj.j4008

Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel C, et al.
 Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC medical research methodology*. 2007;7:10. 10.1186/1471-2288-7-10
 Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The

Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. 10.1136/bmj.d5928

23. Higgins JPT, Li T, Deeks JJ. Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions version 6,0 (updated July 2019).*

24. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21(11):1539-58. 10.1002/sim.1186

25. Schünemann Holger BJ, Guyatt Gordon, and Oxman Andrew Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach. Updated October 2013. *Available at*

http://gdtguidelinedevelopmentorg/app/handbook/handbookhtml#hhnedbo8gqjqk. 2013.

26. Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, et al. GRADE guidelines: 4. Rating the quality of evidence--study limitations (risk of bias). *J Clin Epidemiol.* 2011;64(4):407-15. 10.1016/j.jclinepi.2010.07.017

27. Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 7. Rating the quality of evidence--inconsistency. *J Clin Epidemiol.* 2011;64(12):1294-302. 10.1016/j.jclinepi.2011.03.017

28. Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence--imprecision. *J Clin Epidemiol.* 2011;64(12):1283-93. 10.1016/j.jclinepi.2011.01.012

29. Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 8. Rating the quality of evidence--indirectness. *J Clin Epidemiol.* 2011;64(12):1303-10. 10.1016/j.jclinepi.2011.04.014

30. Guyatt GH, Oxman AD, Montori V, Vist G, Kunz R, Brozek J, et al. GRADE guidelines: 5. Rating the quality of evidence--publication bias. *J Clin Epidemiol*. 2011;64(12):1277-82. 10.1016/j.jclinepi.2011.01.011

31. Faraone SV. Interpreting estimates of treatment effects: implications for managed care. *P T*. 2008;33(12):700-11.

32. Santesso N, Glenton C, Dahm P, Garner P, Akl EA, Alper B, et al. GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions. *J Clin Epidemiol.* 2020;119:126-35. 10.1016/j.jclinepi.2019.10.014

33. Hultcrantz M, Rind D, Akl EA, Treweek S, Mustafa RA, Iorio A, et al. The GRADE Working Group clarifies the construct of certainty of evidence. *J Clin Epidemiol*. 2017;87:4-13. 10.1016/j.jclinepi.2017.05.006

34. Moberg J, Oxman AD, Rosenbaum S, Schunemann HJ, Guyatt G, Flottorp S, et al. The GRADE Evidence to Decision (EtD) framework for health system and public health decisions. *Health Res Policy Syst.* 2018;16(1):45. 10.1186/s12961-018-0320-2

35. Moberg J A-CP, Oxman AD. . GRADE Evidence to Decision (EtD) Frameworks Guidance. Version 1.1 2015 [cited 2015]. Available from: <u>https://ietd.epistemonikos.org/#/help/guidance</u>.

36. Vasudeva M, Mathew JK, Groombridge C, Tee JW, Johnny CS, Maini A, et al. Hypocalcemia in trauma patients: A systematic review. *The Journal of Trauma and Acute Care Surgery*. 2021;90(2):396-402. <u>https://dx.doi.org/10.1097/TA.000000000003027</u>

37. Kamyszek RW, Leraas HJ, Reed C, Ray CM, Nag UP, Poisson JL, et al. Massive transfusion in the pediatric population: A systematic review and summary of best-evidence practice strategies. *Journal of Trauma and Acute Care Surgery*. 2019;86(4):744-54. <u>http://dx.doi.org/10.1097/TA.0000000002188</u>

38. Shih AW, Al Khan S, Wang AYH, Dawe P, Young PY, Greene A, et al. Systematic reviews of scores and predictors to trigger activation of massive transfusion protocols. *Journal of Trauma and Acute Care Surgery*. 2019;87(3):717-29.

http://dx.doi.org/10.1097/TA.000000000002372

39. Lilitsis E, Xenaki S, Athanasakis E, Papadakis E, Syrogianni P, Chalkiadakis G, et al. Guiding Management in Severe Trauma: Reviewing Factors Predicting Outcome in Vastly Injured Patients. *J Emerg Trauma Shock*. 2018;11(2):80-7. 10.4103/jets.Jets_74_17

40. Tran A, Matar M, Lampron J, Steyerberg E, Taljaard M, Vaillancourt C. Early identification of patients requiring massive transfusion, embolization or hemostatic surgery for traumatic hemorrhage: A systematic review and meta-analysis. *J Trauma Acute Care Surg.* 2018;84(3):505-16. 10.1097/ta.000000000001760

41. Levy JH, Rossaint R, Zacharowski K, Spahn DR. What is the evidence for platelet transfusion in perioperative settings? *Vox Sang.* 2017;112(8):704-12. 10.1111/vox.12576

42. Poole D, Cortegiani A, Chieregato A, Russo E, Pellegrini C, De Blasio E, et al. Blood Component Therapy and Coagulopathy in Trauma: A Systematic Review of the Literature from the Trauma Update Group. *PLoS One*. 2016;11(10):e0164090. 10.1371/journal.pone.0164090

43. Baxter J, Cranfield KR, Clark G, Harris T, Bloom B, Gray AJ. Do lactate levels in the emergency department predict outcome in adult trauma patients? A systematic review. *J Trauma Acute Care Surg.* 2016;81(3):555-66. 10.1097/ta.00000000001156

44. Haas T, Fries D, Tanaka KA, Asmis L, Curry NS, Schöchl H. Usefulness of standard plasma coagulation tests in the management of perioperative coagulopathic bleeding: is there any evidence? *Br J Anaesth*. 2015;114(2):217-24. 10.1093/bja/aeu303

45. Abdul-Kadir R, McLintock C, Ducloy AS, El-Refaey H, England A, Federici AB, et al. Evaluation and management of postpartum hemorrhage: consensus from an international expert panel. *Transfusion*. 2014;54(7):1756-68. 10.1111/trf.12550

46. Pacagnella RC, Souza JP, Durocher J, Perel P, Blum J, Winikoff B, et al. A systematic review of the relationship between blood loss and clinical signs. *PloS one*. 2013;8(3):e57594-e. 10.1371/journal.pone.0057594

47. Razzaghi A, Barkun AN. Platelet transfusion threshold in patients with upper gastrointestinal bleeding: a systematic review. *J Clin Gastroenterol.* 2012;46(6):482-6. 10.1097/MCG.0b013e31823d33e3

48. Gaessler H, Helm M, Kulla M, Hossfeld B, Schmid U, Kerschowski J, et al. Prehospital evaluation and detection of induced coagulopathy in trauma: The PREDICT study. *The Journal of Trauma and Acute Care Surgery*. 2021;91(2):344-51.

https://dx.doi.org/10.1097/TA.000000000003246

49. Javali RH, Ravindra P, Patil A, Srinivasarangan M, Mundada H, Adarsh SB, et al. A Clinical Study on the Initial Assessment of Arterial Lactate and Base Deficit as Predictors of Outcome in Trauma Patients. *Indian J Crit Care Med.* 2017;21(11):719-25. 10.4103/ijccm.ijccm_218_17

50. McQuilten ZK, Wood EM, Bailey M, Cameron PA, Cooper DJ. Fibrinogen is an independent predictor of mortality in major trauma patients: A five-year statewide cohort study. *Injury*. 2017;48(5):1074-81. 10.1016/j.injury.2016.11.021

51. Kawatani Y, Nakamura Y, Kurobe H, Suda Y, Hori T. Correlations of perioperative coagulopathy, fluid infusion and blood transfusions with survival prognosis in

endovascular aortic repair for ruptured abdominal aortic aneurysm. *World J Emerg Surg.* 2016;11:29. 10.1186/s13017-016-0087-0

52. Noorbhai MA, Cassimjee HM, Sartorius B, Muckart DJ. Elevated international normalised ratios correlate with severity of injury and outcome. *S Afr Med J.* 2016;106(11):1141-5. 10.7196/SAMJ.2016.v106i11.10356

53. Sawamura A, Hayakawa M, Gando S, Kubota N, Sugano M, Wada T, et al. Disseminated intravascular coagulation with a fibrinolytic phenotype at an early phase of trauma predicts mortality. *Thromb Res.* 2009;124(5):608-13. 10.1016/j.thromres.2009.06.034

54. McQuilten ZK, Bailey M, Cameron PA, Stanworth SJ, Venardos K, Wood EM, et al. Fibrinogen concentration and use of fibrinogen supplementation with cryoprecipitate in patients with critical bleeding receiving massive transfusion: a bi-national cohort study. *British journal of haematology*. 2017;179(1):131-41. 10.1111/bjh.14804

55. Moore HB, Tessmer MT, Moore EE, Sperry JL, Cohen MJ, Chapman MP, et al. Forgot calcium? Admission ionized-calcium in two civilian randomized controlled trials of prehospital plasma for traumatic hemorrhagic shock. *J Trauma Acute Care Surg.* 2020;88(5):588-96. 10.1097/TA.00000000002614

56. Lester ELW, Fox EE, Holcomb JB, Brasel KJ, Bulger EM, Cohen MJ, et al. The impact of hypothermia on outcomes in massively transfused patients. *J Trauma Acute Care Surg.* 2019;86(3):458-63. 10.1097/TA.0000000002144

57. Consunji R, Elseed A, El-Menyar A, Sathian B, Rizoli S, Al-Thani H, et al. The effect of massive transfusion protocol implementation on the survival of trauma patients: A systematic review and meta-analysis. *Blood Transfusion*. 2020;18(6):434-45. http://dx.doi.org/10.2450/2020.0065-20

58. Kinslow K, McKenney M, Boneva D, Elkbuli A. Massive transfusion protocols in paediatric trauma population: A systematic review. *Transfusion Medicine*. 2020;30(5):333-42. <u>https://dx.doi.org/10.1111/tme.12701</u>

59. Sommer N, Schnüriger B, Candinas D, Haltmeier T. Massive transfusion protocols in nontrauma patients: A systematic review and meta-analysis. *J Trauma Acute Care Surg.* 2019;86(3):493-504. 10.1097/ta.00000000002101

60. Maw G, Furyk C. Pediatric Massive Transfusion: A Systematic Review. *Pediatr Emerg Care*. 2018;34(8):594-8. 10.1097/pec.00000000001570

61. Cannon JW, Khan MA, Raja AS, Cohen MJ, Como JJ, Cotton BA, et al. Damage control resuscitation in patients with severe traumatic hemorrhage: A practice management guideline from the Eastern Association for the Surgery of Trauma. *Journal of Trauma and Acute Care Surgery*. 2017;82(3):605-17.

http://dx.doi.org/10.1097/TA.000000000001333

62. Mitra B, O'Reilly G, Cameron PA, Zatta A, Gruen RL. Effectiveness of massive transfusion protocols on mortality in trauma: a systematic review and meta-analysis. *ANZ J Surg.* 2013;83(12):918-23. 10.1111/ans.12417

63. Vogt KN, Van Koughnett JA, Dubois L, Gray DK, Parry NG. The use of trauma transfusion pathways for blood component transfusion in the civilian population: a systematic review and meta-analysis. *Transfus Med.* 2012;22(3):156-66. 10.1111/j.1365-3148.2012.01150.x

64. Kleinveld DJB, van Amstel RBE, Wirtz MR, Geeraedts LMG, Goslings JC, Hollmann MW, et al. Platelet-to-red blood cell ratio and mortality in bleeding trauma patients: A systematic review and meta-analysis. *Transfusion*. 2021;61(S1):S243-S51. http://dx.doi.org/10.1111/trf.16455

65. Phillips AR, Tran L, Foust JE, Liang NL. Systematic review of plasma/packed red blood cell ratio on survival in ruptured abdominal aortic aneurysms. *Journal of Vascular Surgery*. 2021;73(4):1438-44. <u>http://dx.doi.org/10.1016/j.jvs.2020.10.027</u>

66. Rijnhout TWH, Duijst J, Noorman F, Zoodsma M, van Waes OJF, Verhofstad MHJ, et al. Platelet to erythrocyte transfusion ratio and mortality in massively transfused trauma patients. A systematic review and meta-analysis. *J Trauma Acute Care Surg.* 2021;91(4):759-71. 10.1097/TA.00000000003323

67. Meneses E, Boneva D, McKenney M, Elkbuli A. Massive transfusion protocol in adult trauma population. *American Journal of Emergency Medicine*. 2020;38(12):2661-6. <u>https://dx.doi.org/10.1016/j.ajem.2020.07.041</u>

68. Ritchie DT, Philbrook FGA, Leadbitter S, Kokwe KN, Meehan E, McGeady M, et al. Empirical transfusion strategies for major hemorrhage in trauma patients: A systematic review. *Journal of Trauma and Acute Care Surgery*. 2020;88(6):855-65. <u>http://dx.doi.org/10.1097/TA.0000000002685</u>

69. Rodriguez HO, Rios F, Rubio C, Arsanios DM, Herazo AF, Beltran LM, et al. Mortality in civilian trauma patients and massive blood transfusion treated with high vs low plasma: Red blood cell ratio. Systematic review and meta-analysis. *Colombian Journal of Anesthesiology*. 2020;48(3):126-37. http://dx.doi.org/10.1097/CJ9.000000000000161

70. Wirtz MR, Schalkers DV, Goslings JC, Juffermans NP. The impact of blood product ratio and procoagulant therapy on the development of thromboembolic events in severely injured hemorrhaging trauma patients. *Transfusion*. 2020;60(8):1873-82. http://dx.doi.org/10.1111/trf.15917

71. da Luz LT, Shah PS, Strauss R, Mohammed AA, D'Empaire PP, Tien H, et al. Does the evidence support the importance of high transfusion ratios of plasma and platelets to red blood cells in improving outcomes in severely injured patients: a systematic review and meta-analyses. *Transfusion*. 2019;59(11):3337-49. <u>http://dx.doi.org/10.1111/trf.15540</u>

72. McQuilten ZK, Crighton G, Brunskill S, Morison JK, Richter TH, Waters N, et al. Optimal Dose, Timing and Ratio of Blood Products in Massive Transfusion: Results from a Systematic Review. *Transfus Med Rev.* 2018;32(1):6-15. 10.1016/j.tmrv.2017.06.003

73. Rahouma M, Kamel M, Jodeh D, Kelley T, Ohmes LB, de Biasi AR, et al. Does a balanced transfusion ratio of plasma to packed red blood cells improve outcomes in both trauma and surgical patients? A meta-analysis of randomized controlled trials and observational studies. *Am J Surg.* 2018;216(2):342-50. 10.1016/j.amjsurg.2017.08.045

74. Jones AR, Frazier SK. Association of Blood Component Ratio With Clinical Outcomes in Patients After Trauma and Massive Transfusion: A Systematic Review. *Adv Emerg Nurs J.* 2016;38(2):157-68. 10.1097/tme.000000000000003

75. Tapia NM, Suliburk J, Mattox KL. The initial trauma center fluid management of penetrating injury: a systematic review. *Clin Orthop Relat Res.* 2013;471(12):3961-73. 10.1007/s11999-013-3122-4

76. Peralta R, Vijay A, El-Menyar A, Consunji R, Afifi I, Mahmood I, et al. Early high ratio platelet transfusion in trauma resuscitation and its outcomes. *Int J Crit IIIn Inj Sci.* 2016;6(4):188-93. 10.4103/2229-5151.195448

77. Balvers K, Wirtz MR, van Dieren S, Goslings JC, Juffermans NP. Risk factors for trauma-induced coagulopathy- and transfusion-associated multiple organ failure in severely injured trauma patients. *Front Med (Lausanne)*. 2015;2:24. 10.3389/fmed.2015.00024

78. Patel SV, Kidane B, Klingel M, Parry N. Risks associated with red blood cell transfusion in the trauma population, a meta-analysis. *Injury*. 2014;45(10):1522-33. 10.1016/j.injury.2014.05.015

79. Liu S, Fujii Q, Serio F, McCague A. Massive Blood Transfusions and Outcomes in Trauma Patients; An Intention to Treat Analysis. *Bull Emerg Trauma*. 2018;6(3):217-20. 10.29252/beat-060305

80. Hassanien M, El-Talkawy MD, El-Ghannam M, El Ray A, Ali AA, Taleb HA. Predictors of In-Hospital Mortality in patients with hepatocellular carcinoma and Acute Variceal bleeding. *Electron Physician*. 2015;7(6):1336-43. 10.14661/1336

81. McQuilten ZK, Crighton G, Engelbrecht S, Gotmaker R, Brunskill SJ, Murphy MF, et al. Transfusion interventions in critical bleeding requiring massive transfusion: A systematic review. *Transfusion Medicine Reviews*. 2015;29(2):127-37. http://dx.doi.org/10.1016/j.tmrv.2015.01.001

82. Magon N, Babu K. Recombinant Factor VIIa in Post-partum Hemorrhage: A New Weapon in Obstetrician's Armamentarium. *N Am J Med Sci.* 2012;4(4):157-62. 10.4103/1947-2714.94938

83. Okonta KE, Edwin F, Falase B. Is recombinant activated factor VII effective in the treatment of excessive bleeding after paediatric cardiac surgery? *Interactive Cardiovascular and Thoracic Surgery*. 2012;15(4):690-5.

http://dx.doi.org/10.1093/icvts/ivs309

84. Simpson E, Lin Y, Stanworth S, Birchall J, Doree C, Hyde C. Recombinant factor VIIa for the prevention and treatment of bleeding in patients without haemophilia. *Cochrane database of systematic reviews (Online).* 2012;3:CD005011.

85. Curry N, Hopewell S, Doree C, Hyde C, Brohi K, Stanworth S. The acute management of trauma hemorrhage: A systematic review of randomized controlled trials. *Critical Care*. 2011;15 (2) (no pagination)(R92). <u>http://dx.doi.org/10.1186/cc10096</u>

86. Yank V, Tuohy CV, Logan AC, Bravata DM, Staudenmayer K, Eisenhut R, et al. Systematic review: benefits and harms of in-hospital use of recombinant factor VIIa for off-label indications. *Annals of Internal Medicine*. 2011;154(8):529-40.

https://dx.doi.org/10.7326/0003-4819-154-8-201104190-00004

87. Franchini M, Franchi M, Bergamini V, Montagnana M, Salvagno GL, Targher G, et al. The use of recombinant activated FVII in postpartum hemorrhage. *Clinical Obstetrics and Gynecology*. 2010;53(1):219-27. <u>http://dx.doi.org/10.1097/GRF.0b013e3181cc4378</u>

88. Dutton RP, Parr M, Tortella BJ, Champion HR, Bernard GR, Boffard K, et al. Recombinant activated factor VII safety in trauma patients: results from the CONTROL trial. *The Journal of trauma*. 2011;71(1):12-9. 10.1097/TA.0b013e31821a42cf

89. Hauser CJ, Boffard K, Dutton R, Bernard GR, Croce MA, Holcomb JB, et al. Results of the CONTROL trial: efficacy and safety of recombinant activated Factor VII in the management of refractory traumatic hemorrhage. *The Journal of trauma*. 2010;69(3):489-500. 10.1097/TA.0b013e3181edf36e

90. McMullin NR, Wade CE, Holcomb JB, Nielsen TG, Rossaint R, Riou B, et al. Prolonged prothrombin time after recombinant activated factor VII therapy in critically bleeding trauma patients is associated with adverse outcomes. *The Journal of trauma*. 2010;69(1):60-9. 10.1097/TA.0b013e3181e17260

91. Boffard KD, Choong PI, Kluger Y, Riou B, Rizoli SB, Rossaint R, et al. The treatment of bleeding is to stop the bleeding! Treatment of trauma-related hemorrhage. *Transfusion*. 2009;49 Suppl 5:240S-7S. 10.1111/j.1537-2995.2008.01987.x

92. Gill R, Herbertson M, Vuylsteke A, Olsen PS, von Heymann C, Mythen M, et al. Safety and efficacy of recombinant activated factor VII: a randomized placebo-controlled trial in the setting of bleeding after cardiac surgery. *Circulation*. 2009;120(1):21-7. 10.1161/CIRCULATIONAHA.108.834275

93. Bosch J, Thabut D, Albillos A, Carbonell N, Spicak J, Massard J, et al. Recombinant factor VIIa for variceal bleeding in patients with advanced cirrhosis: A randomized, controlled trial. *Hepatology*. 2008;47(5):1604-14. 10.1002/hep.22216

94. Rizoli SB, Boffard KD, Riou B, Warren B, Iau P, Kluger Y, et al. Recombinant activated factor VII as an adjunctive therapy for bleeding control in severe trauma patients with coagulopathy: subgroup analysis from two randomized trials. *Crit Care*. 2006;10(6):R178. 10.1186/cc5133

95. Boffard KD, Riou B, Warren B, Choong PI, Rizoli S, Rossaint R, et al. Recombinant factor VIIa as adjunctive therapy for bleeding control in severely injured trauma patients: two parallel randomized, placebo-controlled, double-blind clinical trials. *The Journal of trauma*. 2005;59(1):8-15; discussion -8. 10.1097/01.ta.0000171453.37949.b7

96. Chuansumrit A, Wangruangsatid S, Lektrakul Y, Chua MN, Zeta Capeding MR, Bech OM, et al. Control of bleeding in children with Dengue hemorrhagic fever using recombinant activated factor VII: a randomized, double-blind, placebo-controlled study. *Blood Coagul Fibrinolysis*. 2005;16(8):549-55. 10.1097/01.mbc.0000186837.78432.2f

97. Pihusch M, Bacigalupo A, Szer J, von Depka Prondzinski M, Gaspar-Blaudschun B, Hyveled L, et al. Recombinant activated factor VII in treatment of bleeding complications following hematopoietic stem cell transplantation. *J Thromb Haemost*. 2005;3(9):1935-44. 10.1111/j.1538-7836.2005.01523.x

98. Bosch J, Thabut D, Bendtsen F, D'Amico G, Albillos A, Gonzalez Abraldes J, et al. Recombinant factor VIIa for upper gastrointestinal bleeding in patients with cirrhosis: a randomized, double-blind trial. *Gastroenterology*. 2004;127(4):1123-30. 10.1053/j.gastro.2004.07.015

99. Lavigne-Lissalde G, Aya AG, Mercier FJ, Roger-Christoph S, Chauleur C, Morau E, et al. Recombinant human FVIIa for reducing the need for invasive second-line therapies in severe refractory postpartum hemorrhage: a multicenter, randomized, open controlled trial. *J Thromb Haemost*. 2015;13(4):520-9. 10.1111/jth.12844

100. Veazie S, Kondo K, Winchell K. Comparative Effectiveness of In-Hospital Use of Recombinant Factor VIIa for Off-Label Indications vs Usual Care. *Agency for Healthcare Research and Quality Systematic Review Surveillance Program*. 2016.

101. Zatta A, McQuilten Z, Kandane-Rathnayake R, Isbister J, Dunkley S, McNeil J, et al. The Australian and New Zealand Haemostasis Registry: ten years of data on off-licence use of recombinant activated factor VII. *Blood Transfus*. 2015;13(1):86-99. 10.2450/2014.0260-13

102. Stabler SN, Li SS, Karpov A, Vu EN. Use of fibrinogen concentrate for traumarelated bleeding: A systematic-review and meta-analysis. *Journal of Trauma and Acute Care Surgery*. 2020;89(6):1212-24. <u>http://dx.doi.org/10.1097/TA.00000000002920</u>

103. van den Brink DP, Wirtz MR, Neto AS, Schochl H, Viersen V, Binnekade J, et al. Effectiveness of prothrombin complex concentrate for the treatment of bleeding: A systematic review and meta-analysis. *Journal of Thrombosis and Haemostasis*. 2020;18(10):2457-67. <u>http://dx.doi.org/10.1111/jth.14991</u>

104. Zaidi A, Kohli R, Daru J, Estcourt L, Khan KS, Thangaratinam S, et al. Early Use of Fibrinogen Replacement Therapy in Postpartum Hemorrhage-A Systematic Review. *Transfusion Medicine Reviews*. 2020;34(2):101-7. <u>http://dx.doi.org/10.1016/j.tmrv.2019.12.002</u>

105. Coccolini F, Pizzilli G, Corbella D, Sartelli M, Agnoletti V, Agostini V, et al. Prehospital plasma in haemorrhagic shock management: current opinion and meta-analysis of randomized trials. *World journal of emergency surgery : WJES*. 2019;14:6. http://dx.doi.org/10.1186/s13017-019-0226-5

106. Fabes J, Brunskill SJ, Curry N, Doree C, Stanworth SJ. Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia. *Cochrane Database of Systematic Reviews*. 2018;12:CD010649. https://dx.doi.org/10.1002/14651858.CD010649.pub2

https://dx.doi.org/10.1002/14651858.CD010649.pub2

107. Mengoli C, Franchini M, Marano G, Pupella S, Vaglio S, Marietta M, et al. The use of fibrinogen concentrate for the management of trauma-related bleeding: A systematic review and meta-analysis. 2017;15(4):318-24. <u>http://dx.doi.org/10.2450/2017.0094-17</u>
108. Aubron C, Reade MC, Fraser JF, Cooper DJ. Efficacy and safety of fibrinogen

concentrate in trauma patients-a systematic review. 2014;29(3):471.e11-.e17. http://dx.doi.org/10.1016/j.jcrc.2013.12.011

109. Lunde J, Stensballe J, Wikkelso A, Johansen M, Afshari A. Fibrinogen concentrate for bleeding - A systematic review. 2014;58(9):1061-74. <u>http://dx.doi.org/10.1111/aas.12370</u>
110. Warmuth M, Mad P, Wild C. Systematic review of the efficacy and safety of fibrinogen concentrate substitution in adults. 2012;56(5):539-48.

https://dx.doi.org/10.1111/j.1399-6576.2011.02586.x

111. Lucena LSd, Rodrigues RdR, Carmona MJC, Noronha FJD, Oliveira HP, Lima NM, et al. Early administration of fibrinogen concentrate in patients with polytrauma with thromboelastometry suggestive of hypofibrinogenemia: A randomized feasibility trial. *Clinics*. 2021;76:-. 10.6061/clinics/2021/e3168

112. Akbari E, Safari S, Hatamabadi H. The effect of fibrinogen concentrate and fresh frozen plasma on the outcome of patients with acute traumatic coagulopathy: A quasi-experimental study. *Am J Emerg Med.* 2018;36(11):1947-50. 10.1016/j.ajem.2018.02.018

113. Curry N, Foley C, Wong H, Mora A, Curnow E, Zarankaite A, et al. Early fibrinogen concentrate therapy for major haemorrhage in trauma (E-FIT 1): results from a UK multi-centre, randomised, double blind, placebo-controlled pilot trial. *Crit Care*. 2018;22(1):164. 10.1186/s13054-018-2086-x

114. Moore HB, Moore EE, Chapman MP, McVaney K, Bryskiewicz G, Blechar R, et al. Plasma-first resuscitation to treat haemorrhagic shock during emergency ground

transportation in an urban area: a randomised trial. *Lancet*. 2018;392(10144):283-91. 10.1016/s0140-6736(18)31553-8

115. Sperry JL, Guyette FX, Brown JB, Yazer MH, Triulzi DJ, Early-Young BJ, et al. Prehospital Plasma during Air Medical Transport in Trauma Patients at Risk for Hemorrhagic Shock. *N Engl J Med*. 2018;379(4):315-26. 10.1056/NEJMoa1802345

116. Innerhofer P, Fries D, Mittermayr M, Innerhofer N, von Langen D, Hell T, et al. Reversal of trauma-induced coagulopathy using first-line coagulation factor concentrates or fresh frozen plasma (RETIC): a single-centre, parallel-group, open-label, randomised trial. *Lancet Haematol.* 2017;4(6):e258-e71. 10.1016/s2352-3026(17)30077-7

117. Nascimento B, Callum J, Tien H, Peng H, Rizoli S, Karanicolas P, et al. Fibrinogen in the initial resuscitation of severe trauma (FiiRST): a randomized feasibility trial. *Br J Anaesth.* 2016;117(6):775-82. 10.1093/bja/aew343

118. Curry N, Rourke C, Davenport R, Beer S, Pankhurst L, Deary A, et al. Early cryoprecipitate for major haemorrhage in trauma: a randomised controlled feasibility trial. *BJA: British Journal of Anaesthesia.* 2015;115(1):76-83. 10.1093/bja/aev134

119. Bilecen S, de Groot JA, Kalkman CJ, Spanjersberg AJ, Brandon Bravo Bruinsma GJ, Moons KG, et al. Effect of Fibrinogen Concentrate on Intraoperative Blood Loss Among Patients With Intraoperative Bleeding During High-Risk Cardiac Surgery: A Randomized Clinical Trial. *JAMA*. 2017;317(7):738-47. 10.1001/jama.2016.21037

120. Rahe-Meyer N, Levy JH, Mazer CD, Schramko A, Klein AA, Brat R, et al. Randomized evaluation of fibrinogen vs placebo in complex cardiovascular surgery (REPLACE): a double-blind phase III study of haemostatic therapy. *Br J Anaesth*. 2016;117(1):41-51. 10.1093/bja/aew169

121. Tanaka KA, Egan K, Szlam F, Ogawa S, Roback JD, Sreeram G, et al. Transfusion and hematologic variables after fibrinogen or platelet transfusion in valve replacement surgery: preliminary data of purified lyophilized human fibrinogen concentrate versus conventional transfusion. *Transfusion*. 2014;54(1):109-18. 10.1111/trf.12248

122. Rahe-Meyer N, Hanke A, Schmidt DS, Hagl C, Pichlmaier M. Fibrinogen concentrate reduces intraoperative bleeding when used as first-line hemostatic therapy during major aortic replacement surgery: results from a randomized, placebo-controlled trial. *J Thorac Cardiovasc Surg.* 2013;145(3 Suppl):S178-85. 10.1016/j.jtcvs.2012.12.083

123. Galas FR, de Almeida JP, Fukushima JT, Vincent JL, Osawa EA, Zeferino S, et al. Hemostatic effects of fibrinogen concentrate compared with cryoprecipitate in children after cardiac surgery: a randomized pilot trial. *J Thorac Cardiovasc Surg.* 2014;148(4):1647-55. 10.1016/j.jtcvs.2014.04.029

124. Collins PW, Cannings-John R, Bruynseels D, Mallaiah S, Dick J, Elton C, et al. Viscoelastometric-guided early fibrinogen concentrate replacement during postpartum haemorrhage: OBS2, a double-blind randomized controlled trial. *Br J Anaesth*. 2017;119(3):411-21. 10.1093/bja/aex181

125. Wikkelso AJ, Edwards HM, Afshari A, Stensballe J, Langhoff-Roos J, Albrechtsen C, et al. Pre-emptive treatment with fibrinogen concentrate for postpartum haemorrhage: randomized controlled trial. *Br J Anaesth*. 2015;114(4):623-33. 10.1093/bja/aeu444

126. Almskog LM, Hammar U, Wikman A, Ostlund A, Svensson J, Wanecek M, et al. A retrospective register study comparing fibrinogen treated trauma patients with an injury severity score matched control group. *Scand J Trauma Resusc Emerg Med.* 2020;28(1):5. 10.1186/s13049-019-0695-2

127. Zeeshan M, Hamidi M, Feinstein AJ, Gries L, Jehan F, Sakran J, et al. Four-factor prothrombin complex concentrate is associated with improved survival in trauma-related hemorrhage: A nationwide propensity-matched analysis. *J Trauma Acute Care Surg.* 2019;87(2):274-81. 10.1097/TA.0000000002262

128. Jehan F, Aziz H, O'Keeffe T, Khan M, Zakaria ER, Hamidi M, et al. The role of fourfactor prothrombin complex concentrate in coagulopathy of trauma: A propensity matched analysis. *J Trauma Acute Care Surg.* 2018;85(1):18-24. 10.1097/TA.000000000001938

129. Holcomb JB, Swartz MD, DeSantis SM, Greene TJ, Fox EE, Stein DM, et al. Multicenter observational prehospital resuscitation on helicopter study. *J Trauma Acute Care Surg.* 2017;83(1 Suppl 1):S83-S91. 10.1097/TA.00000000001484

130. Inokuchi K, Sawano M, Yamamoto K, Yamaguchi A, Sugiyama S. Early administration of fibrinogen concentrates improves the short-term outcomes of severe pelvic fracture patients. *Acute Med Surg.* 2017;4(3):271-7. 10.1002/ams2.268

131. Shackelford SA, Del Junco DJ, Powell-Dunford N, Mazuchowski EL, Howard JT, Kotwal RS, et al. Association of Prehospital Blood Product Transfusion During Medical Evacuation of Combat Casualties in Afghanistan With Acute and 30-Day Survival. *JAMA*. 2017;318(16):1581-91. 10.1001/jama.2017.15097

132. Joseph B, Khalil M, Harrison C, Swartz T, Kulvatunyou N, Haider AA, et al. Assessing the Efficacy of Prothrombin Complex Concentrate in Multiply Injured Patients With High-Energy Pelvic and Extremity Fractures. *J Orthop Trauma*. 2016;30(12):653-8. 10.1097/BOT.00000000000665

133. Joseph B, Aziz H, Pandit V, Hays D, Kulvatunyou N, Yousuf Z, et al. Prothrombin complex concentrate versus fresh-frozen plasma for reversal of coagulopathy of trauma: is there a difference? *World J Surg.* 2014;38(8):1875-81. 10.1007/s00268-014-2631-y

134. O'Reilly DJ, Morrison JJ, Jansen JO, Apodaca AN, Rasmussen TE, Midwinter MJ. Prehospital blood transfusion in the en route management of severe combat trauma: a matched cohort study. *J Trauma Acute Care Surg.* 2014;77(3 Suppl 2):S114-20. 10.1097/TA.000000000000328

135. Innerhofer P, Westermann I, Tauber H, Breitkopf R, Fries D, Kastenberger T, et al. The exclusive use of coagulation factor concentrates enables reversal of coagulopathy and decreases transfusion rates in patients with major blunt trauma. *Injury*. 2013;44(2):209-16. 10.1016/j.injury.2012.08.047

136. Wafaisade A, Lefering R, Maegele M, Brockamp T, Mutschler M, Lendemans S, et al. Administration of fibrinogen concentrate in exsanguinating trauma patients is associated with improved survival at 6 hours but not at discharge. *J Trauma Acute Care Surg.* 2013;74(2):387-3; discussion 93-5. 10.1097/TA.0b013e31827e2410

137. Nienaber U, Innerhofer P, Westermann I, Schochl H, Attal R, Breitkopf R, et al. The impact of fresh frozen plasma vs coagulation factor concentrates on morbidity and mortality in trauma-associated haemorrhage and massive transfusion. *Injury.* 2011;42(7):697-701. 10.1016/j.injury.2010.12.015

138. Schochl H, Nienaber U, Maegele M, Hochleitner G, Primavesi F, Steitz B, et al. Transfusion in trauma: thromboelastometry-guided coagulation factor concentratebased therapy versus standard fresh frozen plasma-based therapy. *Crit Care.* 2011;15(2):R83. 10.1186/cc10078

139. Bilecen S, Peelen LM, Kalkman CJ, Spanjersberg AJ, Moons KG, Nierich AP. Fibrinogen concentrate therapy in complex cardiac surgery. *J Cardiothorac Vasc Anesth*. 2013;27(1):12-7. 10.1053/j.jvca.2012.06.006

140. Rahe-Meyer N, Solomon C, Winterhalter M, Piepenbrock S, Tanaka K, Haverich A, et al. Thromboelastometry-guided administration of fibrinogen concentrate for the treatment of excessive intraoperative bleeding in thoracoabdominal aortic aneurysm surgery. *J Thorac Cardiovasc Surg.* 2009;138(3):694-702. 10.1016/j.jtcvs.2008.11.065

141. Rahe-Meyer N, Pichlmaier M, Haverich A, Solomon C, Winterhalter M, Piepenbrock S, et al. Bleeding management with fibrinogen concentrate targeting a high-normal plasma fibrinogen level: a pilot study. *Br J Anaesth.* 2009;102(6):785-92. 10.1093/bja/aep089 142. Ahmed S, Harrity C, Johnson S, Varadkar S, McMorrow S, Fanning R, et al. The efficacy of fibrinogen concentrate compared with cryoprecipitate in major obstetric haemorrhage--an observational study. *Transfus Med.* 2012;22(5):344-9. 10.1111/j.1365-3148.2012.01178.x

143. Al-Jeabory M, Szarpak L, Attila K, Simpson M, Smereka A, Gasecka A, et al. Efficacy and Safety of Tranexamic Acid in Emergency Trauma: A Systematic Review and Meta-Analysis. *J Clin Med*. 2021;10(5). 10.3390/jcm10051030

144. Almuwallad A, Cole E, Ross J, Perkins Z, Davenport R. The impact of prehospital TXA on mortality among bleeding trauma patients: A systematic review and metaanalysis. *J Trauma Acute Care Surg.* 2021;90(5):901-7. 10.1097/ta.000000000003120

145. Della Corte L, Saccone G, Locci M, Carbone L, Raffone A, Giampaolino P, et al. Tranexamic acid for treatment of primary postpartum hemorrhage after vaginal delivery: a systematic review and meta-analysis of randomized controlled trials. *J Matern Fetal Neonatal Med.* 2020;33(5):869-74. 10.1080/14767058.2018.1500544

146. Chornenki NLJ, Um KJ, Mendoza PA, Samienezhad A, Swarup V, Chai-Adisaksopha C, et al. Risk of venous and arterial thrombosis in non-surgical patients receiving systemic tranexamic acid: A systematic review and meta-analysis. *Thrombosis Research*. 2019;179(pp 81-86). http://dx.doi.org/10.1016/j.thromres.2019.05.003

147. Marsden MER, Rossetto A, Duffield CAB, Woolley TGD, Buxton WP, Steynberg S, et al. Prehospital tranexamic acid shortens the interval to administration by half in Major Trauma Networks: A service evaluation. *Emergency Medicine Journal*. 2019. http://dx.doi.org/10.1136/emermed-2018-208118

148. EI-Menyar A, Sathian B, Asim M, Latifi R, Al-Thani H. Efficacy of prehospital administration of tranexamic acid in trauma patients: A meta-analysis of the randomized controlled trials. *American Journal of Emergency Medicine*. 2018;36(6):1079-87. http://dx.doi.org/10.1016/j.ajem.2018.03.033

149. Gayet-Ageron A, Prieto-Merino D, Ker K, Shakur H, Ageron F-X, Roberts I, et al. Effect of treatment delay on the effectiveness and safety of antifibrinolytics in acute severe haemorrhage: a meta-analysis of individual patient-level data from 40 138 bleeding patients. *Lancet*. 2018;391(10116):125-32. <u>https://dx.doi.org/10.1016/S0140-6736(17)32455-8</u>

150. Shakur H, Beaumont D, Pavord S, Gayet-Ageron A, Ker K, Mousa HA. Antifibrinolytic drugs for treating primary postpartum haemorrhage. *Cochrane Database of Systematic Reviews*. 2018;2018 (2) (no pagination)(CD012964).

http://dx.doi.org/10.1002/14651858.CD012964

151. Huebner BR, Dorlac WC, Cribari C. Tranexamic Acid Use in Prehospital Uncontrolled Hemorrhage. *Wilderness & environmental medicine*. 2017;S. 28(2):S50-S60. <u>http://dx.doi.org/10.1016/j.wem.2016.12.006</u>

152. Nishida T, Kinoshita T, Yamakawa K. Tranexamic acid and trauma-induced coagulopathy. *Journal of Intensive Care*. 2017;5 (1) (no pagination)(5). http://dx.doi.org/10.1186/s40560-016-0201-0

153. Ausset S, Glassberg E, Nadler R, Sunde G, Cap AP, Hoffmann C, et al. Tranexamic acid as part of remote damage-control resuscitation in the prehospital setting: A critical appraisal of the medical literature and available alternatives. *The Journal of Trauma and Acute Care Surgery*. 2015;78(6 Suppl 1):S70-5.

https://dx.doi.org/10.1097/TA.000000000000640

154. Ker K, Roberts I, Shakur H, Coats TJ. Antifibrinolytic drugs for acute traumatic injury. *Cochrane Database of Systematic Reviews*. 2015(5):CD004896. <u>https://dx.doi.org/10.1002/14651858.CD004896.pub4</u>

155. Baskaran D, Rahman S, Salmasi Y, Froghi S, Berber O, George M. Effect of tranexamic acid use on blood loss and thromboembolic risk in hip fracture surgery: systematic review and meta-analysis. *HIP International*. 2018;28(1):3-10. http://dx.doi.org/10.5301/hipint.5000556

156. Gausden EB, Qudsi R, Boone MD, O'Gara B, Ruzbarsky JJ, Lorich DG. Tranexamic Acid in Orthopaedic Trauma Surgery: A Meta-Analysis. *Journal of Orthopaedic Trauma*. 2017;31(10):513-9. <u>https://dx.doi.org/10.1097/BOT.000000000000913</u>

157. Wang W, Yu J. Tranexamic acid reduces blood loss in intertrochanteric fractures: A meta-analysis from randomized controlled trials. *Medicine*. 2017;96(52):e9396. <u>https://dx.doi.org/10.1097/MD.00000000009396</u>

158. Zhang P, He J, Fang Y, Chen P, Liang Y, Wang J. Efficacy and safety of intravenous tranexamic acid administration in patients undergoing hip fracture surgery for hemostasis. *Medicine (United States)*. 2017;96 (21) (no pagination)(e6940). http://dx.doi.org/10.1097/MD.00000000006940 159. Bennett C, Klingenberg LS, Langholz E, Gluud LL. Tranexamic acid for upper gastrointestinal bleeding. *Cochrane Database of Systematic Reviews*. 2014(11).
160. Perel P, Ker K, Morales Uribe CH, Roberts I. Tranexamic acid for reducing mortality in emergency and urgent surgery. *Cochrane database of systematic reviews (Online)*. 2013;1:CD010245.

161. Burke E, Harkins P, Ahmed I. Is There a Role for Tranexamic Acid in Upper GI Bleeding? A Systematic Review and Meta-Analysis. *Surg Res Pract.* 2021;2021:8876991. 10.1155/2021/8876991

162. Lee PL, Yang KS, Tsai HW, Hou SK, Kang YN, Chang CC. Tranexamic acid for gastrointestinal bleeding: A systematic review with meta-analysis of randomized clinical trials. *Am J Emerg Med*. 2021;45:269-79. 10.1016/j.ajem.2020.08.062

163. Ageron FX, Gayet-Ageron A, Ker K, Coats TJ, Shakur-Still H, Roberts I, et al. Effect of tranexamic acid by baseline risk of death in acute bleeding patients: a meta-analysis of individual patient-level data from 28 333 patients. *Br J Anaesth*. 2020;124(6):676-83. 10.1016/j.bja.2020.01.020

164. Guyette FX, Brown JB, Zenati MS, Early-Young BJ, Adams PW, Eastridge BJ, et al. Tranexamic Acid During Prehospital Transport in Patients at Risk for Hemorrhage After Injury: A Double-blind, Placebo-Controlled, Randomized Clinical Trial. *JAMA Surg.* 2020;156(1):11-20. 10.1001/jamasurg.2020.4350

165. Kakaei F, Virani P, Hashemzadeh S, Zarrintan S, Beheshtirouy; S, et al. Effects of Tranexamic Acid on Mortality and Blood Transfusion in Trauma Patients with Significant Hemorrhage: A Clinical Trial. *Advances in Biosciences & Clinical Medicine*. 2017;5(4):24-31. 10.7575/aiac.abcmed.17.05.04.04

166. collaborators C-t, Shakur H, Roberts I, Bautista R, Caballero J, Coats T, et al. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet.* 2010;376(9734):23-32. 10.1016/S0140-6736(10)60835-5

167. Collaborators H-IT. Effects of a high-dose 24-h infusion of tranexamic acid on death and thromboembolic events in patients with acute gastrointestinal bleeding (HALT-IT): an international randomised, double-blind, placebo-controlled trial. *Lancet.* 2020;395(10241):1927-36. 10.1016/S0140-6736(20)30848-5

168. Roberts I, Coats T, Edwards P, Gilmore I, Jairath V, Ker K, et al. HALT-IT--tranexamic acid for the treatment of gastrointestinal bleeding: study protocol for a randomised controlled trial. *Trials*. 2014;15:450. 10.1186/1745-6215-15-450

169. Collaborators WT. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet*. 2017;389(10084):2105-16. 10.1016/S0140-6736(17)30638-4

170. Ducloy-Bouthors AS, Jude B, Duhamel A, Broisin F, Huissoud C, Keita-Meyer H, et al. High-dose tranexamic acid reduces blood loss in postpartum haemorrhage. *Crit Care*. 2011;15(2):R117. 10.1186/cc10143

171. Li C, Zhao Q, Yang K, Jiang L, Yu J. Thromboelastography or rotational thromboelastometry for bleeding management in adults undergoing cardiac surgery: a systematic review with meta-analysis and trial sequential analysis. *J Thorac Dis.* 2019;11(4):1170-81. 10.21037/jtd.2019.04.39

172. Roullet S, de Maistre E, Ickx B, Blais N, Susen S, Faraoni D, et al. Position of the French Working Group on Perioperative Haemostasis (GIHP) on viscoelastic tests: What role for which indication in bleeding situations? *Anaesthesia Critical Care and Pain Medicine*. 2018. <u>http://dx.doi.org/10.1016/j.accpm.2017.12.014</u>

173. Fahrendorff M, Oliveri RS, Johansson PI. The use of viscoelastic haemostatic assays in goal-directing treatment with allogeneic blood products - A systematic review and meta-analysis. *Scandinavian journal of trauma, resuscitation and emergency medicine.* 2017;25(1):39. <u>http://dx.doi.org/10.1186/s13049-017-0378-9</u>

174. Serraino GF, Murphy GJ. Routine use of viscoelastic blood tests for diagnosis and treatment of coagulopathic bleeding in cardiac surgery: Updated systematic review and

meta-analysis. *British Journal of Anaesthesia*. 2017;118(6):823-33. <u>http://dx.doi.org/10.1093/bja/aex100</u>

175. Wikkelso A, Wetterslev J, Moller MA, Afshari A. Thromboelastography (TEG) or thromboelastometry (ROTEM) to monitor haemostatic treatment versus usual care in adults or children with bleeding. *Cochrane Database of Systematic Reviews*. 2017(1).

176. Wikkelso A, Wetterslev J, Moller AM, Afshari A. Thromboelastography (TEG) or rotational thromboelastometry (ROTEM) to monitor haemostatic treatment in bleeding patients: a systematic review with meta-analysis and trial sequential analysis. *Anaesthesia*. 2017;72(4):519-31. http://dx.doi.org/10.1111/anae.13765

177. Saner FH, Kirchner C. Monitoring and Treatment of Coagulation Disorders in End-Stage Liver Disease. *Visc Med.* 2016;32(4):241-8. 10.1159/000446304

178. Deppe AC, Weber C, Zimmermann J, Kuhn EW, Slottosch I, Liakopoulos OJ, et al. Point-of-care thromboelastography/thromboelastometry-based coagulation management in cardiac surgery: A meta-analysis of 8332 patients. *Journal of Surgical Research*. 2016;203(2):424-33. <u>http://dx.doi.org/10.1016/j.jss.2016.03.008</u>

179. Corredor C, Wasowicz M, Karkouti K, Sharma V. The role of point-of-care platelet function testing in predicting postoperative bleeding following cardiac surgery: A systematic review and meta-analysis. *Anaesthesia*. 2015;70(6):715-31. http://dx.doi.org/10.1111/anae.13083

180. Da Luz LT, Nascimento B, Shankarakutty AK, Rizoli S, Adhikari NK. Effect of thromboelastography (TEG) and rotational thromboelastometry (ROTEM) on diagnosis of coagulopathy, transfusion guidance and mortality in trauma: descriptive systematic review. *Critical Care (London, England)*. 2014;18(5):518. <u>https://dx.doi.org/10.1186/s13054-014-0518-9</u>

181. Haas T, Gorlinger K, Grassetto A, Agostini V, Simioni P, Nardi G, et al.
Thromboelastometry for guiding bleeding management of the critically ill patient: a systematic review of the literature. *Minerva Anestesiologica*. 2014;80(12):1320-35.
182. Agarwal S, Johnson RI, Shaw M. Preoperative point-of-care platelet function testing in cardiac surgery. *J Cardiothorac Vasc Anesth*. 2015;29(2):333-41.
10.1053/j.j.yca.2014.06.025

183. Åk K, Isbir CS, Tetik S, Atalan N, Tekeli A, Aljodi M, et al. Thromboelastographybased transfusion algorithm reduces blood product use after elective CABG: a prospective randomized study. *J Card Surg.* 2009;24(4):404-10. 10.1111/j.1540-8191.2009.00840.x

184. Avidan MS, Alcock EL, Da Fonseca J, Ponte J, Desai JB, Despotis GJ, et al. Comparison of structured use of routine laboratory tests or near-patient assessment with clinical judgement in the management of bleeding after cardiac surgery. *Br J Anaesth*. 2004;92(2):178-86. 10.1093/bja/aeh037

185. Cui Y, Hei F, Long C, Feng Z, Zhao J, Yan F, et al. Perioperative monitoring of thromboelastograph on blood protection and recovery for severely cyanotic patients undergoing complex cardiac surgery. *Artif Organs.* 2010;34(11):955-60. 10.1111/j.1525-1594.2010.01148.x

186. De Pietri L, Bianchini M, Montalti R, De Maria N, Di Maira T, Begliomini B, et al. Thrombelastography-guided blood product use before invasive procedures in cirrhosis with severe coagulopathy: A randomized, controlled trial. *Hepatology*. 2016;63(2):566-73. 10.1002/hep.28148

187. Girdauskas E, Kempfert J, Kuntze T, Borger MA, Enders J, Fassl J, et al. Thromboelastometrically guided transfusion protocol during aortic surgery with circulatory arrest: a prospective, randomized trial. *J Thorac Cardiovasc Surg.* 2010;140(5):1117-24 e2. 10.1016/j.jtcvs.2010.04.043

188. Karkouti K, Callum J, Wijeysundera DN, Rao V, Crowther M, Grocott HP, et al. Pointof-Care Hemostatic Testing in Cardiac Surgery: A Stepped-Wedge Clustered Randomized Controlled Trial. *Circulation*. 2016;134(16):1152-62. 10.1161/CIRCULATIONAHA.116.023956 189. Kultufan Turan S AB, Ayik H, Yaar S, Kazanci D, Karadenz Ü, et al. The role of rotational thromboelastgraphy on decision of blood transfusion in open heart surgery [Açik Kalp Cerrahisinde TromboelastografininTransfüzyon Karari Üzerine Etkisi]. *Göğüs-Kalp-Damar Anestezi ve Yoğun Bakım Derneği Dergisi*. 2006;12(4):154-9.

190. Nakayama Y, Nakajima Y, Tanaka KA, Sessler DI, Maeda S, Iida J, et al. Thromboelastometry-guided intraoperative haemostatic management reduces bleeding and red cell transfusion after paediatric cardiac surgery. *Br J Anaesth.* 2015;114(1):91-102. 10.1093/bja/aeu339

191. Rauter M KT, Schwarz S, Fitzgerald R. Reduced number of red blood cell transfusions in cardiac surgery due to perioperative coagulation testing by thromboelastometry. *Acta Anaesthesiologica Scandinavica*. 2007;51 (Suppl 118):40.

192. Royston D, von Kier S. Reduced haemostatic factor transfusion using heparinasemodified thrombelastography during cardiopulmonary bypass. *Br J Anaesth.* 2001;86(4):575-8. 10.1093/bja/86.4.575

193. Shore-Lesserson L, Manspeizer HE, DePerio M, Francis S, Vela-Cantos F, Ergin MA.
Thromboelastography-guided transfusion algorithm reduces transfusions in complex cardiac surgery. *Anesth Analg.* 1999;88(2):312-9. 10.1097/00000539-199902000-00016
194. Wang SC, Shieh JF, Chang KY, Chu YC, Liu CS, Loong CC, et al.

Thromboelastography-guided transfusion decreases intraoperative blood transfusion during orthotopic liver transplantation: randomized clinical trial. *Transplant Proc.* 2010;42(7):2590-3. 10.1016/j.transproceed.2010.05.144

195. Westbrook AJ, Olsen J, Bailey M, Bates J, Scully M, Salamonsen RF. Protocol based on thromboelastograph (TEG) out-performs physician preference using laboratory coagulation tests to guide blood replacement during and after cardiac surgery: a pilot study. *Heart Lung Circ.* 2009;18(4):277-88. 10.1016/j.hlc.2008.08.016

196. Schaden E, Kimberger O, Kraincuk P, Baron DM, Metnitz PG, Kozek-Langenecker S. Perioperative treatment algorithm for bleeding burn patients reduces allogeneic blood product requirements. *Br J Anaesth*. 2012;109(3):376-81. 10.1093/bja/aes186

197. Cao X, Zhang X, Li Q. [Efficacy of thromboelastography to monitor the clinical massive transfusion in scoliosis: a randomized controlled trial]. *Zhonghua Wai Ke Za Zhi*. 2016;54(2):137-41. 10.3760/cma.j.issn.0529-5815.2016.02.013

198. Rourke C, Curry N, Khan S, Taylor R, Raza I, Davenport R, et al. Fibrinogen levels during trauma hemorrhage, response to replacement therapy, and association with patient outcomes. *J Thromb Haemost.* 2012;10(7):1342-51. 10.1111/j.1538-7836.2012.04752.x

199. Johansson PI, Sorensen AM, Larsen CF, Windelov NA, Stensballe J, Perner A, et al. Low hemorrhage-related mortality in trauma patients in a Level I trauma center employing transfusion packages and early thromboelastography-directed hemostatic resuscitation with plasma and platelets. *Transfusion*. 2013;53(12):3088-99. 10.1111/trf.12214 200. Spiess BD, Gillies BS, Chandler W, Verrier E. Changes in transfusion therapy and reexploration rate after institution of a blood management program in cardiac surgical patients. *J Cardiothorac Vasc Anesth*. 1995;9(2):168-73. 10.1016/S1053-0770(05)80189-2

201. Anderson L, Quasim I, Soutar R, Steven M, Macfie A, Korte W. An audit of red cell and blood product use after the institution of thromboelastometry in a cardiac intensive care unit. *Transfus Med.* 2006;16(1):31-9. 10.1111/j.1365-3148.2006.00645.x

202. Spalding GJ, Hartrumpf M, Sierig T, Oesberg N, Kirschke CG, Albes JM. Cost reduction of perioperative coagulation management in cardiac surgery: value of "bedside" thrombelastography (ROTEM). *Eur J Cardiothorac Surg.* 2007;31(6):1052-7. 10.1016/j.ejcts.2007.02.022

203. Rahe-Meyer N, Winterhalter M, Boden A, Froemke C, Piepenbrock S, Calatzis A, et al. Platelet concentrates transfusion in cardiac surgery and platelet function assessment by multiple electrode aggregometry. *Acta Anaesthesiol Scand*. 2009;53(2):168-75. 10.1111/j.1399-6576.2008.01845.x

204. Görlinger K, Dirkmann D, Mueller-Beissenhirtz H, Paul A, Hartmann M, Saner F. Thromboelastometry-Based Perioperative Coagulation Management in Visceral Surgery and Liver Transplantation: Experience of 10 Years and 1105 LTX2010. S86-S p.

205. Noval-Padillo JA, Leon-Justel A, Mellado-Miras P, Porras-Lopez F, Villegas-Duque D, Gomez-Bravo MA, et al. Introduction of fibrinogen in the treatment of hemostatic disorders during orthotopic liver transplantation: implications in the use of allogenic blood. *Transplant Proc.* 2010;42(8):2973-4. 10.1016/j.transproceed.2010.08.011

206. Trzebicki J, Flakiewicz E, Kosieradzki M, Blaszczyk B, Kolacz M, Jureczko L, et al. The use of thromboelastometry in the assessment of hemostasis during orthotopic liver transplantation reduces the demand for blood products. *Ann Transplant*. 2010;15(3):19-24.

207. Gorlinger K, Dirkmann D, Hanke AA, Kamler M, Kottenberg E, Thielmann M, et al. First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: a retrospective, single-center cohort study. *Anesthesiology*. 2011;115(6):1179-91. 10.1097/ALN.0b013e31823497dd

208. Romlin BS, Wahlander H, Berggren H, Synnergren M, Baghaei F, Nilsson K, et al. Intraoperative thromboelastometry is associated with reduced transfusion prevalence in pediatric cardiac surgery. *Anesth Analg.* 2011;112(1):30-6. 10.1213/ANE.0b013e3181fe4674 209. Hvas AM, Boas TW, Jensen M, Lindskov C, Folkersen L, Ravn HB. Change in hemostatic intervention after implementation of thromboelastometry. *J Cardiothorac Vasc Anesth.* 2012;26(2):227-31. 10.1053/j.jvca.2011.07.018

210. Wang SC, Lin HT, Chang KY, Mandell MS, Ting CK, Chu YC, et al. Use of higher thromboelastogram transfusion values is not associated with greater blood loss in liver transplant surgery. *Liver Transpl.* 2012;18(10):1254-8. 10.1002/lt.23494

211. Leon-Justel A, Noval-Padillo JA, Alvarez-Rios AI, Mellado P, Gomez-Bravo MA, Alamo JM, et al. Point-of-care haemostasis monitoring during liver transplantation reduces transfusion requirements and improves patient outcome. *Clin Chim Acta.* 2015;446:277-83. 10.1016/j.cca.2015.04.022

212. Roullet S, Freyburger G, Cruc M, Quinart A, Stecken L, Audy M, et al. Management of bleeding and transfusion during liver transplantation before and after the introduction of a rotational thromboelastometry-based algorithm. *Liver Transpl.* 2015;21(2):169-79. 10.1002/lt.24030

213. Bedreli S, Sowa JP, Gerken G, Saner FH, Canbay A. Management of acute-onchronic liver failure: rotational thromboelastometry may reduce substitution of coagulation factors in liver cirrhosis. *Gut.* 2016;65(2):357-8. 10.1136/gutjnl-2015-309922 214. De Pietri L, Ragusa F, Deleuterio A, Begliomini B, Serra V. Reduced Transfusion During OLT by POC Coagulation Management and TEG Functional Fibrinogen: A Retrospective Observational Study. *Transplantation direct.* 2016;2(1):e49. 10.1097/txd.00000000000559

215. St-Onge S, Lemoine E, Bouhout I, Rochon A, El-Hamamsy I, Lamarche Y, et al. Evaluation of the real-world impact of rotational thromboelastometry-guided transfusion protocol in patients undergoing proximal aortic surgery. *J Thorac Cardiovasc Surg.* 2018. 10.1016/j.jtcvs.2018.07.043

216. Kuiper G, van Egmond LT, Henskens YMC, Roekaerts PM, Maessen JG, Ten Cate H, et al. Shifts of Transfusion Demand in Cardiac Surgery After Implementation of Rotational Thromboelastometry-Guided Transfusion Protocols: Analysis of the HEROES-CS (HEmostasis Registry of patiEntS in Cardiac Surgery) Observational, Prospective Open Cohort Database. *J Cardiothorac Vasc Anesth.* 2019;33(2):307-17. 10.1053/j.jvca.2018.08.203

217. Mallaiah S, Barclay P, Harrod I, Chevannes C, Bhalla A. Introduction of an algorithm for ROTEM-guided fibrinogen concentrate administration in major obstetric haemorrhage. *Anaesthesia*. 2015;70(2):166-75. 10.1111/anae.12859

218. Collins PW, Cannings-John R, Bruynseels D, Mallaiah S, Dick J, Elton C, et al. Viscoelastometry guided fresh frozen plasma infusion for postpartum haemorrhage: OBS2, an observational study. *Br J Anaesth.* 2017;119(3):422-34. 10.1093/bja/aex245

219. Xu L, Wang L, Yang X, Li K, Sun H, Zhang D, et al. Platelet function monitoring guided antiplatelet therapy in patients receiving high-risk coronary interventions. *Chinese medical journal*. 2014;127(19):3364-70.

220. Gorlinger K, Fries D, Dirkmann D, Weber CF, Hanke AA, Schochl H. Reduction of Fresh Frozen Plasma Requirements by Perioperative Point-of-Care Coagulation Management with Early Calculated Goal-Directed Therapy. *Transfus Med Hemother*. 2012;39(2):104-13. 10.1159/000337186

221. Bugaev N, Como JJ, Golani G, Freeman JJ, Sawhney JS, Vatsaas CJ, et al. Thromboelastography and rotational thromboelastometry in bleeding patients with coagulopathy: Practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg.* 2020;89(6):999-1017. 10.1097/TA.0000000000002944

222. Amgalan A, Allen T, Othman M, Ahmadzia HK. Systematic review of viscoelastic testing (TEG/ROTEM) in obstetrics and recommendations from the women's SSC of the ISTH. *J Thromb Haemost*. 2020;18(8):1813-38. 10.1111/jth.14882

223. Gonzalez E, Moore EE, Moore HB, Chapman MP, Chin TL, Ghasabyan A, et al. Goaldirected Hemostatic Resuscitation of Trauma-induced Coagulopathy: A Pragmatic Randomized Clinical Trial Comparing a Viscoelastic Assay to Conventional Coagulation Assays. *Annals of surgery*. 2016;263(6):1051-9. 10.1097/SLA.000000000001608

224. Kempfert. J. Thromboelastography-guided blood component therapy after cardiac surgery: A randomised study. *Interactive Cardiovascular and Thoracic Surgery*. 2011;13:S106–7.

225. NCT00772239. Perioperative coagulation management in cardiac surgery (ROTEM). (first recieved 13 October 2008). [Accessed: 26 November 2019]. Available from: clinicaltrials.gov/ct2/show/NCT00772239.

226. Nuttall GA, Oliver WC, Santrach PJ, Bryant S, Dearani JA, Schaff HV, et al. Efficacy of a simple intraoperative transfusion algorithm for nonerythrocyte component utilization after cardiopulmonary bypass. *Anesthesiology*. 2001;94(5):773-81; discussion 5A-6A. 10.1097/00000542-200105000-00014

227. Paniagua P, Koller T, Requena T, Gil J, Campos J, Galan J. Randomized controlled trial to evaluate postoperative coagulation management with bed-side trombelastometry (Rotem) compared with a transfusion protocol based on laboratory measurements in bleeding patients after cardiac surgery: Preliminary data. *European Journal of Anaesthesiology*. 2011;28:94.

228. Weber CF, Gorlinger K, Meininger D, Herrmann E, Bingold T, Moritz A, et al. Pointof-care testing: a prospective, randomized clinical trial of efficacy in coagulopathic cardiac surgery patients. *Anesthesiology*. 2012;117(3):531-47. 10.1097/ALN.0b013e318264c644 229. Baksaas-Aasen K, Gall L, Eaglestone S, Rourke C, Juffermans NP, Goslings JC, et al. iTACTIC - implementing Treatment Algorithms for the Correction of Trauma-Induced

Coagulopathy: study protocol for a multicentre, randomised controlled trial. *Trials*. 2017;18(1):486. 10.1186/s13063-017-2224-9

230. Baksaas-Aasen K, Gall LS, Stensballe J, Juffermans NP, Curry N, Maegele M, et al. Viscoelastic haemostatic assay augmented protocols for major trauma haemorrhage (ITACTIC): a randomized, controlled trial. *Intensive Care Med*. 2021;47(1):49-59. 10.1007/s00134-020-06266-1

Hanke AA, Herold U, Dirkmann D, Tsagakis K, Jakob H, Gorlinger K.
Thromboelastometry Based Early Goal-Directed Coagulation Management Reduces Blood Transfusion Requirements, Adverse Events, and Costs in Acute Type A Aortic Dissection: A Pilot Study. *Transfus Med Hemother*. 2012;39(2):121-8. 10.1159/000337723
Kashuk JL, Moore EE, Wohlauer M, Johnson JL, Pezold M, Lawrence J, et al. Initial experiences with point-of-care rapid thrombelastography for management of life-

threatening postinjury coagulopathy. *Transfusion*. 2012;52(1):23-33. 10.1111/j.1537-2995.2011.03264.x

233. FassI J, Matt P, Eckstein F, Filipovic M, Gregor M, Zenklusen U, et al. Transfusion of allogeneic blood products in proximal aortic surgery with hypothermic circulatory arrest: effect of thromboelastometry-guided transfusion management. *J Cardiothorac Vasc Anesth*. 2013;27(6):1181-8. 10.1053/j.jvca.2013.02.009

234. Tapia NM, Chang A, Norman M, Welsh F, Scott B, Wall MJ, Jr., et al. TEG-guided resuscitation is superior to standardized MTP resuscitation in massively transfused penetrating trauma patients. *J Trauma Acute Care Surg.* 2013;74(2):378-85; discussion 85-6. 10.1097/TA.0b013e31827e20e0

235. Nardi G, Agostini V, Rondinelli B, Russo E, Bastianini B, Bini G, et al. Traumainduced coagulopathy: impact of the early coagulation support protocol on blood product consumption, mortality and costs. *Crit Care*. 2015;19(1):83. 10.1186/s13054-015-0817-9 236. Barinov SV, Zhukovsky YG, Dolgikh VT, Medyannikova IV. Novel combined strategy of obstetric haemorrhage management during caesarean section using intrauterine balloon tamponade. *J Matern Fetal Neonatal Med*. 2017;30(1):29-33. 10.3109/14767058.2015.1126242

237. Prat NJ, Meyer AD, Ingalls NK, Trichereau J, DuBose JJ, Cap AP. Rotational thromboelastometry significantly optimizes transfusion practices for damage control resuscitation in combat casualties. *J Trauma Acute Care Surg.* 2017;83(3):373-80. 10.1097/TA.00000000001568

238. Snegovskikh D, Souza D, Walton Z, Dai F, Rachler R, Garay A, et al. Point-of-care viscoelastic testing improves the outcome of pregnancies complicated by severe postpartum hemorrhage. *J Clin Anesth*. 2018;44:50-6. 10.1016/j.jclinane.2017.10.003 239. Guth C, Vassal O, Friggeri A, Wey PF, Inaba K, Decullier E, et al. Effects of modification of trauma bleeding management: A before and after study. *Anaesth Crit Care Pain Med*. 2019;38(5):469-76. 10.1016/j.accpm.2019.02.005

240. McNamara H, Kenyon C, Smith R, Mallaiah S, Barclay P. Four years' experience of a ROTEM((R)) -guided algorithm for treatment of coagulopathy in obstetric haemorrhage. *Anaesthesia*. 2019;74(8):984-91. 10.1111/anae.14628

241. Unruh M, Reyes J, Helmer SD, Haan JM. An evaluation of blood product utilization rates with massive transfusion protocol: Before and after thromboelastography (TEG) use in trauma. *Am J Surg.* 2019;218(6):1175-80. 10.1016/j.amjsurg.2019.08.027

242. Wang H, Robinson RD, Phillips JL, Ryon A, Simpson S, Ford JR, et al. Traumatic Abdominal Solid Organ Injury Patients Might Benefit From Thromboelastography-Guided Blood Component Therapy. *J Clin Med Res.* 2017;9(5):433-8. 10.14740/jocmr3005w

243. Nayar SK, Shafiq B. Blood Conservation in Orthopedic Trauma. *Techniques in Orthopaedics*. 2017;32(1):45-50. <u>http://dx.doi.org/10.1097/BTO.0000000000000208</u>

244. Shantikumar S, Patel S, Handa A. The role of cell salvage autotransfusion in abdominal aortic aneurysm surgery. *European Journal of Vascular & Endovascular Surgery*. 2011;42(5):577-84. <u>https://dx.doi.org/10.1016/j.ejvs.2011.04.014</u>

245. Meybohm P, Choorapoikayil S, Wessels A, Herrmann E, Zacharowski K, Spahn DR. Washed cell salvage in surgical patients: A review and meta-analysis of prospective randomized trials under PRISMA. *Medicine*. 2016;95(31):e4490.

https://dx.doi.org/10.1097/MD.000000000004490

246. Li J, Sun LS, Tian HJ, Yang K, Liu R, Li J. Cell salvage in emergency trauma surgery. *Cochrane Database of Systematic Reviews*. 2015(1).

247. Bowley DM, Barker P, Boffard KD. Intraoperative blood salvage in penetrating abdominal trauma: a randomised, controlled trial. *World J Surg.* 2006;30(6):1074-80. 10.1007/s00268-005-0466-2

248. Bhangu A, Nepogodiev D, Doughty H, Bowley DM. Intraoperative cell salvage in a combat support hospital: a prospective proof of concept study. *Transfusion*. 2013;53(4):805-10. 10.1111/j.1537-2995.2012.03835.x

249. Markovic M, Davidovic L, Savic N, Sindjelic R, Ille T, Dragas M. Intraoperative cell salvage versus allogeneic transfusion during abdominal aortic surgery: clinical and financial outcomes. *Vascular*. 2009;17(2):83-92. 10.2310/6670.2009.00009

250. Tawfick WA, O'Connor M, Hynes N, Sultan S. Implementation of the Continuous AutoTransfusion System (C.A.T.S) in open abdominal aortic aneurysm repair: an observational comparative cohort study. *Vasc Endovascular Surg.* 2008;42(1):32-9. 10.1177/1538574407309316

251. Serracino-Inglott F, Awad S, Barclay A, Nasim A. The use of a cell saver during repair of ruptured abdominal aortic aneurysms increases early survival. *Ann R Coll Surg Engl.* 2005;87(6):475.

252. Shuhaiber JH, Whitehead SM. The impact of introducing an autologous intraoperative transfusion device to a community hospital. *Ann Vasc Surg.* 2003;17(4):424-9. 10.1007/s10016-003-0030-1

253. Posacioglu H, Apaydin AZ, Islamoglu F, Calkavur T, Yagdi T, Atay Y, et al. Adverse effects of cell saver in patients undergoing ruptured abdominal aortic aneurysm repair. *Ann Vasc Surg.* 2002;16(4):450-5. 10.1007/s10016-001-0123-7

Appendix A – Literature search results

See Volume 2 of the Technical Report

Appendix B – Literature screening results

See Volume 2 of the Technical Report

Appendix C – List of excluded studies

See Volume 2 of the Technical Report

Appendix D – Critical appraisal

See Volume 2 of the Technical Report

Appendix E – Data extraction forms

See Volume 3 of the Technical Report