



NATIONAL BLOOD AUTHORITY
AUSTRALIA

AUSTRALIAN BLEEDING DISORDERS REGISTRY

Annual Report 2018-19





With the exception of any logos and registered trademarks, and where otherwise noted, all material presented in this document is provided under a Creative Commons Attribution 4.0 license (<https://creativecommons.org/licenses/by/4.0/>)

The details of the relevant license conditions are available on the Creative Commons website (accessible using the links provided) as is the full legal code for the CC BY 4.0 license (<https://creativecommons.org/licenses/by/4.0/legalcode>)

The content obtained from this document or derivative of this work must be attributed as:

Australian Bleeding Disorders Registry (ABDR) Annual Report 2018-19 published by the National Blood Authority.

ISSN 1839-0811 (online version)

This report is available online at <http://www.blood.gov.au/data-analysis-reporting>

Version: 18 December 2019



Locked Bag 8430
Canberra ACT 2601
Phone: 13 000 BLOOD (13000 25663)
Email: data@blood.gov.au
www.blood.gov.au

Table of Contents

List of Tables	6
List of Figures.....	7
PURPOSE OF THIS DOCUMENT	8
KEY FINDINGS	9
BACKGROUND	11
What are bleeding disorders?	11
Bleeding disorders are inherited or acquired.....	11
Haemophilia	11
Types of haemophilia	12
Haemophilia fast facts.....	12
Von willebrand disorder/disease (VWD).....	12
Types of VWD	13
Rare clotting factor deficiencies.....	13
Special issues for girls and women.....	13
Inherited platelet disorders.....	14
What are platelet function disorders?	14
Severity.....	14
Treatment of bleeding disorders.....	15
TREATMENT OF BLEEDING DISORDERS IN AUSTRALIA	16
THE AUSTRALIAN BLEEDING DISORDERS REGISTRY (ABDR).....	17
ABDR management and governance.....	17
Patient privacy in ABDR and MyABDR.....	18
Data governance.....	18
Data quality issues.....	18
ABDR system.....	19
Comparing data from previous ABDR annual reports.....	19
Consistent application of diagnoses and definitions.....	19
Von willebrand disease.....	19
Treatments not included in the ABDR.....	19

Consent.....	19
SUPPLY OF PRODUCTS FOR TREATMENT.....	20
Extended Half Life (EHL) products.....	20
Patient suitability and prioritisation.....	21
ABDR PATIENT DEMOGRAPHICS.....	22
Diagnoses.....	22
Patients with multiple bleeding disorders.....	23
Age, diagnosis and severity	26
By age group and detailed diagnosis.....	29
By location	31
By gender and age distribution	32
Inhibitor status	44
Incidence of major disorders.....	47
PATIENT TREATMENT IN 2018-19.....	48
Products issued to patients	48
Volume (IU) of products issued for HMA and HMB	51
Volume of products issued and patient counts by treatment regimen, severity, product and state.....	55
APPENDIX A CHARACTERISTICS OF RARE CLOTTING FACTOR DEFICIENCIES	67
APPENDIX B HAEMOPHILIA TREATMENT CENTRES	68
The objectives of HTC's	68
Operating concept.....	68
Data quality of HTC data collections	69
Data Projects	70
List of HTC's	71
APPENDIX C NATIONAL SUPPLY OF PRODUCTS	72
National supply plan and budget	72
Issues of clotting factors.....	73
Supply of extended half life products.....	76
Patient suitability and prioritisation.....	76
Priority Criteria	76
Other considerations.....	77
Chronology of products supplied	78

APPENDIX D HISTORY OF THE ABDR	79
Benefits of the 4 th generation ABDR.....	80
Current position of the development of the ABDR.....	80
APPENDIX E PATIENT REGISTRATION FORM.....	81
ACRONYMS AND GLOSSARY OF TERMS.....	84
Acronyms.....	84
Glossary of terms.....	84

LIST OF TABLES

Table 1 - Major bleeding disorders and their cause	11
Table 2 - Severity and the concentration of clotting factors	14
Table 3 - Number of people in the registry and treated by broad diagnosis	22
Table 4 - Number of hereditary patients who received Extended Half Life Products by broad diagnosis and financial year	23
Table 5 - Number of people in the registry with multiple bleeding disorders	23
Table 6 - Number of people in the registry and treated by detailed diagnosis	24
Table 7 - Number of adult patients in the registry and treated by broad diagnosis and severity for HMA, HMB	27
Table 8 - Number of paediatric and adolescent patients in the registry and treated by broad diagnosis and severity for HMA, HMB.....	28
Table 9 - Number of people in the registry diagnosed with HMA or HMB by age group and disease classification	29
Table 10 - Number of people in the registry diagnosed with VWD by age group and disease classification	30
Table 11 - Numbers of patients with severe HMA and HMB by location	31
Table 12 - Patient numbers by Bleeding Disorder and gender	32
Table 13 - Patient numbers by Bleeding Disorder and Age Group.....	34
Table 14 - Hereditary HMA, HMB and VWD patients by gender and age range at 30 June 2019	35
Table 15 - Description of inhibitor status used in ABDR	45
Table 16 - Patient inhibitor status numbers.....	46
Table 17 - Incidence statistics from World Federation of Hemophilia Global Survey 2018.....	47
Table 18 - Volume (IU) of product issued for HMA, HMB and VWD patients, by severity and treatment regimen in 2018-19 - hereditary bleeding disorders.....	49
Table 19 - Volume (IU) of product issued for HMA, HMB and VWD patients over time, including EHL products 2014-15 to 2018-19 - hereditary bleeding disorders	50
Table 20 - Volume (IU) of product issued for HMA, HMB and VWD patients, by severity and treatment regimen in 2018-19 - acquired bleeding disorders	50
Table 21 - Volume (IU) of products issued for other patients, by severity and treatment regimen in 2018-19 - other diagnoses.....	50
Table 22 - Volume (IU) of products issued in 2018-19 (including EHL products) by treatment regimen - hereditary bleeding disorders	56
Table 23 - Patient numbers and Volume (IU) of products issued in 2018-19 by treatment regimen - hereditary bleeding disorders – Adult patients.....	57
Table 24 - Patient numbers and Volume (IU) of products issued in 2018-19 by treatment regimen - Acquired bleeding disorders.....	59
Table 25 - Volume (IU) of products issued in 2018-19 by treatment regimen - other diagnoses.....	60
Table 26 - Number of patients for hereditary HMA, HMB and VWD by state	61
Table 27 - Volume (IU) of product issued for hereditary HMA, HMB and VWD by state	62
Table 28 - Volume (IU), patient counts and IU or mg/kg/year of products issued in 2018-19 by treatment regimen - Hereditary	63
Table 29 - Volume (IU) of product issued and patient counts for hereditary HMA and HMB by severity and regimen type	64
Table 30 - Volume (IU) of product issued and patient counts for hereditary HMA, HMB and VWD by regimen type and product.....	65
Table 31 - Characteristics of rare clotting factor deficiencies	67
Table 32 - Haemophilia Treatment Centres	71

LIST OF FIGURES

Figure 1 - Location of haemophilia treatment centres.....	16
Figure 2 - Market share of recombinant FVIII issues 2014-15 to 2018-19	20
Figure 3 - Total Units (IU) of extended half life products issued 2018-19.....	21
Figure 4 - Numbers of active patients in the Registry as at 30 June 2019	31
Figure 5 - Distribution of hereditary female HMA patients by age in 2018-19.....	36
Figure 6 - Distribution of hereditary female HMB patients by age in 2018-19	37
Figure 7 - Distribution of hereditary male HMA patients by age in 2018-19	38
Figure 8 - Distribution of hereditary male HMA severe patients by age in 2018-19	39
Figure 9 - Distribution of hereditary male HMB patients by age in 2018-19	40
Figure 10 - Distribution of hereditary male HMB severe patients by age in 2018-19.....	41
Figure 11 - Distribution of hereditary female VWD patients by age in 2018-19.....	42
Figure 12 - Distribution of hereditary male VWD patients by age in 2018-19	43
Figure 13 - Percentage of patients receiving product by severity for HMA - hereditary bleeding disorders.....	48
Figure 14 - Percentage of patients receiving product by severity for HMB - hereditary bleeding disorders.....	49
Figure 15 - FVIII Product usage (IU/kg/year) in severe HMA patients on prophylaxis.....	51
Figure 16 - FVIII Product usage (IU/kg/year) in severe HMA patients on demand.....	52
Figure 17 - FIX Product usage (IU/kg/year) in severe HMB patients on prophylaxis	53
Figure 18 - FIX Product usage (IU/kg/year) in severe HMB patients on demand	54
Figure 19 - National issues by product category 2018-19.....	72
Figure 20 - Expenditure on clotting factors and percentage of blood budget 2009-10 to 2018-19	73
Figure 21 - Issues of factor VIII products, 2014-15 to 2018-19 per '000 population	74
Figure 22 - Issues of factor IX products, 2014-15 to 2018-19 per '000 population	74
Figure 23 - Issues of recombinant factor VIIa products, 2014-15 to 2018-19 per '000 population.....	75
Figure 24 - Issues of FEIBA, 2014-15 to 2018-19 per '000 population	75
Figure 25 - Uptake of EHL product use December 2017- July 2019	76

Purpose of this document

The intention of this document is to present the reader with an integrated view of current clinical and demographic information on people with inherited bleeding disorders in Australia and the resultant demand for clotting factor products. It draws on data from the Australian Bleeding Disorders Registry (ABDR) and other National Blood Authority (NBA) supply and contract sources. Some international data comparisons have also, where meaningful, been included.

The ABDR is a clinical registry for patients in Australia with bleeding disorders. It is used on a daily basis by clinicians in all Australian Haemophilia Treatment Centres (HTCs) to assist in managing the treatment of people with bleeding disorders and to gain a better understanding of the incidence and prevalence of bleeding disorders. This information will also be used by the NBA to understand demand for, and to facilitate ordering of, clotting factor product.

This document will be used by people involved in providing care for patients with bleeding disorders, and may also be useful for patient advocacy groups and those in administrative and government positions.

Key findings

Total patients in ABDR 2018-19

- 6,355 patients
- 5,151 with hereditary HMA, HMB or VWD
- 111 with acquired HMA, HMB or VWD
- 1,093 with other bleeding disorders
- 1,804 received product

HMA

- 2,372 hereditary HMA patients (668 severe)
- 1,104 patients received product
- 187,097,550 IU of Factor VIII products used by hereditary HMA patients (154,947,300 IU standard half life (SHL) and 32,150,250 IU extended half life (EHL))
- 125,028,250 IU (80.7%) SHL and 31,745,750 (98.7%) EHL for prophylactic use

HMB

- 558 hereditary HMB patients (109 severe)
- 247 patients received product
- 27,714,752 IU of Factor IX products used by hereditary HMB patients (16,845,502 IU SHL and 10,869,250 EHL)
- 8,813,000 IU (52.3%) SHL and 10,298,000 (94.7%) for prophylactic use

VWD

- 2,221 hereditary VWD patients
- 307 patients received product
- 7,352,750 IU of Factor VIII products used by hereditary VWD patients
- 3,899,250 (53.0%) for prophylactic use

Other bleeding disorders

- 1,093 patients with other bleeding disorders
- 121 patients received product
- 324,780 IU of product used by patients with other bleeding disorders
- 260,148 (74.9%) for prophylactic use

Demand for clotting factors in 2018-19

- 13.7% of total blood and blood product expenditure in 2018-19 compared with 14% in 2017-18

Factor VIII

- Demand for Factor VIII products increased by 6.7% compared with 2017-18
- Recombinant FVIII increased by 8.3%
- Plasma derived FVIII decreased by 3.6%

Factor IX

- Demand for Factor IX decreased by 12.6% compared to 2017-18
- Plasma derived FIX decreased by 1.3% due to a reduction in specific patient requirements
- Recombinant FIX decreased 12.8% compared to 2017-18

Source: NBA Annual Report 2018-19

Background

The information in this section has been drawn from the materials and websites of two peak bodies for haemophilia; the World Federation of Hemophilia (www.wfh.org) and the Haemophilia Foundation Australia (www.haemophilia.org.au).

WHAT ARE BLEEDING DISORDERS?

In people with bleeding disorders, the clotting process doesn't work properly. As a result, people with bleeding disorders can bleed for longer than normal, and some may experience spontaneous bleeding into joints, muscles, or other parts of their bodies.

BLEEDING DISORDERS ARE INHERITED OR ACQUIRED

Bleeding disorders are almost always inherited or passed through families; they have a genetic basis and the genes responsible for the disorders are passed from parents to children. However, a person can also spontaneously develop a bleeding disorder, although this is rare.

Acquired bleeding disorders are not inherited or passed through families. Most acquired bleeding disorders have an identifiable root cause. Men and women are equally likely to be affected by an acquired bleeding disorder, and the potential for problems is high.

TABLE 1 - MAJOR BLEEDING DISORDERS AND THEIR CAUSE

Disorder group	Cause
Haemophilia A	Deficiency of Factor VIII
Haemophilia B	Deficiency of Factor IX
von Willebrand Disease	Deficiency, or dysfunction, of von Willebrand Factor
Other Factor deficiencies	Deficiency of other coagulation factors
Platelet Disorder	Inherited deficiency of effective platelet function

HAEMOPHILIA

Haemophilia causes excessive bleeding following trauma or surgery and can be related to spontaneous haemorrhages into muscles and joints. People with haemophilia do not bleed any faster than normal, but they can bleed for a longer time.

Haemophilia is an X-linked disorder that typically affects males, whereas females are normally classified as carriers. However, affected males will pass on the haemophilia gene to their daughters, and women carrying a F8 or F9 gene mutation may have reduced factor levels and should therefore be classified as having haemophilia. Most carriers are asymptomatic. Carriers with clotting factor levels in the haemophilia range may be symptomatic, with bleeding manifestations commensurate with their degree of clotting factor deficiency, particularly during trauma and surgery. Symptomatic carriers are classified as haemophilia in line with the World Federation of Hemophilia (www.wfh.org) guidelines.

TYPES OF HAEMOPHILIA

- The most common type of haemophilia is called Haemophilia A. This means the person does not have enough clotting Factor VIII (factor eight).
- Haemophilia B is less common. A person with Haemophilia B does not have enough Factor IX (factor nine). The symptoms are the same for people with Haemophilia A and B; that is, they bleed for a longer time than normal.

HAEMOPHILIA FAST FACTS

- Haemophilia occurs in 1 in 6,000-10,000 males internationally.
- Currently in Australia there are 3,009 people with Haemophilia A and B, (including 79 with Acquired Haemophilia) with varied degrees of severity, in the Australian Bleeding Disorders Registry (ABDR).
- Bleeding is most commonly internal into the joints and/or muscles. Less commonly, bleeding into internal organs can also occur. It can happen without an obvious cause (sometimes called 'spontaneous'), or as a result of injury.
- Over time this internal bleeding into joints ('bleeds') can cause severe arthritis, chronic pain and disability.
- Specialised treatment is needed to help blood clot normally. With appropriate treatment haemophilia can be managed effectively.
- Haemophilia is an inherited condition and occurs in families; however in 1/3 of cases it appears in families with no previous history of the disorder. The haemophilia gene is passed down from parent to child through generations. Men with haemophilia will pass the gene on to their daughters but not their sons. Women who carry the haemophilia gene can pass the haemophilia gene on to their sons and daughters. Sons with the gene will have haemophilia. Some women and girls who carry the gene may also experience bleeding problems.

VON WILLEBRAND DISORDER/DISEASE (VWD)

Von Willebrand disease (VWD) is the most common type of bleeding disorder. People with VWD have a problem with von Willebrand Factor (VWF), a protein in their blood that would normally help control bleeding. When a blood vessel is injured and bleeding occurs, VWF helps cells in the blood, called platelets, adhere to damaged blood vessels and mesh together and form a clot to stop the bleeding. People with VWD do not have enough VWF, or it does not work the way it should. It takes longer for blood to clot and for bleeding to stop.

VWD is generally less severe than other bleeding disorders. Many people with VWD may not know that they have the disorder because their bleeding symptoms are very mild. For most people with VWD, the disorder causes little or no disruption to their lives except when there is a serious injury or need for surgery. However, with all forms of VWD, there can be bleeding problems. VWD is difficult to accurately diagnose as laboratory values can fluctuate and values in those with mild bleeding symptoms can overlap with normal laboratory values.

From some studies, it is estimated that up to 1% of the world's population has VWD, but because many people have only very mild symptoms, only a small number of them are diagnosed. Research has shown that as many as 9 out of 10 people with VWD have not been diagnosed. It is estimated that VWD affects approximately 200,000 people in Australia, but symptomatic individuals possibly less. Currently there are 2,253 people with VWD in the ABDR including 32 with acquired VWD.

TYPES OF VWD

There are three main types of VWD. Bleeding symptoms can be quite variable within each type depending in part on the VWF activity. It is important to know which type of VWD a person has, because treatment is different for each type.

- Type 1 VWD is the most common form. People with Type 1 VWD have lower than normal levels of VWF. Symptoms are usually mild. Still, it is possible for someone with Type 1 VWD to have serious bleeding.
- Type 2 VWD involves a defect in the VWF structure. The VWF protein does not work properly, causing lower than normal VWF activity. There are different Type 2 VWD defects. Severity of symptoms can vary.
- Type 3 VWD is usually the most serious form. People with Type 3 VWD have very little or no VWF. Symptoms are more severe. People with Type 3 VWD can have bleeding into muscles and joints, sometimes without injury.

RARE CLOTTING FACTOR DEFICIENCIES

Rare clotting factor deficiencies are a group of inherited bleeding disorders caused by a problem with one of several clotting factors. Clotting factors are proteins in the blood that control bleeding. Many different clotting factors work together in a series of chemical reactions to stop bleeding. This is called the clotting process.

Problems with Factor VIII and Factor IX are known as Haemophilia A and B, respectively. Rare clotting factor deficiencies are bleeding disorders in which one of the other clotting factors (i.e. factors I, II, V, V+VIII, VII, X, XI, or XIII) is missing or not working properly. The World Federation of Hemophilia produced a summary (Table 25) of the characteristics of rare clotting factor deficiencies, the severity of bleeds associated with them, and the treatment typically required.

SPECIAL ISSUES FOR GIRLS AND WOMEN

Women with clotting factor deficiencies may have additional symptoms because of menstruation and childbirth. Girls may have especially heavy bleeding when they begin to menstruate. Women with clotting factor deficiencies may have heavier and/or longer menstrual flow, which can cause anaemia (with low levels of iron, which results in weakness and fatigue). Women with clotting factor deficiencies should receive genetic counselling about the risks of having an affected child well in advance of any planned pregnancies and should see an obstetrician as soon as they suspect they are pregnant. The obstetrician should work closely with the staff of the haemophilia/bleeding disorder treatment centre in order to provide the best care during pregnancy and childbirth and to minimize the potential complications for both the mother and the newborn child.

Women with certain rare factor deficiencies (such as Factor XIII deficiency and afibrinogenemia) may be at greater risk of miscarriage and placental abruption (a premature separation of the placenta from the uterus that disrupts the flow of blood and oxygen to the foetus). Therefore, these women require treatment throughout the pregnancy to prevent these complications.

The main risk related to pregnancy is postpartum haemorrhage. All bleeding disorders are associated with a greater risk of increased bleeding after delivery. The risk and the severity of the bleeding can be reduced with appropriate treatment. This treatment is different for each woman and depends on her personal and family history of bleeding symptoms, the severity of the factor deficiency, and the mode of delivery (vaginal birth vs. caesarean section). Factor replacement may be necessary in some cases.

INHERITED PLATELET DISORDERS

Platelets are small parts of cells that circulate in the blood. They are involved in the formation of blood clots and the repair of damaged blood vessels.

When a blood vessel is injured, platelets stick to the damaged area and spread along the surface to stop the bleeding (this process is called adhesion). At the same time, chemical signals are released from small sacks inside the platelets called granules (this process is called secretion). These chemicals attract other platelets to the site of injury and make them clump together to form what is called a platelet plug (this process is called aggregation).

Sometimes the platelet plug is enough to stop the bleeding. However if the wound is large, other proteins called clotting factors are recruited to the site of injury. These clotting factors work together on the surface of the platelets to form and strengthen the blood clot.

WHAT ARE PLATELET FUNCTION DISORDERS?

Platelet function disorders are conditions in which platelets don't work the way they should, resulting in a tendency to bleed or bruise. Since the platelet plug does not form properly, bleeding can continue for longer than normal. Since platelets have many roles in blood clotting, platelet function disorders can lead to bleeding disorders of various intensities.

SEVERITY

Haemophilia A and B are classified according to their severity, as this informs the treatment regimens required. The definitions of severity that are applied within the ABDR are listed in Table 2. Definition of severity of VWD and other coagulation factor deficiencies is not standardised but variable.

TABLE 2 - SEVERITY AND THE CONCENTRATION OF CLOTTING FACTORS¹

Severity	Concentration of Clotting Factor	Typical Bleeding Picture
Severe	<0.01 IU/ml (<1% of normal [†])	Frequent bleeding episodes common, predominantly into joints & muscles. Bleeding can occur spontaneously or after minor injury.
Moderate	0.01 – 0.05 IU/ml (1–5% of normal)	Can bleed after minor injury. May have joint bleeding. Severe bleeding with trauma, surgery, invasive procedures.
Mild	>0.05 – 0.40 IU/ml (5-40% of normal) [‡]	Spontaneous bleeding does not occur. Bleeding with major trauma, surgery, invasive procedures.

Notes † Normal concentration of Factor VIII or IX is defined as 100% or one unit of Factor VIII activity per ml of plasma - 100 U/dL (Kasper, CK 2004, Hereditary plasma clotting factor disorders and their management. Treatment of Hemophilia Monograph Series, No. 4, World Federation of Hemophilia, Montreal, Canada)
‡ Levels of FVIII above 40% are usually considered sufficient for normal haemostasis

¹ Modified from Srivastava A, Brewer AK, Mauser-Bunschoten EP, Key NS, Kitchen S, Llinas A, Ludlam CA, Mahlangu JN, Mulder K, Poon MC, Street A; Treatment Guidelines Working Group on Behalf of The World Federation Of Hemophilia (2013). Guidelines for the management of hemophilia, Haemophilia 19(1):e1-47.

TREATMENT OF BLEEDING DISORDERS

Mild conditions may require no treatment or treatment only under special circumstances, such as surgery. More severe conditions may require regular interventions. Treatment may occur in hospital or other medical facilities, or at home. The treatments may be regular and preventative (prophylaxis), or on demand (when a bleed occurs). In some patients, therapy is complicated when their body develops inhibitors that destroy the replacement clotting factors and other treatment is necessary.

Often the treatments involve providing replacement for the missing or defective clotting factors. Products used include plasma derived and recombinant clotting factors, cryoprecipitate and Desmopressin (1-desamino-8-D-arginine vasopressin; DDAVP) which can stimulate the release of Factor VIII and VWF from stores in the body (this is not used in Haemophilia B or Factor IX deficiency).

Treatment of bleeding disorders in Australia

The majority of people with these conditions are treated at HTC which are specialist centres that provide comprehensive care to people with haemophilia and other bleeding disorders. The comprehensive care model ensures that preventative and general treatment on the complex aspects of haemophilia are given in a co-ordinated way by a multi-disciplinary team with specialised expertise within the one centre.

HTCs were established following a decision by Australian Health Ministers Advisory Council (AHMAC) in 1998, to provide a leadership role within their hospital, city and outlying areas to ensure optimal care and an equitable distribution of professional and therapeutic resources, together with responsible record-keeping. The roles, aims and governance of these Centres are defined in [Appendix B](#). The locations of the HTCs in Australia are shown in Figure 1.



FIGURE 1 - LOCATION OF HAEMOPHILIA TREATMENT CENTRES

The model for HTCs varies between jurisdictions in relation to centralisation of services, size and age of patient population.

There are also some patients whose treatment is managed by clinicians who are not associated with a HTC. The proportion of product that is used in these circumstances varies across jurisdictions and there is some variability in the data capture for this activity between jurisdictions. Accordingly, data on total volume of products recorded from the ABDR may not be consistent with data from other sources.

The Australian Bleeding Disorders Registry (ABDR)

The Australian Bleeding Disorders Registry (ABDR) is a database that is designed to collect all clinical information related to the treatment of people with inherited bleeding disorders. This includes information about patient diagnosis, viral status, treatment details, hospital admissions and administrative information as well as details on ordering, supply and use of clotting factor products. Information is entered into the ABDR web enabled software by staff at HTC's. The current version of the ABDR has been in existence since December 2008 and background on the development of the system is at [Appendix D](#). In August 2012 the 4th generation ABDR was implemented.

The ABDR provides health care teams and support staff with a record enabling them to monitor and manage treatment over time to improve patients' quality of life. De-identified information from the ABDR may be used for research purposes by authorised organisations to understand and improve treatment for bleeding disorders. Considerations for the release of any information for research are made under specific governance arrangements. The ABDR also provides governments with information on total clotting factor product requirements to inform supply planning to meet the needs of all Australians with bleeding disorders.

The ABDR has evolved and improved with improvements in technology and feedback from stakeholders. In 2014 the ABDR entered a new phase with MyABDR - a secure app for smartphones (Android and iOS) and a web site for people with bleeding disorders or parents/caregivers to record home treatments and bleeds. It is an internet-based online system that gives patients a quick, easy and reliable way to:

- Record treatments and bleeds
- Manage treatment product stock
- Share the information with a Haemophilia Treatment Centre through the Australian Bleeding Disorders Registry (ABDR)
- Update contact and personal details.

ABDR MANAGEMENT AND GOVERNANCE

The ABDR is managed on a day to day basis by the National Blood Authority (NBA) in accordance with the guidance and policy oversight provided by the ABDR Steering Committee. The Committee consists of representatives of the key stakeholders involved in the clinical management, advocacy and funding of treatment for people with bleeding disorders.

Endorsement from Haemophilia Foundation Australia

Haemophilia Foundation Australia supports the ABDR. It helps doctors and other treating health professionals to understand more about the care and treatment needs of people affected by bleeding disorders. The ABDR will assist and guide planning to ensure treatment product is available when it is needed. We are confident the steps in place will mean accurate, reliable and confidential data is available and that no patient details can be identified outside haemophilia centres.

www.haemophilia.org.au

Endorsement from Australian Haemophilia Centre Directors' Organisation

The ABDR is a valuable tool that provides a summary of those affected with haemophilia and other bleeding disorders in Australia. Data from the ABDR is the best information available for clinicians to advise governments making policy decisions regarding treatment needs and product availability.

National statistics available through the ABDR will give AHCCDO an overview of practice and allow opportunities for improvement. This data can be pooled to compare Australian treatment standards with international benchmarks. The ABDR will continue to provide the ability to assess quality of life and other important clinical questions arising across Australia.

AHCCDO's partnership on this initiative with the National Blood Authority, Haemophilia Foundation Australia and other specialist health professional groups is vital to the pursuit of excellence in clinical treatment practices.

www.ahcco.org.au

In 2018-19 the Steering Committee representatives were:

- Dr Simon McRae (Chair) – Australian Haemophilia Centre Directors’ Organisation
- Dr Huyen Tran – Chair of Australian Haemophilia Centre Directors’ Organisation
- Ms Sharon Caris – Executive Director, The Haemophilia Foundation Australia
- Mr Michael Furey, VIC Health – Jurisdictional Blood Committee nominee
- Mr Ian Kemp – National Blood Authority

PATIENT PRIVACY IN ABDR AND MYABDR

The ABDR and MyABDR are provided by the National Blood Authority (NBA). The NBA is required to ensure that patient information in ABDR and MyABDR is collected and managed in a way which complies with the Commonwealth *Privacy Act 1988*. There are also parallel requirements which may apply under state and territory laws. Privacy requirements under the *Privacy Act* were tightened in 2014, and a new Privacy Policy for these systems was implemented from 26 January 2015.

More information about the management of patient privacy in ABDR and MyABDR can be found at <http://www.blood.gov.au/privacy-info-abdr-myabdr>, including a copy of the ABDR/MyABDR Privacy Policy together with further information, forms and other implementation resources.

In order to maintain the anonymity of individual patients and health providers, small cell data published or released, showing less than five (5) may be suppressed or aggregated if there is a potential to re-identify or exceptions are agreed between national and state/territory data custodians.

DATA GOVERNANCE

There is an extremely robust governance framework that oversees the management and operation of the ABDR. An AHCDO member chairs the Steering Committee tasked with these responsibilities. The Steering Committee also includes the Executive Director of Haemophilia Foundation Australia to ensure patient needs are met. Patient privacy and confidentiality are paramount to these arrangements.

In addition, there are stringent security protocols embedded into the technical architecture of the ABDR. These effectively control access to personal data ensuring this information is only accessible to treating health professionals and authorised support staff.

The database provides a capability for all HTC staff to enter data on the interactions with patients to provide treating clinicians with a comprehensive picture of the health and wellbeing of patients. The database provides for both real time ordering of product and retrospective collection of data to provide national clotting factor usage data to inform and assist planning and funding. The system also provides for inclusion of information on physiotherapy and social work interactions with patients.

To ensure appropriate management of the information, the NBA has instigated a detailed governance framework for data use and release.

DATA QUALITY ISSUES

There are a number of data quality issues in the ABDR. These include incomplete records with empty fields or entries. The data entered into some fields has also been characterised by a lack of consistency. This issue in the interpretation of specific fields has been addressed with the development of data standards for users. Application of the data standards will improve data quality. The ABDR Steering Committee has initiated strategies to improve the data quality and over time the reporting from the ABDR has become more robust. However, there are still some data quality issues that impact the data presented in this report and review of these issues continues to be addressed.

ABDR SYSTEM

The 4th Generation ABDR was successfully implemented on 13 August 2012. System enhancements are ongoing and approved by the ABDR Steering Group to enhance performance and ease of use.

COMPARING DATA FROM PREVIOUS ABDR ANNUAL REPORTS

Comprehensive automated and manual data cleansing and validation processes (that occurred as part of the implementation of the new system) enhanced the ABDR data accuracy and consistency presented in this report. This will make it difficult to undertake comparisons with data published in previous reports particularly in regards to multiple diagnoses, treatment plans, ages and dates of death. In 2014-15 historical data was refreshed for the four previous years. Continued work on the data integrity of the registry has been undertaken in 2018-19.

CONSISTENT APPLICATION OF DIAGNOSES AND DEFINITIONS

The application of definitions for bleeding disorders (e.g. VWD subtypes) varies between HTC's, and work will continue to ensure consistent approaches are used, including alignment of the severity ratings and treatment regimens for some patient records.

Commencing 2014-15 the data has been categorised by hereditary and acquired.

VON WILLEBRAND DISEASE

Not all patients with VWD are treated through HTC's and the figures in this report do not represent the total number of VWD patients in Australia.

The diagnosis of VWD subtypes and the assignment of a severity rating to the disorder can vary between HTC's. Often the treatments for VWD involve providing replacement for the missing or defective clotting factors, and use of these products is included in this report.

TREATMENTS NOT INCLUDED IN THE ABDR

The treatments for bleeding disorders often involve providing replacement for the missing or defective clotting factors. The use of commercially produced clotting factors is the subject of this report.

However, there are other clinically appropriate treatments for bleeding disorders which are not counted in this report. Other products used include cryoprecipitate (a fresh blood product), platelets (a fresh blood product) and Desmopressin (1-desamino-8-D-arginine vasopressin, abbreviated as DDAVP).

Mild cases of HMA, HMB and VWD are often treated with DDAVP. Platelet disorders may be treated with DDAVP, platelet infusion or FVIIa.

CONSENT

Patient information in the Australian Bleeding Disorders Registry (ABDR) and MyABDR is collected and managed in a way which complies with the Commonwealth *Privacy Act 1988* and parallel requirements under state and territory laws. Privacy requirements under the *Privacy Act* were tightened in 2014, and a new ABDR/MyABDR Privacy Policy applied from 26 January 2015.

A patient's personal information may be entered into the ABDR, either at a Haemophilia Treatment Centre (HTC) or when a patient enters data directly via MyABDR, and becomes part of an electronic record about the patient's bleeding disorder condition.

In accordance with the ABDR/[MyABDR Privacy Policy](#), a patient's consent is required for the recording of their data in ABDR (consent may be given by a parent, guardian or authorised representative where relevant). Where a patient does not consent then details will not be aggregated in this report, and therefore patient numbers and product use may be understated.

Supply of products for treatment

A key element in ensuring security of supply of products for the treatment of bleeding disorders is the NBA's role in developing, coordinating and monitoring the annual national supply plan and budget, including obtaining annual approval from health ministers. Further details on national supply and demand trends can be found in [Appendix C](#).

The range of products available to clinicians has changed over the years. Figure 2 shows the total issues and market shares for recombinant products from 2014-15 to 2018-19.




FIGURE 2 - MARKET SHARE OF RECOMBINANT FVIII ISSUES 2014-15 TO 2018-19

Figure 2 illustrates the changes that occurred during 2014 to 2019, brought about by new national supply arrangements, with a transition away from Kogenate and Recombinate, an increase in the issue of Xyntha and the introduction of Advate. In 2014-15 the NBA implemented new contracts for the supply of Recombinant Factor VIII and IX. The new supply arrangements have provided high level national efficiencies without detriment to the patient population. Advate accounted for approximately 52 per cent and Xyntha for 48 per cent of the market share of Recombinant FVIII issues during 2018-19 excluding EHL products or 42 per cent and 39 per cent of the market including EHL products.

The most challenging aspect of HMA management is the development of FVIII inhibitors; previously untreated patients are at the highest risk for inhibitor formation.

EXTENDED HALF LIFE (EHL) PRODUCTS

In 2018 the NBA governments endorsed limited interim supply arrangements for extended half life products to enable a limited number of patients to access EHL products under nationally funded arrangements. The agreed arrangements were to enable:

- around 140 haemophilia A patients to have access to the Shire product Adynovate (around 100 patients) or the Sanofi-aventis product Eloctate (around 40 patients), and
- around 60 haemophilia B patients to have access to the Sanofi-aventis product Alprolix.

PATIENT SUITABILITY AND PRIORITISATION

Prioritisation criteria and other considerations were agreed by a Reference Group to ensure that EHL products were directed to those patients where the greatest benefit would be obtained. Patient suitability criteria and other considerations are set out in [Appendix C](#).

During 2018-19, in total, 209 patients used EHL products (see Table 4). Some patients used product for only part of the year and other patients were able to be added to the arrangements.

Figure 3 shows the total units (IU) of EHL products supplied in 2017-18 (part year only) and 2018-19.




FIGURE 3 - TOTAL UNITS (IU) OF EXTENDED HALF LIFE PRODUCTS ISSUED 2018-19

ABDR patient demographics

This section of the report presents the key patient demographic data collected in the ABDR.

DIAGNOSES

The following tables present the numbers of patients in the ABDR and the numbers of patients who received therapeutic products during the years 2014-15 to 2018-19. As noted in the section on *Data quality issues* (page 18) comprehensive automated and manual data cleansing and validation processes that occurred as part of the 4th Generation ABDR Redevelopment project released in August 2012 and the continuation in 2018-19 enhanced the ABDR data accuracy and consistency presented in this report. This may make it difficult to undertake comparisons with data published in previous reports.

Table 3 shows the number of people in the registry and the number treated by latest broad diagnosis for the years 2014-15 to 2018-19. Table 6 expands the data in Table 3 to show the number of people in the registry and the number treated by detailed diagnosis for the years 2014-15 to 2018-19.

TABLE 3 - NUMBER OF PEOPLE IN THE REGISTRY AND TREATED BY BROAD DIAGNOSIS

Diagnosis	Number in ABDR Registry*					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
HMA	2,158	2,301	2,365	2,302	2,372	992	1,022	1,009	1,040	1104
HMB	530	548	564	541	558	218	219	218	227	247
VWD	2,012	2,092	2,141	2,146	2,221	255	287	248	239	307
Acquired										
HMA	59	74	68	74	78	23	13	11	12	15
HMB			<5	<5	<5		<5			
VWD	19	22	25	27	32	<5	8	10	5	10
Other Diagnoses										
Other‡	193	179	193	162	181	11	15	14	12	18
Other Factor Deficiency	344	391	427	449	469	36	52	50	51	58
Platelet Disorder	255	271	288	302	323	15	19	10	8	22
Vascular	7	7	9	7	7					
Fibrinogen Disorder	49	62	74	91	113	10	11	20	13	23
Total	5,626	5,947	6,155	6,102	6,355	1,563	1,647	1,590	1,607	1,804

Note: Includes asymptomatic carriers in Hereditary

* As noted in the section *Data quality issues* (p18) the data has improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

‡The ABDR allows for a diagnosis of 'Other' to be recorded for patients with rare and less prevalent disorders or difficult to classify disorders eg mild VWD

TABLE 4 - NUMBER OF HEREDITARY PATIENTS WHO RECEIVED EXTENDED HALF LIFE PRODUCTS BY BROAD DIAGNOSIS AND FINANCIAL YEAR

Diagnosis	Number of patients who received EHL products during the year	
	2017-18 (part year)	2018-19
Hereditary		
HMA	58	140
HMB	30	69
Total	88	209

PATIENTS WITH MULTIPLE BLEEDING DISORDERS

Individual patients may have more than one bleeding disorder, and will be registered with more than one diagnosis. There are patients with multiple diagnoses in the registry for 2018-19. In these cases, a patient may be counted more than once in the data in this report (e.g. if a patient has two bleeding disorders, that patient may be counted in the totals for each disorder).

In 2018-19 there were 104 patients with two diagnoses and <5 patients with three diagnoses. Of the patients with more than one diagnosis 17 patients received product.

TABLE 5 - NUMBER OF PEOPLE IN THE REGISTRY WITH MULTIPLE BLEEDING DISORDERS

Diagnosis	Patient Numbers in ABDR Registry*			Number of Patients with Multiple Disorders who Received Product during the year
	Bleeding Disorder 1	Bleeding Disorder 2	Bleeding Disorder 3	
HMA	2,450	44	<5	12
HMB	559	5		<5
VWD	2,253	20	<5	<5
Other‡	181	<5		
Other Factor Deficiency	469	21	<5	
Platelet Disorder	323	12		<5
Vascular	7	<5		
Fibrinogen Disorders	113			
Total	6,355	<112	<15	<27

Note: Includes Acquired and Hereditary disorders

* As noted in the section *Data quality issues* (p18) the data has improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

‡The ABDR allows for a diagnosis of 'Other' to be recorded for patients with rare and less prevalent disorders or difficult to classify disorders eg mild VWD

TABLE 6 - NUMBER OF PEOPLE IN THE REGISTRY AND TREATED BY DETAILED DIAGNOSIS

	Number in ABDR Registry*					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
HMA										
Asymptomatic Carrier Factor VIII Deficiency (HmA)	190	226	235	150	146	6	<5	<5	<5	<5
Factor VIII Deficiency (HmA)	1,793	1,972	2,085	2,119	2,187	972	1,011	1,005	1,036	1,100
Symptomatic Carrier Factor VIII Deficiency (HmA)	175	103	45	33	39	14	9	<5	<5	<5
HMB										
Asymptomatic Carrier Factor IX Deficiency (HmB)	47	47	53	40	39		<5			<5
Factor IX Deficiency (HmB)	426	471	489	488	503	209	213	216	225	243
Symptomatic Carrier Factor IX Deficiency (HmB)	57	30	22	13	16	9	5	<5	<5	<5
VWD										
von Willebrand Disease – Uncharacterised	279	219	180	176	181	13	8	9	9	13
von Willebrand Disease Type 1	1,233	1,328	1,387	1,379	1,430	127	137	115	98	147
von Willebrand Disease Type 2	459	502	533	545	561	84	108	94	95	108
von Willebrand Disease Type 3	41	43	41	46	49	31	34	30	37	39
Hereditary Total	4,700	4,941	5,070	4,989	5,151	1,465	1,528	1,475	1,506	1,658
Acquired										
HMA	59	74	68	74	78	23	13	11	12	15
HMB			<5	<5	<5		<5			
VWD	19	22	25	27	32	<5	8	10	5	10
Acquired Total	78	96	<97	<105	<115	<27	<22	21	17	25
Other Factor Deficiency										
Factor V Deficiency	11	17	15	15	20	<5	<5	<5	<5	6
Factor VII Deficiency	61	67	73	83	87	8	7	9	8	12

	Number in ABDR Registry*					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Factor X Deficiency	17	20	19	19	20	5	<5	6	<5	6
Factor XI Deficiency	217	249	273	286	299	14	24	18	22	19
Factor XII Deficiency†	17	17	18	18	15	<5	<5			
Factor XIII Deficiency	19	21	24	24	28	6	11	14	15	15
Platelet Disorder										
Platelet - Bernard-Soulier	5	5	7	8	10		<5			<5
Platelet - Glanzmann's Thrombasthenia	18	21	22	25	27	<5	8	6	7	7
Platelet - Macrothrombocytopenias	12	13	13	13	13					
Platelet - May Hegglin	<5	<5	<5	<5	<5		<5			
Platelet - Primary Secretion Defect	10	10	9	7	8	<5		<5	<5	
Platelet - Storage Pool (Dense Granule) Deficiency	43	46	52	59	71		<5	<5		6
Platelet – Uncharacterised	164	173	182	186	190	10	7	<5	5	7
Vascular										
Vascular Disorders - Ehlers Danlos Syndrome	7	7	9	7	7					
Fibrinogen										
Fibrinogen – Afibrinogenemia	7	7	7	7	8	5	<5	5	<5	5
Fibrinogen – Dysfibrinogenemia	29	36	45	58	73	<5	<5	9	9	9
Fibrinogen – Hypofibrinogenemia	12	17	19	23	28	<5	<5	6	<5	8
Fibrinogen Dysfunction - Uncharacterised	<5	<5	<5	<5	<5					<5
Other (Including Unclassified)	193	179	198	166	181	11	15	14	12	18
Other Diagnoses Total	848	910	991	1,011	1,093	72	97	94	84	121
Total	5,626	5,947	6,155	6,102	6,355	1,563	1,647	1,590	1,607	1,804

* As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

†Factor XII Deficiency does not require treatment with products, but is included as a diagnostic category.

AGE, DIAGNOSIS AND SEVERITY

In the following tables patients are categorised as either Adult (aged 18 years and over) or Paediatric and Adolescent (aged under 18 years).²

Table 7 and Table 8 detail the numbers of patients in the registry who received product (therapeutic treatment) during the period 2014-15 to 2018-19; by age group, broad diagnosis and by severity.

Table 9 and Table 10 set out age group and detailed diagnosis for patients with HMA, HMB and VWD.

The majority of patients receiving treatment for bleeding disorders have HMA, specifically those patients with severe HMA.

There are some discrepancies in the data regarding the coding of severity when a patient receives treatment, and data cleansing and patient record updates are continuing. This will improve the forecasting for the national supply plan and budget for future years. It should be noted that the national forecasting and supply management process continue to perform very well.

Whilst the data discrepancies affect the analysis for this annual report, there is minimal impact on patient care as Haemophilia Treatment Centre staff have full access to their patient records for the provision of care and treatment.

In 2018-19 the results show variations. The patterns indicate that the implemented strategies are improving data quality, completeness and accuracy. This will make it difficult to undertake comparisons with data published in previous reports particularly in regards to multiple diagnoses. Continued work on the data integrity of the registry has been undertaken again in 2018-19.

² In ABR Annual Reports prior to 2011-12 the threshold age between paediatric and adult patients was 20 years of age. This threshold has been adjusted in subsequent reports to better reflect the manner in which patients are treated in HTCs.

TABLE 7 - NUMBER OF ADULT PATIENTS IN THE REGISTRY AND TREATED BY BROAD DIAGNOSIS AND SEVERITY FOR HMA, HMB

Adult (aged 18 years and over)	Number in ABDR Registry					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
HMA										
Mild	996	1,040	1,007	1,030	1,063	208	227	220	215	241
Moderate	145	159	157	156	161	94	99	88	98	101
Severe	374	385	392	394	413	327	340	355	355	385
HMB										
Mild	235	240	227	228	229	55	53	54	51	57
Moderate	97	96	98	99	100	49	54	52	51	56
Severe	56	60	63	60	68	47	50	51	54	62
Total Hereditary	1,903	1,980	1,944	1,967	2,034	780	823	820	824	902
Total Acquired HMA	44	23	22	22	18	13	<5	<5	<5	<5
Total	1,947	2,003	1,966	1,989	2,052	793	<828	<825	<829	<907

Note: As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year. Patients can have their severity categorised as 'unknown' or 'not applicable' during the initial diagnosis procedures, and these figures are not shown in this table. Excludes those severities recorded as *Unknown, Not Applicable and Blank*.

TABLE 8 - NUMBER OF PAEDIATRIC AND ADOLESCENT PATIENTS IN THE REGISTRY AND TREATED BY BROAD DIAGNOSIS AND SEVERITY FOR HMA, HMB

Paediatric and Adolescent (aged less than 18 years)	Number in ABDR Registry					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
HMA										
Mild	183	205	206	220	235	54	56	49	51	62
Moderate	68	66	70	64	66	56	52	48	54	47
Severe	266	275	274	278	275	249	247	246	263	264
HMB										
Mild	51	54	47	48	52	14	13	12	13	12
Moderate	21	21	19	19	23	16	13	15	17	19
Severe	41	42	43	43	41	37	35	34	39	38
Total Hereditary	630	663	659	672	692	426	416	404	437	442
Total Acquired HMA	<5	<5	<5			<5				

Note: As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. Patients can have their severity categorised as 'unknown' or 'not applicable' during the initial diagnosis procedures, and these figures are not shown in this table. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year. Excludes those severities recorded as *Unknown, Not Applicable and Blank*.

BY AGE GROUP AND DETAILED DIAGNOSIS

TABLE 9 - NUMBER OF PEOPLE IN THE REGISTRY DIAGNOSED WITH HMA OR HMB BY AGE GROUP AND DISEASE CLASSIFICATION

	Number in ABDR Registry*					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
HMA – Adult (aged 18 years and over)										
Asymptomatic Carrier Factor VIII Deficiency	181	216	223	142	136	6	<5		<5	<5
Factor VIII Deficiency	1,293	1,441	1,536	1,562	1,618	615	658	662	668	727
Symptomatic Carrier Factor VIII Deficiency **	154	86	38	26	31	11	7	<5	<5	<5
HMA – Paediatric (aged less than 18 years)										
Asymptomatic Carrier Factor VIII Deficiency	9	10	12	8	10			<5		<5
Factor VIII Deficiency	500	531	549	557	569	357	353	343	368	373
Symptomatic Carrier Factor VIII Deficiency	21	17	7	7	8	<5	<5			
HMB – Adult (aged 18 years and over)										
Asymptomatic Carrier Factor IX Deficiency	42	44	48	36	35		<5			<5
Factor IX Deficiency	321	360	382	380	389	142	152	155	156	174
Symptomatic Carrier Factor IX Deficiency	51	25	18	10	13	9	5	<5	<5	<5
HMB – Paediatric (aged less than 18 years)										
Asymptomatic Carrier Factor IX Deficiency	5	<5	5	<5	<5					
Factor IX Deficiency	105	111	107	108	114	67	61	61	69	69
Symptomatic Carrier Factor IX Deficiency	6	5	<5	<5	<5					
Acquired										
HMA – Adult (aged 18 years and over)	57	73	67	74	78	22	13	11	12	15
HMA – Paediatric (aged less than 18 years)	<5	<5	<5			<5				
HMB – Adult (aged 18 years and over)			<5	<5	<5		<5			

* As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

** Symptomatic carriers transitioned to asymptomatic carriers and Haemophilia Factor VIII Deficiency patients, accounts for ongoing data quality changes in patient counts in 2016-17.

TABLE 10 - NUMBER OF PEOPLE IN THE REGISTRY DIAGNOSED WITH VWD BY AGE GROUP AND DISEASE CLASSIFICATION

	Number in ABDR Registry*					Number who Received Product during the year				
	2014-15	2015-16	2016-17	2017-18	2018-19	2014-15	2015-16	2016-17	2017-18	2018-19
Hereditary										
VWD – Adult (aged 18 years and over)										
von Willebrand Disease - Uncharacterised	263	231	176	148	149	11	9	6	7	11
von Willebrand Disease Type 1	1,016	1,023	1,123	1,202	1,244	108	107	116	89	123
von Willebrand Disease Type 2	320	354	394	436	453	60	68	86	77	93
von Willebrand Disease Type 3	34	35	35	35	36	24	26	27	29	28
VWD – Paediatric (aged less than 18 years)										
von Willebrand Disease - Uncharacterised	55	48	43	28	32	<5	<5	<5	<5	<5
von Willebrand Disease Type 1	220	210	205	177	186	22	20	21	9	24
von Willebrand Disease Type 2	97	105	108	109	108	13	16	22	18	15
von Willebrand Disease Type 3	8	6	8	11	13	8	5	7	8	11
Acquired										
VWD – Adult (aged 18 years and over)	17	19	22	27	32	5	<5	8	5	10
VWD – Paediatric (aged less than 18 years)	<5									

* As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

BY LOCATION

Figure 4 depicts the geographic distribution of patients in the ABDR. Patient distribution is largely in line with the distribution of the general population. However, a more detailed analysis of geographic distribution could be expected to reveal the clustering effects often associated with the distribution of genetic disorder. Excluded from Figure 4 are 11 patients that have unknown locations (10 in 2017-18).




FIGURE 4 - NUMBERS OF ACTIVE PATIENTS IN THE REGISTRY AS AT 30 JUNE 2019

Table 11 shows the numbers of patients with severe hereditary HMA and HMB, acquired HMA and the numbers of male patients with severe HMA and HMB by state and territory.

TABLE 11 - NUMBERS OF PATIENTS WITH SEVERE HMA AND HMB BY LOCATION

State/Territory	HMA			HMB	
	Severe Hereditary	Severe Hereditary Males	Severe Acquired	Severe Hereditary	Severe Hereditary Males
ACT	14	14		<5	<5
NSW	204	202	6	38	38
NT	<5	<5			
QLD	152	151	<5	20	20
SA	54	54	<5	5	5
TAS	18	18		<5	<5
VIC	169	169		35	35
WA	74	74	<5	8	7
Total	688	685	<22	109	108

As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year. This table excludes patients with an unknown location.

BY GENDER AND AGE DISTRIBUTION

The figures in this section present the gender and age distribution of patients in the ABDR in 2018-19, compared to the general Australian population³. Table 12 sets out the split between male and female patients with bleeding disorders, and Table 13 sets out numbers of patients by age group compared with Australian population figures. Table 14 sets out HMA, HMB and VWD patients by age group and gender.

TABLE 12 - PATIENT NUMBERS BY BLEEDING DISORDER AND GENDER

Bleeding Disorder	Female	Male	Grand Total
Asymptomatic Carrier Factor VIII Deficiency (Haemophilia A)	146		146
Factor VIII Deficiency (Haemophilia A)	210	1,977	2,187
Symptomatic Carrier Factor VIII Deficiency (Haemophilia A)	38	<5	<43
Asymptomatic Carrier Factor IX Deficiency (Haemophilia B)	38	<5	<43
Factor IX Deficiency (Haemophilia B)	60	443	503
Symptomatic Carrier Factor IX Deficiency (Haemophilia B)	15	<5	<20
Von Willebrand Disease - Uncharacterised	107	74	181
Von Willebrand Disease Type 1	952	478	1,430
Von Willebrand Disease Type 2 - Uncharacterised	69	48	117
Von Willebrand Disease Type 2A	74	60	134
Von Willebrand Disease Type 2B	33	35	68
Von Willebrand Disease Type 2M	114	82	196
Von Willebrand Disease Type 2N	34	12	46
Von Willebrand Disease Type 3	25	24	49
Factor V Deficiency	9	9	18
Factor VII Deficiency	44	43	87
Factor X Deficiency	9	9	18
Factor XI Deficiency	199	99	298
Factor XII Deficiency	9	6	15
Factor XIII Deficiency	11	17	28
Fibrinogen - Afibrinogenemia	<5	6	<11
Fibrinogen - Dysfibrinogenemia	49	24	73
Fibrinogen - Hypofibrinogenemia	15	13	28
Fibrinogen Dysfunction - Uncharacterised	<5		<5
Platelet Dysfunction - Bernard-Soulier	5	5	10
Platelet Dysfunction - Glanzmann's Thrombasthenia	16	11	27

³ Australian Bureau of Statistics, Australian Demographic Statistics, Cat. No. 3101.0, Population by Age and Sex, released June 2019 (Table 6) – data as at December 2018

TABLE 12 CONTINUED - PATIENT NUMBERS BY BLEEDING DISORDER AND GENDER

Bleeding Disorder	Female	Male	Grand Total
Platelet Dysfunction - Macrothrombocytopenias	7	6	13
Platelet Dysfunction - May Hegglin	<5	<5	<10
Platelet Dysfunction - Primary Secretion Defect	7	<5	<12
Platelet Dysfunction - Storage Pool (Dense Granule) Deficiency	43	28	71
Platelet Dysfunction - Uncharacterised	119	71	190
Vascular Disorders - Ehlers Danlos Syndrome	<5	5	<10
(Acquired) Factor VIII Deficiency (Haemophilia A)	32	46	78
(Acquired) Factor IX Deficiency (Haemophilia B)	<5		<5
(Acquired) Von Willebrand Disease - Uncharacterised	11	12	23
(Acquired) Von Willebrand Disease Type 1	<5	<5	<10
(Acquired) Von Willebrand Disease Type 2 - Uncharacterised		<5	<5
(Acquired) Von Willebrand Disease Type 2A	<5		<5
(Acquired) Von Willebrand Disease Type 3		<5	<5
(Acquired) Factor V Deficiency	<5	<5	<10
(Acquired) Factor X Deficiency		<5	<5
(Acquired) Factor XI Deficiency	<5		<5
(Acquired) Other	<5	<5	<10
Other	111	52	163
No Bleeding Disorder recorded	14	<5	<19
Total	2,643	3,712	6,355

TABLE 13 - PATIENT NUMBERS BY BLEEDING DISORDER AND AGE GROUP

Age Range Jun30 2019	Haemophilia A	Haemophilia B	Von Willebrand Disease	Factor V Deficiency	Factor VII Deficiency	Factor X Deficiency	Factor XI Deficiency	Factor XII Deficiency	Factor XIII Deficiency	Other Factor Deficiency	Fibrinogen	Platelet Disorder	Vascular	Other	Null	Total with bleeding disorders	% for bleeding disorder patients	Total for population	% for population
0 - 4	119	22	33	<5	<5		5		<5		5	<5				189	3.0%	1,582,296	6.3%
5 - 9	175	27	87	<5	<5	<5	12	<5	<5		7	12	<5	6		334	5.3%	1,604,580	6.4%
10 - 14	201	48	135	<5	<5	<5	9		<5		6	22		<5		433	6.8%	1,515,838	6.1%
15 - 19	153	38	132	<5	<5	<5	17	<5	5		11	27	<5	8	<5	400	6.3%	1,490,664	6.0%
20 - 24	176	35	209	<5	9	<5	23	<5	<5		12	40		19	<5	535	8.4%	1,740,205	7.0%
25 - 29	192	44	193		13	<5	21	<5	5		5	23	<5	16		518	8.2%	1,876,948	7.5%
30 - 34	190	41	202	<5	12	<5	25	<5	<5		14	25	<5	11	<5	531	8.4%	1,862,445	7.5%
35 - 39	196	45	177	<5	13	<5	25				6	28		18	<5	516	8.1%	1,722,521	6.9%
40 - 44	166	45	176	<5	<5	<5	29	<5	5		<5	24		9	<5	464	7.3%	1,594,067	6.4%
45 - 49	170	49	197		<5	<5	25	<5			15	16	<5	15	<5	494	7.8%	1,671,426	6.7%
50 - 54	120	30	149	<5	<5	<5	14	<5		<5	7	19		16	<5	364	5.7%	1,528,926	6.1%
55 - 59	109	29	128		7		13				<5	18		11	<5	319	5.0%	1,529,464	6.1%
60 - 64	121	33	108	<5	<5		13				7	19		12		316	5.0%	1,359,299	5.4%
65 - 69	122	19	100	<5	<5		17	<5		<5	<5	18	<5	<5		288	4.5%	1,206,750	4.8%
70 - 74	89	28	97	<5	<5	<5	15	<5			6	12		12		267	4.2%	1,017,730	4.1%
75 - 79	60	13	59	<5	<5	<5	14			<5		9		6		170	2.7%	700,298	2.8%
80 - 84	37	<5	34		<5		11	<5			<5	<5		<5		100	1.6%	485,620	1.9%
85 - 89	34	<5	18				<5				<5	<5				63	1.0%	309,055	1.2%
90 - 94	14	<5	15		<5		<5					<5				38	0.6%	149,761	0.6%
95 and over	6	<5	<5				<5				<5			<5		16	0.2%	44,854	0.2%
Total	2,450	559	2,253	18	87	18	298	15	28	5	113	323	7	165	16	6,355		24,992,747	

TABLE 14 - HEREDITARY HMA, HMB AND VWD PATIENTS BY GENDER AND AGE RANGE AT 30 JUNE 2019

Age Range Jun30 2019	Haemophilia A		Haemophilia B		Von Willebrand Disease		Total
	Female	Male	Female	Male	Female	Male	
0 - 4	5	114	<5	19	16	17	<176
5 - 9	19	156	<5	23	32	55	<290
10 - 14	16	185	8	40	54	81	384
15 - 19	15	138	9	29	63	69	323
20 - 24	20	156	8	27	136	73	420
25 - 29	26	166	8	36	121	72	429
30 - 34	49	138	7	34	143	59	430
35 - 39	52	139	8	37	121	52	409
40 - 44	44	118	9	36	133	43	383
45 - 49	41	128	13	36	138	54	410
50 - 54	27	91	<5	26	98	49	<299
55 - 59	21	87	6	23	85	41	263
60 - 64	19	99	6	27	64	41	256
65 - 69	21	94	<5	15	67	31	<236
70 - 74	9	70	7	21	63	30	200
75 - 79	6	39	6	6	35	20	112
80 - 84	<5	27	<5	<5	17	14	<73
85 - 89	<5	22	<5	<5	10	7	<54
90 - 94		6		<5	9	<5	<25
95 - 104		5		<5	<5	<5	<15
Grand Total	394	1,978	113	445	1,408	813	5,151

Figure 5 and Figure 6 chart the distribution of all female hereditary HMA and HMB patients against the female population. The tables next to each figure show the numbers and percentages used in the charts.




FIGURE 5 - DISTRIBUTION OF HEREDITARY FEMALE HMA PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 5 – DISTRIBUTION OF HEREDITARY FEMALE HMA PATIENTS BY AGE 2018-19

Age group	2018 Australian Female Population	% 2018 Australian Female Population	HMA female patients	% HMA female patients	Patient average weight 2018-19
0-4	769,381	6.0%	5	1.3%	13
5-9	781,148	6.2%	19	4.8%	23
10-14	736,599	5.9%	16	4.1%	39
15-19	725,727	5.7%	15	3.8%	54
20-24	849,641	6.7%	20	5.1%	67
25-29	936,647	7.4%	26	6.6%	70
30-34	941,358	7.5%	49	12.4%	68
35-39	864,948	7.0%	52	13.2%	76
40-44	800,767	6.3%	44	11.2%	67
45-49	852,103	6.7%	41	10.4%	79
50-54	779,358	6.1%	27	6.9%	76
55-59	779,427	6.2%	21	5.3%	71
60-64	697,814	5.6%	19	4.8%	70
65-69	616,925	4.9%	21	5.3%	70
70+	1,464,084	11.9%	19	4.8%	80
All ages	12,595,927		394		65

Notes:

- Patient weight values are averaged across the year.




FIGURE 6 - DISTRIBUTION OF HEREDITARY FEMALE HMB PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 6 – DISTRIBUTION OF HEREDITARY FEMALE HMB PATIENTS BY AGE 2018-19

Age group	2018 Australian Female Population	% 2018 Australian Female Population	HMB female patients	% HMB female patients	Patient average weight 2018-19
0–4	769,381	6.0%	<5	<4.4%	12
5–9	781,148	6.2%	<5	<4.4%	20
10–14	736,599	5.9%	8	7.1%	41
15–19	725,727	5.7%	9	8.0%	53
20–24	849,641	6.7%	8	7.1%	70
25–29	936,647	7.4%	8	7.1%	61
30–34	941,358	7.5%	7	6.2%	75
35–39	864,948	7.0%	8	7.1%	61
40–44	800,767	6.3%	9	8.0%	64
45–49	852,103	6.7%	13	11.5%	76
50–54	779,358	6.1%	<5	<4.4%	78
55–59	779,427	6.2%	6	5.3%	75
60–64	697,814	5.6%	6	5.3%	75
65–69	616,925	4.9%	<5	<4.4%	76
70+	1,464,084	11.9%	16	14.2%	66
All ages	12,595,927		113		61

Notes:

- Patient weight values are averaged across the year.

Figure 7 and Figure 8 chart the distribution of all male hereditary HMA patients and all male severe hereditary HMA patients against the male population. The disorder is genetically linked to a patient's gender, and usually affects males. There are a relatively lower number of older patients (from the age grouping of 75-79 years onwards). The life expectancy of HMA patients has improved dramatically⁴ in recent decades. The younger cohorts can expect to survive longer, which will increase the overall patient population and the demand for product in the future.

The number of acquired HMA severe male patients totalled 8.




FIGURE 7 - DISTRIBUTION OF HEREDITARY MALE HMA PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 7 – DISTRIBUTION OF HEREDITARY MALE HMA PATIENTS BY AGE 2018-19

Age group	2018 Australian Male Population	% 2018 Australian Male Population	HMA male patients	% HMA male patients	Patient average weight 2018-19
0-4	812,915	6.4%	114	5.8%	13
5-9	823,432	6.6%	156	7.9%	25
10-14	779,239	6.4%	185	9.4%	42
15-19	764,937	6.1%	138	7.0%	66
20-24	890,564	7.2%	156	7.9%	76
25-29	940,301	7.6%	166	8.4%	80
30-34	921,087	7.4%	138	7.0%	86
35-39	857,573	7.0%	139	7.0%	89
40-44	793,300	6.3%	118	6.0%	91
45-49	819,323	6.6%	128	6.5%	90
50-54	749,568	6.0%	91	4.6%	89
55-59	750,037	6.0%	87	4.4%	87
60-64	661,485	5.4%	99	5.0%	89
65-69	589,825	4.7%	94	4.8%	87
70+	1,243,234	10.3%	169	8.5%	82
All ages	12,396,820		1,978		71

Notes:

- Patient weight values are averaged across the year.

⁴ Oldenburg J, Dolan G, Lemm G (2009). Haemophilia care then, now and in the future. Haemophilia 15, S1: 2-7.




FIGURE 8 - DISTRIBUTION OF HEREDITARY MALE HMA SEVERE PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 8 – DISTRIBUTION OF HEREDITARY MALE HMA SEVERE PATIENTS BY AGE 2018-19

Age group	2018 Australian Male Population	% 2018 Australian Male Population	HMA severe male patients	% HMA severe male patients	Patient average weight 2018-19
0-4	812,915	6.4%	61	8.9%	14
5-9	823,432	6.6%	78	11.4%	25
10-14	779,239	6.4%	94	13.7%	42
15-19	764,937	6.1%	61	8.9%	67
20-24	890,564	7.2%	71	10.4%	76
25-29	940,301	7.6%	57	8.3%	82
30-34	921,087	7.4%	64	9.3%	86
35-39	857,573	7.0%	46	6.7%	90
40-44	793,300	6.3%	29	4.2%	82
45-49	819,323	6.6%	42	6.1%	88
50-54	749,568	6.0%	22	3.2%	85
55-59	750,037	6.0%	15	2.2%	84
60-64	661,485	5.4%	15	2.2%	81
65-69	589,825	4.7%	16	2.3%	76
70+	1,243,234	10.3%	14	2.0%	82
All ages	12,396,820		685		63

Notes:

- Patient weight values are averaged across the year.

Figure 9 and Figure 10 chart the distribution of all male hereditary HMB patients and all male severe hereditary HMB patients against the male population. As with HMA, HMB is also genetically linked to a patient’s gender, and usually affects males. The observed male severe HMB population does not conform to the same pattern as the general male population, however there are low patient numbers (n=108) in this group and no conclusions should be drawn.

There were no acquired HMB severe male patients.




FIGURE 9 - DISTRIBUTION OF HEREDITARY MALE HMB PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 9 – DISTRIBUTION OF HEREDITARY MALE HMB PATIENTS BY AGE 2018-19

Age group	2018 Australian Male Population	% 2018 Australian Male Population	HMB male patients	% HMB male patients	Patient average weight 2018-19
0–4	812,915	6.4%	19	4.3%	13
5–9	823,432	6.6%	23	5.2%	23
10–14	779,239	6.4%	40	9.0%	42
15–19	764,937	6.1%	29	6.5%	63
20–24	890,564	7.2%	27	6.1%	78
25–29	940,301	7.6%	36	8.1%	81
30–34	921,087	7.4%	34	7.6%	90
35–39	857,573	7.0%	37	8.3%	90
40–44	793,300	6.3%	36	8.1%	91
45–49	819,323	6.6%	36	8.1%	97
50–54	749,568	6.0%	26	5.8%	88
55–59	750,037	6.0%	23	5.2%	88
60–64	661,485	5.4%	27	6.1%	82
65–69	589,825	4.7%	15	3.4%	85
70+	1,243,234	10.3%	37	8.3%	93
All ages	12,396,820		445		75

Notes:

- Patient weight values are averaged across the year.




FIGURE 10 - DISTRIBUTION OF HEREDITARY MALE HMB SEVERE PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 10 – DISTRIBUTION OF HEREDITARY MALE HMB SEVERE PATIENTS BY AGE 2018-19

Age group	2018 Australian Male Population	% 2018 Australian Male Population	HMB severe male patients	% HMB severe male patients	Patient average weight 2018-19
0-4	812,915	6.4%	11	10.2%	13
5-9	823,432	6.6%	9	8.3%	24
10-14	779,239	6.4%	13	12.0%	50
15-19	764,937	6.1%	11	10.2%	67
20-24	890,564	7.2%	13	12.0%	80
25-29	940,301	7.6%	6	5.6%	75
30-34	921,087	7.4%	10	9.3%	87
35-39	857,573	7.0%	10	9.3%	89
40-44	793,300	6.3%	8	7.4%	95
45-49	819,323	6.6%	6	5.6%	91
50-54	749,568	6.0%	5	4.6%	86
55-59	750,037	6.0%	<5	<4.6%	67
60-64	661,485	5.4%	<5	<4.6%	76
65-69	589,825	4.7%			
70+	1,243,234	10.3%	<5	<4.6%	100
All ages	12,396,820		108		67

Notes:

- Patient weight values are averaged across the year.

Figure 11 and Figure 12 chart the distribution of female and male VWD patients against the female and male populations.




FIGURE 11 - DISTRIBUTION OF HEREDITARY FEMALE VWD PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 11 – DISTRIBUTION OF HEREDITARY FEMALE VWD PATIENTS BY AGE 2018-19

Age group	2018 Australian Female Population	% 2018 Australian Female Population	VWD female patients	% VWD female patients	Patient average weight 2018-19
0-4	769,381	6.0%	16	1.1%	14
5-9	781,148	6.2%	32	2.3%	22
10-14	736,599	5.9%	54	3.8%	37
15-19	725,727	5.7%	63	4.5%	54
20-24	849,641	6.7%	136	9.7%	70
25-29	936,647	7.4%	121	8.6%	68
30-34	941,358	7.5%	143	10.2%	71
35-39	864,948	7.0%	121	8.6%	70
40-44	800,767	6.3%	133	9.4%	74
45-49	852,103	6.7%	138	9.8%	74
50-54	779,358	6.1%	98	7.0%	77
55-59	779,427	6.2%	85	6.0%	73
60-64	697,814	5.6%	64	4.5%	74
65-69	616,925	4.9%	67	4.8%	76
70+	1,464,084	11.9%	137	9.7%	72
All ages	12,595,927		1,408		68

Notes:

- Patient weight values are averaged across the year.




FIGURE 12 - DISTRIBUTION OF HEREDITARY MALE VWD PATIENTS BY AGE IN 2018-19

DATA TABLE - FIG 12 – DISTRIBUTION OF HEREDITARY MALE VWD PATIENTS BY AGE 2018-19

Age group	2018 Australian Male Population	% 2018 Australian Male Population	VWD male patients	% VWD male patients	Patient average weight 2018-19
0-4	812,915	6.4%	17	2.1%	13
5-9	823,432	6.6%	55	6.8%	22
10-14	779,239	6.4%	81	10.0%	38
15-19	764,937	6.1%	69	8.5%	57
20-24	890,564	7.2%	73	9.0%	70
25-29	940,301	7.6%	72	8.9%	78
30-34	921,087	7.4%	59	7.3%	78
35-39	857,573	7.0%	52	6.4%	87
40-44	793,300	6.3%	43	5.3%	92
45-49	819,323	6.6%	54	6.6%	86
50-54	749,568	6.0%	49	6.0%	90
55-59	750,037	6.0%	41	5.0%	84
60-64	661,485	5.4%	41	5.0%	86
65-69	589,825	4.7%	31	3.8%	96
70+	1,243,234	10.3%	76	9.3%	85
All ages	12,396,820		813		69

Notes:

- Patient weight values are averaged across the year.

INHIBITOR STATUS

Table 15 provides a description of the inhibitor status used in the ABDR. Table 16 shows the status of inhibitors for patients as at 30 June 2019. Inhibitors are immunoglobulins made by the body's immune system to react against replacement clotting factor proteins. This occurs when the immune system perceives the proteins as foreign or harmful to the body. When this happens, the inhibitors prevent the usual replacement factors (Factor VIII or IX) from working properly to stop bleeding.

Inhibitor detection is conducted using the Bethesda assay, with or without the Nijmegen modification (Verbruggen et al. 1995), and results are expressed in Bethesda units (BU)⁵. If the inhibitor titre is high (>5 BU/ml), factor replacement therapy is ineffective and bleeding persists. With low titre inhibitor (<5 BU/ml), haemostasis may be achieved with higher doses. Patients with severe Haemophilia A with high-titre inhibitors are most at risk for recurrent bleeds and chronic haemarthroses.

FEIBA and Recombinant Factor VIIa (brand name NovoSeven) are both used to treat patients that have developed inhibitors. In the setting of managing inhibitors for haemophilia, the drivers for clinical demand for FEIBA are similar to those for NovoSeven. Predicting or interpreting changing demand trends is not possible with any accuracy, as the product is only used in a small number of patients each year. Use patterns will vary from year to year and will not only depend on the number of patients treated, but their severity of disease, the potency of inhibitors, whether secondary prophylaxis is practiced, the number and severity of spontaneous bleeds, and the amount of elective surgery undertaken in this patient group.

⁵ Bethesda units (BU) = a measure of inhibitor activity – the amount of inhibitor that inactivates 50% or 0.5 units of a coagulation factor during the incubation period

TABLE 15 - DESCRIPTION OF INHIBITOR STATUS USED IN ABDR

Inhibitor Event Type	Screening or Inhibitor Status
Initial Inhibitor Status	<ul style="list-style-type: none"> • <i>Inhibitor Testing Not Performed</i> - No inhibitor test has ever been performed for this patient • <i>Unknown</i> – Used if a patient has been tested but the results are unknown (i.e. transferred from overseas)
Screening Test Result	<ul style="list-style-type: none"> • <i>Negative</i> - Patient has a negative screening test result (then enter Inhibitor Status) • <i>Equivocal</i> - Not determined • <i>Present</i> - Patient has a positive screening test result
Screening Test (Result is Negative) or Inhibitor Test	<ul style="list-style-type: none"> • <i>Currently present – not on ITI</i> - Patient has an inhibitor but is not currently on ITI therapy • <i>Never Present</i> – No inhibitor detected for this test or previous tests performed • <i>Previously present – high responder (>5 BU/mL)</i> – Patient is negative this occasion but previously had a high inhibitor level to FVIII / FIX where the titre level is greater than 5 BU/mL • <i>Previously present – low responder (<5 BU/mL)</i> – Patient is negative this occasion but previously had a low inhibitor level to FVIII / FIX where the titre level less than 5 BU/mL • <i>On ITI</i> – Patient is on Immune Tolerance Induction (ITI) therapy or Tolerisation • <i>Unknown</i> – recorded as blank • <i>Present</i> – Patient has a positive inhibitor test result (Migrated data from previous version of ABDR and can no longer be used) • <i>Historic</i> - Patient does not currently have an inhibitor but has previously had one (Migrated data from previous version of ABDR and can no longer be used) • <i>Tolerised</i> - Patient has previously had an inhibitor in the past and been successfully tolerised (Migrated data from previous version of ABDR and can no longer be used) previous titre eg high or low responder – not known

TABLE 16 - PATIENT INHIBITOR STATUS NUMBERS

	30-Jun-18	30-Jun-19
HMA	2,376	2,450
Currently Present - Not on ITI	51	68
Equivocal	6	6
Historic	<5	<5
Inhibitor Testing Not Performed	788	804
Negative	10	11
Never Present	1,273	1,321
On ITI	25	23
Present	11	7
Previously Present - High Responder (≥ 5 BU/mL)	105	103
Previously Present - Low Responder (< 5 BU/mL)	104	105
Tolerised	<5	<5
HMB	542	559
Currently Present - Not on ITI	<5	<5
Equivocal	<5	
Inhibitor Testing Not Performed	249	255
Negative	5	5
Never Present	279	288
On ITI	<5	<5
Previously Present - High Responder (≥ 5 BU/mL)	<5	5
Previously Present - Low Responder (< 5 BU/mL)	<5	<5
VWD	2,173	2,253
Currently Present - Not on ITI	<5	<5
Inhibitor Testing Not Performed	2,099	2,177
Never Present	65	67
On ITI	<5	<5
Present	<5	<5
Previously Present - High Responder (≥ 5 BU/mL)	<5	<5
Previously Present - Low Responder (< 5 BU/mL)	<5	<5

* As noted in the section *Data quality issues* (p18) the data has been improved since previous ABDR Annual Reports. The figures presented here represent the most accurate data currently available. The census date for number of people in the registry is 30 June, the last day of the financial year.

INCIDENCE OF MAJOR DISORDERS

When we consider the incidence of bleeding disorders in global terms we see great variety in data and the reported prevalence. Table 17 details the incidence statistics from the World Federation of Hemophilia (WFH) global survey 2018 released in 2019.

TABLE 17 - INCIDENCE STATISTICS FROM WORLD FEDERATION OF HEMOPHILIA GLOBAL SURVEY 2018

Country	Population	HMA/HMB	VWD	OBD	HMA/HMB per100,000	VWD per 100,000	OBD per 100,000	Factor VIII per capita	Factor IX per capita
Australia	24,992,369	2,653	2,146	840	10.62	8.59	3.36	6.78	1.15
New Zealand	4,885,500	556	405	101	11.38	8.29	2.07		
UK	66,488,991	8,348	10,969	9,504	12.56	16.50	14.29	8.83	1.19
USA	327,167,434	17,757	11,805	6,700	5.43	3.61	2.05	9.96	1.65
Canada	37,058,856	3,687	4,321	2,057	9.95	11.66	5.55	8.15	1.58
France	66,987,244	7,944	2,479	1,033	11.86	3.70	1.54		
Sweden	10,183,175	936	203	-	9.19	1.99	-	10.13	2.23
Germany	82,927,922	4,139	3,777	-	4.99	4.55	-	7.60	0.70
South Africa	57,779,622	2,332	647	220	4.04	1.12	0.38	1.22	0.17
Japan	126,529,100	6,457	1,325	387	5.10	1.05	0.31	6.11	0.90

Abbreviations; OBD - other bleeding disorders (i.e. not HMA, HMB, VWD)

In 2010, Stonebraker *et al*⁶ reported on HMA prevalence data for 106 countries from the WFH annual global surveys and the literature. They found that the reported HMA prevalence varied considerably among countries, even among the wealthiest of countries. Prevalence data reported from the WFH compared well with prevalence data from the literature, but patient registries (such as the ABDR) generally provided the highest quality prevalence data.

In 2011, the same group reported on the prevalence of Haemophilia B⁷. Data was reported for 105 countries from the WFH annual global surveys. They reported that the prevalence varied considerably among countries, even among the wealthiest of countries.

Prevalence data is extremely valuable information for the planning efforts of national healthcare agencies in setting priorities and allocating resources for the treatment of bleeding disorders.

⁶ Stonebraker JS, Bolton-Maggs PHB, Soucie JM, Walker I, Brooker M. (2010). A study of variations in the reported hemophilia A prevalence around the world. *Haemophilia* 16(1): 20–32.

⁷ Stonebraker JS, Bolton-Maggs PHB, Soucie JM, Walker I, Brooker M. (2011). A study of variations in the reported hemophilia B prevalence around the world. *Haemophilia* 18(3): 1-4.

Patient Treatment in 2018-19

The data in this section relates to patients who received treatment (products) during the 2018-19 financial year. Figure 13 and Figure 14 show data for the period 2014-15 to 2018-19, and chart the relative volume of therapeutic products issued according to patient severity. Patients with greater severity of bleeding disorders received more products.

PRODUCTS ISSUED TO PATIENTS

Figure 13 shows the proportion of hereditary HMA patients receiving treatment (1,104 patients in 2018-19) by severity. For the five financial years, around 60% (by volume) of all FVIII products issued were for patients with severe HMA.

Figure 14 shows the proportion of hereditary HMB patients receiving treatment (247 patients in 2018-19) by severity. For the five financial years, around 40% (by volume) of all FIX products issued were for patients with severe HMB. There are far fewer HMB patients in the registry than there are HMA patients.

Around 38% of the patients in the ABDR are diagnosed with HMA (see Table 3). In relative terms, HMA is the most important consideration for national supply planning, and the key factor is the issue of product to severe HMA patients.




FIGURE 13 - PERCENTAGE OF PATIENTS RECEIVING PRODUCT BY SEVERITY FOR HMA - HEREDITARY BLEEDING DISORDERS
Note: A very small number of patients have a severity recorded as Not Applicable or Unknown. These are not shown in the above chart.




FIGURE 14 - PERCENTAGE OF PATIENTS RECEIVING PRODUCT BY SEVERITY FOR HMB - HEREDITARY BLEEDING DISORDERS

Note: Proportion of patients receiving product by severity for HmB – Acquired bleeding disorders are too small to present in graphical format. A very small number of patients have a severity recorded as Not Applicable or Unknown. These are not shown in the above chart.

Tables 18-21 detail the volume (IU) of product issued for HMA, HMB, VWD and other diagnosis patients in 2018-19 and across 5 years. The volumes are subdivided by severity and treatment regimen as stated in the ABRD record. The largest and most important sectors are products for severe HMA patients for *on demand* and *prophylactic* treatment regimens. The volume issued for prophylactic treatment of severe HMA is the single greatest determining factor for supply planning.

TABLE 18 - VOLUME (IU) OF PRODUCT ISSUED FOR HMA, HMB AND VWD PATIENTS, BY SEVERITY AND TREATMENT REGIMEN IN 2018-19 - HEREDITARY BLEEDING DISORDERS

	Mild	Moderate	Severe	Unknown**	Total**
HMA (IU FVIII Products)†	6,722,000	13,234,250	132,079,550	4,000	152,039,800
On Demand	4,556,750	3,748,750	11,418,050		19,723,550
Prophylaxis	1,505,250	9,438,000	112,764,500		123,707,750
Tolerisation	494,000		7,817,750		8,311,750
Unknown*	166,000	47,500	79,250	4,000	296,750
HMB (IU FIX Products)‡	2,198,750	6,549,500	8,092,250	5,000	16,845,500
On Demand	2,009,750	3,846,500	1,530,750	3,500	7,390,500
Prophylaxis	132,000	2,633,500	6,047,500		8,813,000
Tolerisation			488,000		488,000
Unknown*	57,000	69,500	26,000	1,500	154,000
VWD (IU FVIII Product) ++	477,750	699,000	3,990,750	2,185,250	7,352,750
On Demand	246,250	612,750	1,260,750	784,250	2,904,000
Prophylaxis	196,000	64,750	2,265,000	1,373,500	3,899,250
Tolerisation			465,000		465,000
Unknown*	35,500	21,500		27,500	84,500

† FVIII Products included are Advate, Biostate and Xyntha

‡ FIX Products included are BeneFIX, MonoFIX and Rixubis

++ FVIII Products include Advate and Biostate

* This represents a blank/not completed/empty field for the treatment regimen in the ABRD

** The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown severities.

Table 19 shows the volume of product in IUs issued to hereditary HMA, HMB and VWD patients across the five years 2014-15 to 2018-19. Both patient numbers and IUs issued have increased over time. The introduction of EHL products in 2018 has seen some HMA and HMB patients move to those products. Supply and uptake of EHL products are discussed further in [Appendix C](#).

TABLE 19 - VOLUME (IU) OF PRODUCT ISSUED FOR HMA, HMB AND VWD PATIENTS OVER TIME, INCLUDING EHL PRODUCTS 2014-15 TO 2018-19 - HEREDITARY BLEEDING DISORDERS

	2014-15	2015-16	2016-17	2017-18	2018-19
HMA	147,515,250	156,355,618	156,701,760	157,756,670	152,039,800
HMB	26,442,100	26,292,500	26,631,900	27,193,875	16,845,500
VWD	6,088,000	5,904,750	6,734,250	7,101,002	7,352,750
HMA - EHL products				3,846,000	32,150,250
HMB - EHL products				1,484,250	10,869,250
Total	180,045,350	188,552,868	190,067,910	197,381,797	219,257,550

TABLE 20 - VOLUME (IU) OF PRODUCT ISSUED FOR HMA, HMB AND VWD PATIENTS, BY SEVERITY AND TREATMENT REGIMEN IN 2018-19 - ACQUIRED BLEEDING DISORDERS

	Mild	Moderate	Severe	Unknown**	Total**
HMA (IU FVIII Products)†				99,000	99,000
On Demand					
Unknown*				99,000	99,000
VWD (IU FVIII Product) ++	2,000		37,000	764,750	803,750
On Demand	2,000		37,000	638,500	677,500
Unknown*				126,250	126,250

++ FVIII Products include Advate and Biostate

* This represents a blank/not completed/empty field for the treatment regimen in the ABDR

** The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown severities.

TABLE 21 - VOLUME (IU) OF PRODUCTS ISSUED FOR OTHER PATIENTS, BY SEVERITY AND TREATMENT REGIMEN IN 2018-19 - OTHER DIAGNOSES

	Mild	Moderate	Severe	Unknown**	Total**
Other Factor Deficiency	69,211	21,656	208,223	29,000	328,090
On Demand	6,711	7,406	24,014	3,500	41,631
Prophylaxis	62,500	14,250	184,208		260,958
Unknown*			1	25,500	25,501
Other				20,000	20,000
On Demand				4,000	4,000
Prophylaxis					
Unknown*				16,000	16,000

* This represents a blank/not completed/empty field for the treatment regimen in the ABDR

** The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown severities.

VOLUME (IU) OF PRODUCTS ISSUED FOR HMA AND HMB

Severe haemophilia requires lifelong treatment with expensive products. Clotting factor consumption is often expressed in IU/kg/year, and the ranges reported vary by population.^{8,9} Figure 15 shows the clotting factor consumption of FVIII during 2018-19 for severe HMA patients on prophylaxis. There is a wide range of use across these age groups, which are not normally distributed. Median values for each age bracket are listed below. Note there are significant outliers which require further investigation.

Median IU/Kg/year	0-4 years	5-9 years	10-14 years	15-17 years	Adult
2018-19	4,983	4,995	4,109	4,629	3,339




FIGURE 15 - FVIII PRODUCT USAGE (IU/KG/YEAR) IN SEVERE HMA PATIENTS ON PROPHYLAXIS

⁸ Schramm W, Royal S, Kroner B, Berntorp E, Giangrande P, Ludlam CA, et al. (2002). Clinical outcomes and resource utilization associated with haemophilia care in Europe. *Haemophilia* 8(1): 33-43.

⁹ Aledort LM, Haschmeyer RH, Pettersson H (1994) A longitudinal study of orthopaedic outcomes for severe factor-VIII-deficient haemophiliacs. The Orthopaedic Outcome Study Group. *J Intern Med.* 236(4): 391-399.

Figure 16 shows the clotting factor consumption of FVIII during 2018-19 for severe HMA patients on demand regimen. As in previous years there is a wide range of use across the paediatrics (includes adolescents) and adult age groups, which are not normally distributed.

Median IU/Kg/year	Paediatric	Adult
2018-19	419	1,202




FIGURE 16 - FVIII PRODUCT USAGE (IU/KG/YEAR) IN SEVERE HMA PATIENTS ON DEMAND

Figure 17 shows the clotting factor consumption during 2018-19 for severe HMB patients on prophylaxis regimen.

Median IU/Kg/year	0-4 years	5-9 years	10-14 years	15-17 years	Adult
2018-19	1,479	845	769	216	2,440




FIGURE 17 - FIX PRODUCT USAGE (IU/KG/YEAR) IN SEVERE HMB PATIENTS ON PROPHYLAXIS

Figure 18 shows the clotting factor consumption during 2018-19 for severe HMB patients on demand regimen.

Median IU/Kg/year	Paediatric	Adult
2018-19	403	727




FIGURE 18 - FIX PRODUCT USAGE (IU/KG/YEAR) IN SEVERE HMB PATIENTS ON DEMAND

These figures are higher than some of those reported in the literature for adult patients, but reflect the shift in treatment practice towards regular prophylactic infusions to prevent bleeds, especially in children. Recent theoretical work allowed for the comparison of different treatment strategies, ranging from long-term on demand therapy to different prophylactic strategies.¹⁰ In time the ABDR data should provide further insight into these issues.

¹⁰ Fischer K, Pouw ME, Lewandowski D, Janssen MP, van den Berg HM, van Hout BA (2011). A modelling approach to evaluate long-term outcome of prophylactic and on demand treatment strategies for severe hemophilia A. *Haematologica* 96(5): 738-743.

VOLUME OF PRODUCTS ISSUED AND PATIENT COUNTS BY TREATMENT REGIMEN, SEVERITY, PRODUCT AND STATE

Table 22 shows the volumes issued by product and treatment regimen, for hereditary HMA, HMB, VWD. In both the adult and paediatric (includes adolescents) age groups the majority of product is issued for patients on prophylactic treatment regimens. The ABDR product issues data contains records where the treatment regimen is blank, unknown and not specified.

Table 23 and Table 24 show further breakdowns by whether patients have or do not have inhibitors.

Table 25 shows the volumes issued by product and treatment regimen, for diagnoses other than HMA, HMB, VWD.

Table 26 and Table 27 show the number of patients and volumes issued by product and state. The totals are distinct counts of patients who received product. A patient may be counted more than once under each state as they may have received product from more than one state throughout the year. This applies to both hereditary and acquired HMA, HMB and VWD.

Table 28 shows the number of patients, volume issued and IU or mg/kg/year of products issued in 2018-19 by treatment regimen for hereditary HMA, HMB and VWD.

Table 29 shows the number of patients and IUs issued by severity and regimen type for hereditary HMA and HMB. Values in this table exclude products issued to patients with unknown severity classification or treatment regimen, so they will vary from those figures shown in other parts of this report. Also, patients may receive more than one regimen type and may therefore be counted multiple times.

Table 30 shows the number of patients and volume of products issued by regimen type and product for hereditary HMA, HMB and VWD. Values in this table exclude products issued to patients with unknown treatment regimen, so they may vary from those figures shown in other parts of this report. Also, patients may receive more than one regimen type and may therefore be counted multiple times.

TABLE 22 - VOLUME (IU) OF PRODUCTS ISSUED IN 2018-19 (INCLUDING EHL PRODUCTS) BY TREATMENT REGIMEN - HEREDITARY BLEEDING DISORDERS

	Adult					Paediatric				
	On Demand	Prophylaxis	Tolerisation	Not specified	Adult Total*	On Demand	Prophylaxis	Tolerisation	Not specified	Paediatric Total*
HMA (IUs)	20,200,050	104,570,250	1,712,500	592,500	127,075,300	1,006,000	52,203,750	6,773,750	38,750	60,022,250
Advate	10,370,300	35,773,500	1,218,500	140,500	47,502,800	784,000	20,838,250	1,317,750	37,750	22,977,750
Biostate	824,000	2,174,000			2,998,000	3,000	5,955,000	5,281,500		11,239,500
*FEIBA (Units)	1,412,500	80,000			1,492,500		1,240,500	174,500		1,415,000
**NovoSeven (mgs)	4,367	2,842	70		7,279	1,491	723	369		2,583
Xyntha	7,593,250	44,938,750	494,000	118,500	53,144,500	149,000	14,028,250			14,177,250
Adynovate		16,013,000		333,500	16,346,500	70,000	6,986,750		1,000	7,057,750
Eloctate		5,591,000			5,591,000		3,155,000			3,155,000
HMB (IUs)	7,079,502	14,653,500	488,000	117,500	22,338,502	880,500	4,457,500		38,250	5,376,250
BeneFIX	6,347,500	7,313,000		70,500	13,731,000	642,000	1,392,000		36,500	2,070,500
Factor XI bpl	2				2					
MonoFIX	75,000		488,000		563,000					
**NovoSeven (mgs)	1,782	120			1,902	2				2
Rixubis	288,000			20,000	308,000	38,000	58,000			96,000
Xyntha		50,000		27,000	77,000					
Alprolix	369,000	7,290,500			7,659,500	200,500	3,007,500		1,750	3,209,750
VWD (IUs)	2,707,250	3,363,500		71,500	6,142,250	196,750	535,750	465,000	13,000	1,210,500
Biostate	2,707,250	3,363,500		71,500	6,142,250	196,750	535,750	465,000	13,000	1,210,500

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 23 - PATIENT NUMBERS AND VOLUME (IU) OF PRODUCTS ISSUED IN 2018-19 BY TREATMENT REGIMEN - HEREDITARY BLEEDING DISORDERS – ADULT PATIENTS

Table 23 shows product issued in 2018-19 by treatment regimen and patients with or without inhibitors for hereditary bleeding disorders. ‘With inhibitors’ means patients had an inhibitor status of Currently present – not on ITT, On ITT or Present as the final inhibitor status for 2018-19. ‘Without inhibitors’ means patients had an inhibitor status of Never Present or Negative as the final inhibitor status for 2018-19. The total rows (eg FVIII products) include all inhibitor statuses.

	Adult									
	On Demand		Prophylaxis		Tolerisation		Not specified		Adult Total *	
Hereditary	No pts	Total IU	No pts	Total IU	No pts	Total IU	No pts	Total IU	No pts	Total IU
HMA (Total IU Products)†										
FVIII products (all inhibitor statuses)	336	18,787,550	290	82,886,250	<5	1,712,500	17	259,000	646	103,645,300
Patients with inhibitors	9	873,000	<5	572,000	<5	1,712,500			14	3,157,500
Patients without inhibitors	273	16,145,250	236	65,435,000			13	230,500	522	81,810,750
*FEIBA (Units) (all inhibitor statuses)	5	1,412,500	<5	80,000					6	1,492,500
Patients with inhibitors	5	1,412,500	<5	80,000					6	1,492,500
Patients without inhibitors										
**NovoSeven (mgs) (all inhibitor statuses)	24	4,367	6	2,842	<5	70			31	7,279
Patients with inhibitors	14	2,836	5	2,682	<5	70			20	5,588
Patients without inhibitors	<5	264							<5	264
HMB (Total IU Products)‡										
FIX products (all inhibitor statuses)	110	6,710,500	37	7,153,000	<5	488,000	<5	90,500	152	14,442,000
Patients with inhibitors			<5	529,000	<5	488,000			<5	1,017,000
Patients without inhibitors	76	5,439,000	32	5,805,000			<5	32,000	110	11,276,000
Factor XI bpl (all inhibitor statuses)	<5	2							<5	2
Patients with inhibitors										
Patients without inhibitors	<5	2							<5	2
*NovoSeven (mgs) (all inhibitor statuses)	<5	1,782	<5	120					<5	1,902
Patients with inhibitors			<5	120					<5	120
Patients without inhibitors										
Xyntha (all inhibitor statuses)			<5	50,000			<5	27,000	<5	77,000
Patients with inhibitors										
Patients without inhibitors			<5	50,000			<5	27,000	<5	77,000
VWD (Total IU Products)										
FVIII products (all inhibitor statuses)	159	2,707,250	18	3,363,500			18	71,500	195	6,142,250
Patients with inhibitors			<5	121,000					<5	121,000
Patients without inhibitors	15	797,000	6	1,283,000					21	2,080,000

TABLE 23 CONTINUED- PATIENT NUMBERS AND VOLUME (IU) OF PRODUCTS ISSUED IN 2018-19 BY TREATMENT REGIMEN - HEREDITARY BLEEDING DISORDERS – PAEDIATRIC PATIENTS

	Paediatric									
	On Demand		Prophylaxis		Tolerisation		Not specified		Paediatric Total *	
Hereditary	No pts	Total IU	No pts	Total IU	No pts	Total IU	No pts	Total IU	No pts	Total IU
HMA (Total IU Products)†										
FVIII products (all inhibitor statuses)	89	936,000	230	40,821,500	15	6,599,250	7	37,750	341	48,394,500
Patients with inhibitors	<5	61,250	8	3,779,250	13	6,561,750	<5	14,750	23	10,417,000
Patients without inhibitors	43	593,250	143	22,458,000			<5	12,500	187	23,063,750
FEIBA (Units) (all inhibitor statuses)			6	1,240,500	<5	174,500			9	1,415,000
Patients with inhibitors			6	1,240,500	<5	174,500			9	1,415,000
Patients without inhibitors										
**NovoSeven (mgs) (all inhibitor statuses)	<5	1,491	<5	723	7	369			13	2,583
Patients with inhibitors	<5	1,491	<5	723	7	369			13	2,583
Patients without inhibitors										
HMB (Total IU Products)‡										
FIX products (all inhibitor statuses)	29	680,000	23	1,450,000			<5	36,500	55	2,166,500
Patients with inhibitors	<5	125,250					<5	6,000	<5	131,250
Patients without inhibitors	12	140,750	18	1,276,750			<5	30,000	31	1,447,500
Factor XI bpl (all inhibitor statuses)										
Patients with inhibitors										
Patients without inhibitors										
*NovoSeven (mgs) (all inhibitor statuses)	<5	2							<5	2
Patients with inhibitors										
Patients without inhibitors	<5	2							<5	2
Xyntha (all inhibitor statuses)										
Patients with inhibitors										
Patients without inhibitors										
VWD (Total IU Products)										
FVIII products (all inhibitor statuses)	27	196,750	6	535,750	<5	465,000	<5	13,000	36	1,210,500
Patients with inhibitors	<5	16,000			<5	465,000			<5	481,000
Patients without inhibitors	<5	15,000	<5	351,500					7	366,500

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 24 - PATIENT NUMBERS AND VOLUME (IU) OF PRODUCTS ISSUED IN 2018-19 BY TREATMENT REGIMEN - ACQUIRED BLEEDING DISORDERS

Table 24 shows product issued in 2018-19 by treatment regimen and patients with or without inhibitors for acquired bleeding disorders. ‘With inhibitors’ means patients had an inhibitor status of Currently present – not on ITT, On ITT or Present as the final inhibitor status for 2018-19. ‘Without inhibitors’ means patients had an inhibitor status of Never Present or Negative as the final inhibitor status for 2018-19. The total rows (eg FVIII products) include all inhibitor statuses.

	Adult									
	On Demand		Prophylaxis		Tolerisation		Not specified		Adult Total *	
	Number of patients	Total IU								
Acquired										
HMA (Total IU Products)†										
FVIII products (all inhibitor statuses)							<5	99,000	<5	99,000
Patients with inhibitors										
Patients without inhibitors										
FEIBA (Units) (all inhibitor statuses)	<5	10,500							<5	10,500
Patients with inhibitors										
Patients without inhibitors										
**NovoSeven (mgs) (all inhibitor statuses)	<5	590					10	3,701	13	4,291
Patients with inhibitors	<5	290					5	2,537	7	2,827
Patients without inhibitors										
VWD (Total IU Products)										
FVIII products	6	677,500					<5	126,250	10	803,750
Patients with inhibitors	<5	4,500							<5	4,500
Patients without inhibitors	<5	596,000							<5	596,000
**NovoSeven (mgs)	<5	30							<5	30
Patients with inhibitors										
Patients without inhibitors	<5	30							<5	30

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient’s condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 25 - VOLUME (IU) OF PRODUCTS ISSUED IN 2018-19 BY TREATMENT REGIMEN - OTHER DIAGNOSES

	Adult				Paediatric			
	On Demand	Prophylaxis	Not specified	Adult Total *	On Demand	Prophylaxis	Not specified	Paediatric Total *
Other Factor Deficiency	328,629	221,398	25,501	575,528	24,002	38,750		62,752
BeneFIX					2,000			2,000
Factor XI bpl	8,629		25,501	34,130	2			2
FEIBA (Units)	311,000			311,000				
Fibrogammin	2,500	5,000		7,500		38,750		38,750
*NovoSeven (mgs)	564	688	16	1,268		175		175
NovoThirteen		127,898		127,898				
Prothrombinex - VF	6,500	88,500		95,000	22,000			22,000
Platelet Disorder	106		7	113	151			151
*NovoSeven (mgs)	106		7	113	151			151
Fibrinogen	215	105	12	332	42	106		148
Human Fibrinogen RiaSTAP (gms)	215	105	12	332	42	106		148
Other	102,500		16,000	118,500				
Biostate	4,000		16,000	20,000				
FEIBA (Units)	98,500			98,500				
*NovoSeven (mgs)	78			78				

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 26 - NUMBER OF PATIENTS FOR HEREDITARY HMA, HMB AND VWD BY STATE

	Number of Patients who received product during the year								
	ACT	NSW	NT	QLD	SA	TAS	VIC	WA	Total*
HMA									
Advate	20	173	5	149	90	11	92	44	579
Biostate	<5	21	<5	12	<5		6	5	47
FEIBA		<5		6		<5	<5	<5	13
NovoSeven		<5		9	7	<5	15	<5	39
Xyntha	5	95	<5	40	20	13	102	65	334
HMB									
BeneFIX	6	56		57	14	<5	44	13	190
Factor XI bpl								<5	<5
MonoFIX	<5	<5		<5					<15
NovoSeven		<5					<5		<10
Rixubis		<5		<5			<5	<5	8
Trial Material		<5							<5
Xyntha	<5	<5							<10
VWD									
Biostate	8	62	<5	65	16	6	26	45	229

* The Totals are distinct counts of Patients who received product and may be counted more than once under each state or across different states as they may have received more than one product or been treated in more than one state throughout the year.

TABLE 27 - VOLUME (IU) OF PRODUCT ISSUED FOR HEREDITARY HMA, HMB AND VWD BY STATE

	Volume of Product Issued through the year								
	ACT	NSW	NT	QLD	SA	TAS	VIC	WA	Total
HMA (IUs)†	1,586,500	57,408,550	690,500	34,537,000	13,231,000	4,973,500	27,892,250	14,628,000	154,947,300
Advate	972,000	27,057,550	684,250	19,099,000	7,954,250	1,171,000	11,354,500	2,188,000	70,480,550
Biostate	6,000	6,387,500	3,500	6,096,000	8,000		1,457,500	279,000	14,237,500
FEIBA		780,000		1,165,500		108,000	574,000	280,000	2,907,500
**NovoSeven (mgs)		80		2,052	646	120	6,132	832	9,862
Xyntha	608,500	23,183,500	2,750	8,176,500	5,268,750	3,694,500	14,506,250	11,881,000	67,321,750
HMB (IUs)‡	829,000	7,544,250		2,982,750	851,500	407,000	3,584,500	646,502	16,845,502
BeneFIX	291,000	7,464,250		2,728,750	851,500	407,000	3,488,500	570,500	15,801,500
Factor XI bpl								2	2
MonoFIX	488,000	33,000		42,000					563,000
**NovoSeven (mgs)		122					1,782		1,904
Rixubis		20,000		212,000			96,000	76,000	404,000
Xyntha	50,000	27,000							77,000
VWD (IUs)	341,500	3,882,250	16,500	1,380,250	388,250	35,500	348,250	960,250	7,352,750
Biostate	341,500	3,882,250	16,500	1,380,250	388,250	35,500	348,250	960,250	7,352,750

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 28 - VOLUME (IU), PATIENT COUNTS AND IU OR MG/KG/YEAR OF PRODUCTS ISSUED IN 2018-19 BY TREATMENT REGIMEN - HEREDITARY

Table 28 shows volume of product issued by IU/mg/units, number of patients and Average IU/kg for the 2018-19 year, by treatment regimen.

	Adult											
	On Demand			Prophylaxis			Tolerisation			Not specified*		
	No of patients	Total Units	Avg. IU/KG	No of patients	Total Units	Avg. IU/KG	No of patients	Total Units	Avg. IU/KG	No of patients	Total Units	Avg. IU/KG
HMA (IUs)	339	18,787,550	395	292	82,886,250	1,137	<5	1,712,500	469	17	259,000	101
Advate	242	10,370,300	95	130	35,773,500	273	<5	1,218,500	101	11	140,500	59
Biostate	18	824,000	109	6	2,174,000	592						
FEIBA	5	1,412,500	153	<5	80,000	201						
*NovoSeven (mgs)	24	4,367		6	2,842		<5	70				
Xyntha	79	7,593,250	191	156	44,938,750	272	<5	494,000	368	6	118,500	42
HMB (IUs)	113	6,710,502	377	39	7,363,000	644	<5	488,000	185	5	117,500	633
BeneFIX	104	6,347,500	131	38	7,313,000	252				<5	70,500	176
Factor XI bpl	<5	2										
MonoFIX - VF	<5	75,000	74				<5	488,000	185			
NovoSeven	<5	1,782	1	<5	120							
Rixubis	6	288,000	172							<5	20,000	
Xyntha				<5	50,000	392				<5	27,000	457
VWD (IUs)	159	2,707,250	59	18	3,363,500	150				19	71,500	39
Biostate	159	2,707,250	59	18	3,363,500	150				19	71,500	39

* The total in this table combines the values for patients with mild, moderate and severe conditions. The severity of a patient's condition is not always known at initial presentation. This table includes product issues to patients with unknown/not specified treatment regimens. All products listed above are in IUs unless stated.

**IUs sums all the products except NovoSeven.

TABLE 29 - VOLUME (IU) OF PRODUCT ISSUED AND PATIENT COUNTS FOR HEREDITARY HMA AND HMB BY SEVERITY AND REGIMEN TYPE

Severity*	Regimen type**	Haemophilia A		Haemophilia B	
		Total IUs	Number of patients	Total IUs	Number of patients
Adult - Mild	On demand	4,172,500	186	1,882,502	55
	Prophylaxis	1,314,500	9	311,500	<5
	Tolerisation	494,000	<5		
Adult - Moderate	On demand	3,487,250	64	3,313,500	40
	Prophylaxis	9,070,250	34	3,379,000	15
Adult - Severe	On demand	11,127,800	86	1,880,000	16
	Prophylaxis	94,105,500	292	10,963,000	45
	Tolerisation	1,218,500	<5	488,000	<5
Adult - Total		124,990,300		22,217,502	
Paediatric - Mild	On demand	384,250	44	127,250	12
	Prophylaxis	190,750	5		
Paediatric - Moderate	On demand	261,500	25	533,000	14
	Prophylaxis	3,615,500	25	573,000	7
Paediatric - Severe	On demand	360,250	20	220,250	<5
	Prophylaxis	47,157,000	238	3,884,500	35
	Tolerisation	6,599,250	15		
Paediatric - Total		58,568,500		5,338,000	

* Values in this table exclude products issued to patients with unknown severity classification or treatment regimen, so they will vary from those figures shown previously.

**Patients may receive more than one regimen type and may therefore be counted multiple times.

TABLE 30 - VOLUME (IU) OF PRODUCT ISSUED AND PATIENT COUNTS FOR HEREDITARY HMA, HMB AND VWD BY REGIMEN TYPE AND PRODUCT

Bleeding Disorder	Paediatric / Adult	Regimen type	Advate		BeneFIX		Biostate		Factor XI bpl		FEIBA	
			Total IUs	Number of patients	Total IUs	Number of patients	Total IUs	Number of patients	Total IUs	Number of patients	Total Units	Number of patients
Haemophilia A	Adult	On demand	10,370,300	242			824,000	18			1,412,500	5
		Prophylaxis	35,773,500	130			2,174,000	6			80,000	<5
		Tolerisation	1,218,500	<5								
	Paediatric	On demand	784,000	66			3,000	<5				
		Prophylaxis	20,838,250	137			5,955,000	16			1,240,500	6
		Tolerisation	1,317,750	7			5,281,500	8			174,500	<5
	Total		70,302,300	<587			14,237,500	<53			2,907,500	<21
Haemophilia B	Adult	On demand			6,347,500	104			2	<5		
		Prophylaxis			7,313,000	38						
		Tolerisation										
	Paediatric	On demand			642,000	28						
		Prophylaxis			1,392,000	23						
	Total				15,694,500	193			2	<5		
Von Willebrand Disease	Adult	On demand					2,707,250	159				
		Prophylaxis					3,363,500	18				
	Paediatric	On demand					196,750	27				
		Prophylaxis					535,750	6				
		Tolerisation					465,000	<5				
	Total					7,268,250	<215					

*Values in this table exclude products issued to patients with unknown treatment regimen, so they will vary from those figures shown previously.

**Patients may receive more than one regimen type and may therefore be counted multiple times.

TABLE 30 CONTINUED - VOLUME (IU) OF PRODUCT ISSUED AND PATIENT COUNTS FOR HEREDITARY HMA, HMB AND VWD BY REGIMEN TYPE AND PRODUCT

Bleeding Disorder	Paediatric / Adult	Regimen type	MonoFIX - VF		NovoSeven		Rixubis		Xyntha	
			Total IUs	Number of patients	Total mgs	Number of patients	Total IUs	Number of patients	Total IUs	Number of patients
Haemophilia A	Adult	On demand			4,367	24			7,593,250	79
		Prophylaxis			2,842	6			44,938,750	156
		Tolerisation			70	<5			494,000	<5
	Paediatric	On demand			1,491	<5			149,000	22
		Prophylaxis			723	<5			14,028,250	80
		Tolerisation			369	7				
	Total				9,862	44			67,203,250	<342
Haemophilia B	Adult	On demand	75,000	<5	1,782	<5	288,000	6		
		Prophylaxis			120	<5			50,000	<5
		Tolerisation	488,000	<5						
	Paediatric	On demand			2	<5	38,000	<5		
		Prophylaxis					58,000	<5		
	Total		563,000	<10	1,904	<15	384,000	<16	50,000	<5
Von Willebrand Disease	Adult	On demand								
		Prophylaxis								
	Paediatric	On demand								
		Prophylaxis								
		Tolerisation								
	Total									

*Values in this table exclude products issued to patients with unknown treatment regimen, so they will vary from those figures shown previously.

**Patients may receive more than one regimen type and may therefore be counted multiple times. Patient totals are the total number of distinct patients, excluding patients which are counted multiple times, so may not match the individual values.

Appendix A Characteristics of Rare Clotting Factor Deficiencies

TABLE 31 - CHARACTERISTICS OF RARE CLOTTING FACTOR DEFICIENCIES

Missing Factor	Incidence*	Inheritance	Severity of Bleeding	Treatment
Factor I Afibrinogenemia Hypofibrinogenemia Dysfibrinogenemia	5 in 10 million not available 1 in 1 million	Autosomal recessive Recessive or dominant Recessive or dominant	Usually mild, except in afibrinogenemia	<ul style="list-style-type: none"> •Fibrinogen conc. (Not funded in Australia) •Cryoprecipitate •Fresh frozen plasma
Factor II	1 in 2 million	Autosomal recessive	Moderate to severe when factor levels are low; usually mild	<ul style="list-style-type: none"> •Prothrombin complex conc. •Fresh frozen plasma
Factor V	1 in 1 million	Autosomal recessive	Moderate to severe when factor levels are low; usually mild	<ul style="list-style-type: none"> •Fresh frozen plasma
Combined Factor V and Factor VIII	1 in 1 million†	Autosomal recessive‡	Usually mild	<ul style="list-style-type: none"> •Fresh frozen plasma •Factor VIII conc. •Desmopressin
Factor VII	1 in 500,000	Autosomal recessive	Severe when factor levels are low	<ul style="list-style-type: none"> •Recombinant Factor VIIa conc. •Factor VII conc. •Fresh frozen plasma
Factor X	1 in 1 million	Autosomal recessive	Moderate to severe when factor levels are low	<ul style="list-style-type: none"> •Prothrombin complex conc. •Fresh frozen plasma
Combined deficiency of vitamin K-dependent clotting factors	not available	Autosomal recessive	Usually mild, but a few families have reported very low levels and more severe symptoms	<ul style="list-style-type: none"> •Vitamin K •Prothrombin complex conc. •Fresh frozen plasma
Factor XI	1 in 100,000	Recessive or dominant	Mild to moderate when factor levels are low	<ul style="list-style-type: none"> •Factor XI concentrate •Antifibrinolytic drugs •Fibrin glue •Fresh frozen plasma
Factor XIII	1 in 3 million	Autosomal recessive	Moderate to severe when factor levels are low	<ul style="list-style-type: none"> •Factor XIII conc. •Cryoprecipitate •Fresh frozen plasma

Note: Australian Prothrombin Complex Concentrate is not used for FVII deficiency

* Estimates only

† 1 in 100,000 in some populations, including Israel, Iran, and Italy

‡ Very rarely, Factor VIII deficiency can be inherited separately from only one parent

Appendix B Haemophilia Treatment Centres

THE OBJECTIVES OF HTCS

Haemophilia Treatment Centres provide comprehensive care for people with haemophilia. Their roles include:

- Compilation and distribution of guidelines for best practice directed toward optimal care of patients with disorders of haemostasis
- Providing protocols for the accurate diagnosis of patients with bleeding disorders
- Providing protocols for the regular review of infectious disease markers in patients and their families
- The allocation and distribution of therapeutic blood and recombinant products together with advice regarding the use of blood and recombinant products, at a state and national level
- The establishment of review programs to assess outcomes of therapies
- Provision of regularly updated data to the national Haemophilia Registry (ABDR)
- Participation in basic and clinical research

OPERATING CONCEPT

Haemophilia Treatment Centres coordinate and, where possible, integrate patient care, research and education to provide the optimal use of expertise and resources within hospitals and the community. One collaborative centre for each state and territory may suffice but this must include adult and paediatric type centres.

Haemophilia Centres provide:

- a single point of accountability for the care of patients with bleeding disorders with responsibility for the coordination, allocation and distribution of therapeutic resources for the treatment of patients, i.e. coagulation products derived either from blood donors or recombinant technologies
- a clinical service by experienced staff for patients with bleeding disorders and their families at short notice at any time of the day or night
- organisation of home therapy programs by the centre or in collaboration with other haemophilia treatment facilities
- a counselling and advisory service for people with haemophilia and their families including genetic counselling and family planning
- specialist medical expertise, principally haematology, surgery (the surgeons would have to be accredited to the Haemophilia Centre) rheumatology, infectious diseases and dental services
- specialist allied health services to include physiotherapy, social work and podiatry
- a laboratory service able to carry out all investigations required for the accurate diagnosis of haemophilia and other inherited disorders of haemostasis and to have access, in association with other centres, to specialised testing facilities, for example gene typing
- a system of record for all investigations, treatments, allocation of therapeutic products and adverse reactions
- a capability to participate in research including clinical trials
- educational programs for medical staff, other personnel, patients and their families which promote care of patients with disorders of haemostasis

- an outreach service to isolated patients and treating medical services. The outreach service may include:-
 - A haemophilia treatment facility located in a hospital that does not provide all the specialist services
 - Designated supervising medical practitioner
- data management to facilitate the use of an information system database, such as the Australian Bleeding Disorders Registry, used in the clinical environment to aid in the capturing of data critical to HTC staff for the day to day management of people with bleeding disorders and for supply management and policy purposes.

All isolated patients (where care is managed in an outreach program) should be registered at, and be reviewed regularly by, a Haemophilia Treatment Centre which would arrange delivery of and monitor the supply of therapeutic coagulation products.

The HTC must maintain on-going dialogue with the client group in each state and territory. The role of State and Territory Governments is to designate 'Haemophilia Treatment Centres' and negotiate the funding of the HTC including the purchase of therapeutic blood and recombinant products for distribution within states (or regions) and territories. In some states committees have been established to consider and schedule elective surgery.

DATA QUALITY OF HTC DATA COLLECTIONS

The following organisations are represented at various HTCs nationally:

- Australian Haemophilia Nurses Group (AHNG)
- Australia/New Zealand Haemophilia Social Workers' and Counsellors' Group (ANZHSWCG)
- Australia/New Zealand Physiotherapist Group (ANZHPG)
- Haemophilia Foundation Australia (HFA).

These member representatives have provided input into the initial design of the ABDR and continue to provide input from their respective areas of speciality.

The Data Managers at each HTC are members of the Data Managers' Group (DMG). DMG Co-Chairs are elected and coordinate teleconferences, between all Data Managers, on a regular basis. The DMG Co-Chairs also have the functionality of raising issues, to the NBA, on behalf of their group. AHCDO has a major role in providing support to ABDR Data Managers through the funded model for Data Managers.

The advantages of this model of Haemophilia Data Co-ordination are:

- Accurate and complete data entry
- Dedicated and focused data management
- Regular reporting and analysis of collated information
- New product initiation of unresolved haemophilia care related questions
- Clinical audit of current policies and monitoring of agreed national standards.

A number of ongoing data quality initiatives were first implemented in 2010-11, including:

- Regular teleconferences for ABDR DMG
- 'Advanced Search' functionality of the ABDR whereby Data Managers are able to extract information from the ABDR on an ad hoc basis
- Reviews of data definitions undertaken by DMG Co Chairs
- NBA financial support, through AHCDO funding, for HTC Data Managers
- The ABDR Update is a functional tool in the form of a Newsletter. This provides an update on issues such as changes to the ABDR and functionality enhancements. This update is a means of keeping all ABDR stakeholders informed.

Comprehensive automated and manual data cleansing and validation processes that occurred as part of the 4th Generation ABDR Redevelopment project released in August 2012 enhanced the ABDR data accuracy and consistency presented in this report. The 4th Generation ABDR was successfully implemented on 13 August 2012.

However, there are still some data quality issues that impact the data presented in this report. Some post migration tasks for Data Managers to clean the data include:

- Verify patients with more than one diagnosis
- Duplicate diagnoses to be deleted and Inhibitor Tests to be combined under the persisting diagnosis
- Verify severity ratings and treatment regimens for some patient records
- There are also a number of low level data verification activities.

DATA PROJECTS

As data quality improves, various data projects are able to be undertaken to provide insights into further opportunities for improvement in data entry, or additional information to assist with managing patients and treatments. The following projects were progressed during 2018-19:

- Hepatitis C project – this project is looking at the prevalence of Hepatitis C (HCV) among patients with a bleeding disorder and the impact of subsidised medication for HCV. Results of the project show that treatment uptake has improved dramatically. Data gaps are being followed up to improve ongoing data quality.
- SIPPET (Survey of Inhibitors in Plasma-Products Exposed Toddlers) project – this project included Previously Untreated Patients (PUPs) born between 2011 and 2017. There was little change in prescribing practice in terms of product choice.
- Genetic Landscape Project – a review of the genetic profile of patients with bleeding disorders and the correlation between particular types of mutations and the risk of inhibitor development. Intron Inversion, Large Deletion and Frameshift Mutation were the groups that were most likely to develop inhibitors.
- Switch Project – 857 Haemophilia A patients switched from one recombinant FVIII product to another. The results indicate that switching products did not increase the risk of inhibitor development, however switching between product types may impact inhibitor development.
- Inhibitor Project – 24.9% of severe patients developed an inhibitor. Overall development of inhibitors was 17.5%. After more than 50 exposure days, the risk decreases drastically.
- EHL Project – the aim of this project was to look at medical factors around EHL product use. The most common products used were Eloctate and Adynovate. The most popular regimen for Haemophilia A patients was 41-50 IU per kg twice weekly, and for Haemophilia B patients 41-50 IU per kg once a week. Overall bleed rates decreased tremendously, and the proportion of patients with no bleeds increased significantly (44% to 64%).

These projects will continue over time and other data analysis projects will also be undertaken and reported in future years.

LIST OF HTCS

TABLE 32 - HAEMOPHILIA TREATMENT CENTRES

Hospital	Haemophilia Treatment Centre	State
The Canberra Hospital	Haemophilia Clinic	ACT
Calvary Mater Newcastle	Haemophilia Treatment Centre	NSW
Royal Prince Alfred Hospital	Haemophilia Treatment Centre	NSW
Sydney Children's Hospital	Centre for Children's Cancer and Blood Disorders	NSW
The Children's Hospital at Westmead	Haemophilia Treatment Centre	NSW
Prince of Wales Hospital	Bleeding Disorders Clinic	NSW
Westmead Hospital	Bleeding Disorders Clinic	NSW
Royal Darwin Hospital	Haemophilia Treatment Centre	NT
Royal Brisbane and Women's Hospital	Queensland Haemophilia Centre	QLD
Queensland Children's Hospital	Queensland Haemophilia Centre Child and Adolescent Service	QLD
Royal Adelaide Hospital	South Australia Haemophilia Treatment Centre	SA
Women's and Children's Hospital	South Australia Haemophilia Treatment Centre	SA
Royal Hobart Hospital	Tasmanian Haemophilia Treatment Centre	TAS
The Alfred Hospital	Ronald Sawyers Haemophilia Centre	VIC
Royal Children's Hospital	Henry Ekert Haemophilia Treatment Centre	VIC
Perth Children's Hospital	Paediatric Haemophilia Centre	WA
Hollywood Private Hospital	Hollywood Hospital Haemophilia Treatment Centre	WA
Fiona Stanley Hospital	Adult Haemophilia Centre	WA

Appendix C National Supply of Products

It is the responsibility of the NBA to manage the national blood supply to ensure that healthcare providers have sustainable, reliable and efficient access to blood and blood products needed for patient care. The NBA ensures blood supply security by working with states and territories to determine and manage an annual supply plan and budget and negotiating and managing blood supply contracts and arrangements with local and overseas suppliers.

NATIONAL SUPPLY PLAN AND BUDGET

A key element of the NBA's role in ensuring security of supply is to develop, coordinate and monitor the annual national supply plan and budget, including obtaining annual approval from health ministers.

This is achieved by:

- developing a national estimate of product demand
- liaising with jurisdictions and stakeholders to refine the estimated demand for products
- collecting and distributing data on product issued and reporting variations to jurisdictions on the approved supply plan
- intensively managing products if they are in short supply.

Figure 19 illustrates the national supply by product category for 2018-19, and shows issues of clotting factor products was 13.7% (\$168.1 million).




FIGURE 19 - NATIONAL ISSUES BY PRODUCT CATEGORY 2018-19

Note: Plasma for Fractionation costs paid to the Blood Service for collection has been attributed to IVIg and Hyperimmunes




FIGURE 20 - EXPENDITURE ON CLOTTING FACTORS AND PERCENTAGE OF BLOOD BUDGET 2009-10 TO 2018-19

Figure 20 illustrates the variations in total expenditure on clotting factors and the percentage of the blood and blood products budget that comprised each year for 2009-10 to 2018-19. It also shows that the number of patients who received products has grown significantly over the 10 years to 2018-19. Overall expenditure is down over the 10 year period, remaining relatively steady in the last 5 years. Contract negotiation processes have led to falls in average costs per IU from 2012-13 to 2018-19.

Throughout 2018-19, products were supplied to meet clinical demand and supply risks were effectively managed. The approved budget for 2018–19 covering the supply and management of blood and blood products and services under contract was \$1,259.48 million, comprising \$677.23 million for fresh blood products and plasma collection and \$569.2 million for plasma and recombinant products. The remaining \$13.04 million included items such as support for the publication of PBM Guidelines, maintenance of the Australian Haemophilia Centre Directors' Organisation (AHCDO) and administration of the Australian Bleeding Disorders Registry (ABDR).

ISSUES OF CLOTTING FACTORS

Issues of clotting factor products represent those deliveries from suppliers to all Australian Health Providers, including hospitals and Haemophilia Treatment Centres.

Figure 21 indicates that the demand for Factor VIII products in 2018-19 increased by 6.7 per cent when compared to 2017-18. The demand for recombinant Factor VIII increased by 8.3 per cent over 2017-18. Plasma derived Factor VIII demand decreased by 3.6 per cent. Patient participation in company clinical trials for recombinant Factor VIII products continues to contribute to the variability of year-to-year product growth.




FIGURE 21 - ISSUES OF FACTOR VIII PRODUCTS, 2014-15 TO 2018-19 PER '000 POPULATION

Figure 22 indicates that demand for Factor IX products in 2018-19 decreased by 12.6 per cent compared to 2017-18. Plasma derived Factor IX demand decreased by 1.3 per cent in 2018-19 due to a reduction in specific patient requirements. Demand for Recombinant Factor IX decreased by 12.8 per cent in 2018-19. Patient participation in company clinical trials for recombinant Factor IX products continues to contribute to the variability of year-to-year product growth.




FIGURE 22 - ISSUES OF FACTOR IX PRODUCTS, 2014-15 TO 2018-19 PER '000 POPULATION

Figure 23 and Figure 24 show demand for Recombinant Factor VIIa decreased by 0.8 per cent and increased 17.3 per cent for FEIBA compared to 2017-18. Demand for Recombinant Factor VIIa and FEIBA can change significantly from year to year as a result of the variable needs of a small number of patients.




FIGURE 23 - ISSUES OF RECOMBINANT FACTOR VIIA PRODUCTS, 2014-15 TO 2018-19 PER '000 POPULATION




FIGURE 24 - ISSUES OF FEIBA, 2014-15 TO 2018-19 PER '000 POPULATION

SUPPLY OF EXTENDED HALF LIFE PRODUCTS

In 2018 the NBA implemented limited interim arrangements to provide temporary access to EHL recombinant factor VIII and factor IX clotting factor products under the national supply arrangements for a limited number of haemophilia A and B patients with high priority needs.

These arrangements provide recombinant factor VIII and factor IX products from the supplier companies Shire and Bioverativ for approximately 200 patients, under the coordination and monitoring of the Australian Haemophilia Centre Directors' Organisation (AHCDO) by arrangement with the NBA.

These limited initial arrangements will remain in place pending the outcomes of the next national tender for clotting factors and related products.

Figure 25 shows patient numbers and uptake of EHL product use, which rose sharply in the May-July period of 2018 as patients transitioned from trial periods (under initial limited arrangements in place from December 2017) and commenced recording their usage of products. Note that this figure which shows treatments recorded by patients may differ slightly to Figure 25 which shows total quantity supplied.




FIGURE 25 - UPTAKE OF EHL PRODUCT USE DECEMBER 2017- JULY 2019

PATIENT SUITABILITY AND PRIORITISATION

Prioritisation criteria and other considerations were agreed by a Reference Group to ensure that EHL products were directed to those patients where the greatest benefit would be obtained. Patient suitability criteria are:

PRIORITY CRITERIA

1. Patients currently on extension programs for Adynovate, Eloctate or Alprolix in Australia. (These patients have been covered under the initial limited arrangements since December 2017).

2. Patients for whom infusion is currently accomplished using an infusion port or central line, which could be avoided by using an EHL product.
3. Severe or moderate haemophilia A or B patients who are currently adherent to recommended prophylactic therapy who nonetheless experience frequent bleeds, where this could be reduced or avoided by using an EHL product.
4. Severe or moderate haemophilia A or B patients where current patient care is likely to be substantially improved by using an EHL, because of (in descending priority):
 - a. improved adherence to recommended therapeutic regime
 - b. the opportunity to move from on-demand to prophylactic therapy
 - c. the opportunity to reduce current excessive use of clotting factor products
 - d. the opportunity for reduced dosing, or
 - e. the opportunity to support therapy with improved data recording in ABDR.

OTHER CONSIDERATIONS

1. Patients will not be considered suitable for participation where:
 - a. the patient has less than 50 exposure days to clotting factor therapy
 - b. the patient has a history of inhibitors to clotting factor therapy, or
 - c. data recording for the patient within ABDR/MyABDR is not possible, or is not satisfactorily maintained while the patient has access to EHL products under the limited interim arrangements.
2. In addition, a patient may not be considered suitable for participation, or may have their participation reconsidered, where:
 - a. the patient's clinician does not consider that clinical benefit will be obtained, or considers that clinical benefit is not being demonstrated, by the patient having access to EHL products under the limited interim arrangements, or
 - b. the patient is not able to or chooses not to attend ongoing monitoring and review appointments as determined by the treating HTC clinician.

CHRONOLOGY OF PRODUCTS SUPPLIED

Various products have been supplied through national arrangements. Since 2009-10 the following arrangements for the supply of products have occurred.

2009-10	<ul style="list-style-type: none"> Commenced supply of Flebogamma
2010-11	<ul style="list-style-type: none"> Ceased supply of Sandoglobulin
2011-12	<ul style="list-style-type: none"> Ceased supply of WinRho Commenced supply of Kogenate
2012-13	<ul style="list-style-type: none"> Ceased supply of Flebogamma, vFVIII/Recombinate and vFVIII/Advate Commenced supply of Kiovig, Rhophylac, Normal Immunoglobulin, CMV Immunoglobulin, Hepatitis B Immunoglobulin, Tetanus Immunoglobulin and Zoster Immunoglobulin
2013-14	<ul style="list-style-type: none"> Commenced supply of Evogam and Gammanorm
2014-15	<ul style="list-style-type: none"> Commenced supply of Advate and Rixubus
2015-16	<ul style="list-style-type: none"> Ceased supply of Factor VII Concentrate Commenced supply of RiaSTAP, Flebogamma DIF, Privigen and Hizentra
2016-17	<ul style="list-style-type: none"> Ceased supply of Kogenate FS, Gammanorm and Octagam Commenced supply of Berinert Intragam P transitioned to Intragam 10
2017-18	<ul style="list-style-type: none"> Commenced supply of Eloctate, Alprolix, Adynovate and Novo Thirteen

Appendix D History of the ABDR

The ABDR was first established in 1988 using a 'Paradox' database at each Haemophilia Treatment Centre in Australia. The aims of the ABDR were to provide a clinical tool for improved management and national demographics of patients with haemophilia and other inherited bleeding disorders.

The first demographic Haemophilia registry was established by the Haemophilia Foundation Australia (HFA), under auspices of the Medical Advisory Panel (MAP), in 1991 with an initial survey of Haemophilia Treatment Centres (HTC) established in Australia. Following on this initial survey the MAP took on responsibility for developing an ongoing registry and database associated with a University. The registry was based on a Paradox database with a comprehensive data collection including demographics, factor usage and bleed data. It was intended that software would be updated regularly by circulation of floppy disc updates and annual reports produced. Issues identified included no dedicated data entry staff, variability of IT support in institutions, unstable database requiring significant maintenance, time for data entry, and complexity. Unfortunately the registry did not progress.

In view of issues identified, in 2000 a new database using Access was developed with a single initial page collecting demographic and basic clinical data – 'medical registry'. Financial support was provided for data entry. Identification was by a code including multiple initials of name and date of birth as used by National HIV registries in Australia. Duplicate entries were identified and individual HTCs were asked to resolve differences based on activity of PWH and HTC. Initial demographics and diagnoses were provided for an annual report – initially to Department of Health and Aging, subsequently to National Blood Authority and presented at various forums. Data was vital for identifying product needs of the PWH community at a time of introduction of recombinant products. The ABDR achieved Quality Assurance status with the Commonwealth to assist with concerns about privacy. Ongoing issues identified were related to privacy, data collection (with one state not being involved) and coverage of the database, and it appeared total product usage was not complete.

The National Blood Authority (NBA) was established in 2003 and in 2007 it was proposed to develop the ABDR further with a web based clinical registry. Funding from the NBA allowed updating of the database. Widespread consultation was undertaken with HTCs to draw up specifications for a clinical database. The project was tendered to a commercial provider to enable 'third party custody' of data. The ABDR was to be capable of ordering products in 'real time' at HTCs. Governance of the development and operation was by a steering committee consisting of Australian Haemophilia Centre Directors Organisation (AHCDO), HFA, NBA and jurisdictional representatives.

An internet-based, standardised data entry database involving all states was introduced in December 2008. But the database highlighted significant resource and IT issues in HTCs and hospitals with slow response and significant variation of practice within HTCs. This hampered complete data collection with lack of feedback to HTCs, inability to provide ad hoc reporting for HTCs and nationally available reports. Annual reports only provided broad information with NBA providing figures for factor usage. The commercial provider was unable to address these issues.

Issues with existing software and support by the commercial provider necessitated a different approach. Further funding from the NBA enabled redevelopment of the ABDR using industry standard software in a 'Like for like' development. Data is now being held within NBA – requiring strict security protocols and separation of staff analysing data from those managing the system. Deficiencies of previous software were addressed with development of online reports to assist HTC management. Further expansion to include data from physiotherapy and social work, counselling pages and adverse events were developed. The 4th generation ABDR was released on August 13, 2012.

The ABDR has evolved with improvements in technology and feedback from stakeholders. In 2014 the ABDR entered a new phase with MyABDR – a smartphone application to enable patient input of bleed data and factor usage directly to the ABDR. The ABDR project has improved communication between HTC's for transfers and knowledge of 'travellers'.

The NBA delivered a number of updates and improvements to the Registry in 2014-15 to enhance the functionality and the user experience with MyABDR. The innovation delivered by the patient portal to ABDR, MyABDR, was recognised by the ICT industry through the receipt of two national iAwards merits in the Health and Government categories in August 2014 and through ITnews naming the NBA's Chief Information Officer as 'Healthcare CIO of the Year' in February 2015.

There has been further identification of PWH and opportunity for standardisation of terminology. There is wide involvement of other professionals – nursing, physiotherapy, social workers/counselling. Adverse event reporting has commenced. Benchmarking between HTC's is possible with improvement in data recoding enabling opportunities for improvement.

BENEFITS OF THE 4TH GENERATION ABDR

The NBA redeveloped the ABDR and deployed the 4th generation ABDR on August 13, 2012. It provides the following benefits:

- Single point of access for clinicians for treatment of patients
- Patient information relating to all clinical information associated with the treatment of haemophilia
- Information exchange between states and Haemophilia Treatment Centres
- National demographic information (age, gender etc.) of persons with bleeding disorders
- National data on inhibitor incidence and outcomes of treatment
- Allied health (physiotherapy and social work) monitoring and outcomes
- Recording of personal usage of factor replacement for clinical monitoring
- Data for forward planning and funding of factor concentrates on a national basis
- MyABDR is a secure app for smartphones and web site for people with bleeding disorders or parents/caregivers to record home treatments and bleeds. As an alternative, there is also a MyABDR paper-based treatment diary.

CURRENT POSITION OF THE DEVELOPMENT OF THE ABDR

Today the Australian Bleeding Disorders Registry and MYABDR are fully operational. The ABDR Steering committee continues to oversee the project.

The National Blood Authority's role continues around provision of resources to maintain ABDR operations and to ensure timely and accurate reporting from the ABDR through provision of support to Data Managers. Data Managers, funded and supported by AHCD, are located at HTC's across Australia.

Appendix E Patient Registration Form

PATIENT REGISTRATION FORM			
Clinician/Nurse to complete. Fields marked with an *asterisk are mandatory, optional fields are shaded grey.			
<input type="checkbox"/> New patient <input type="checkbox"/> Change of name <input type="checkbox"/> Change of address			
Patient			
ABDR ID <small>(Existing patients only)</small>	Title	Australian Resident Status <small>(Please tick)</small>	
		<input type="checkbox"/> Australian Citizen/Permanent Resident <input type="checkbox"/> Overseas Visitor <input type="checkbox"/> Temporary Visa	
*First name	Second name / Initial	*Family name	
Known as / Alias	*Gender	*Date of birth	Previous family name/s
	<input type="checkbox"/> Male <input type="checkbox"/> Female	/ /	
*Address		*Suburb	
1		*State	
2		*Postcode	
3		Country	
<input type="checkbox"/> Home phone	<input type="checkbox"/> Work phone	<input type="checkbox"/> Mobile	*Tick preferred contact method; at least one contact must be supplied.
<input type="checkbox"/> Home email	<input type="checkbox"/> Work email		
Patient contact <small>(mandatory if patient is under 18)</small>			
<input type="checkbox"/> Mother <input type="checkbox"/> Father <input type="checkbox"/> Spouse <input type="checkbox"/> Grandparent <input type="checkbox"/> Emergency <input type="checkbox"/> Other Please specify: _____			
Title	First name	Second name / Initial	Last name
Address		Suburb	
1		State	
2		Postcode	
3		Country	
<input type="checkbox"/> Home phone	<input type="checkbox"/> Work phone	<input type="checkbox"/> Mobile	<input type="checkbox"/> Home email
<input type="checkbox"/> Home email	<input type="checkbox"/> Work email	Tick best contact method	
Best contact number or email address _____			
Diagnosis <small>See overleaf for # options</small>			
* Date diagnosed	*Bleeding disorder #		
/ /			
*Severity	Baseline factor date	Baseline factor level	*Weight in kilograms
	/ /	%	
<small>Mild / Moderate / Severe / Unknown / Not applicable</small>	<small>(Where applicable)</small>	<small>(Where applicable)</small>	
Treatment <small>See overleaf for + ^ options</small>			
*Regimen +	*Product name ^	*Total dose	*Frequency
Comments _____			
Attending Physician and Clinic / Hospital Address <small>Missing data will be requested by an ABDR Data Manager.</small>			
*Title	*First name	*Last name	
*Name of Clinic / Hospital		*Best contact number or email address	
*Address		*Suburb	
1		*State	
2		*Postcode	
3			
DECLARATION:			
These details are true and correct at the time of completing this form. I have read the ABDR User Conditions and the Clinicians FAQ on the ABDR and I understand my role and obligations in populating the ABDR. The patient is also aware of the purpose for capturing their details in the ABDR and is aware of privacy and confidentiality protection arrangements as described overleaf. The ABDR Pamphlet has been given to patient.			
Name _____	Signature _____	Date / /	

When complete fax to your nearest Treatment Centre or Clinic – see www.ahcdo.org.au for details.

Effective November 2016.

#Bleeding Disorder

Factor II deficiency (Prothrombin)
 Factor V deficiency
 Factor VII deficiency
 Factor VIII deficiency (Haemophilia A)
 Factor IX deficiency (Haemophilia B)
 Factor X deficiency
 Factor XI deficiency
 Factor XII deficiency
 Factor XIII deficiency
 Symptomatic Carrier Factor VIII deficiency (Haemophilia A)
 Symptomatic Carrier Factor IX deficiency (Haemophilia B)
 Asymptomatic Carrier Factor VIII deficiency (Haemophilia A)
 Asymptomatic Carrier Factor IX deficiency (Haemophilia B)
 von Willebrand Disease Type 1
 von Willebrand Disease Type 2 – Uncharacterised
 von Willebrand Disease Type 2A
 von Willebrand Disease Type 2B
 von Willebrand Disease Type 2M
 von Willebrand Disease Type 2N
 von Willebrand Disease Type 3
 von Willebrand Disease – Uncharacterised
 Fibrinogen – Afibrinogenemia
 Fibrinogen – Hypofibrinogenemia
 Fibrinogen – Dysfibrinogenemia
 Fibrinogen dysfunction – Uncharacterised
 Platelet – Glanzmann's thrombasthenia
 Platelet – Bernard-Soulier
 Platelet – May Hegglin
 Platelet – Macrothrombocytopenias
 Platelet – Storage pool (dense granule) deficiency
 Platelet – Primary secretion defect
 Platelet – Uncharacterised
 Acquired factor VIII inhibitor (Acquired Haemophilia A)
 Acquired von Willebrand's Disease
 Vascular disorders – Ehlers Danlos Syndrome
 Vascular disorders – Uncharacterised
 Other, please specify

+Treatment Regimen

On demand
 Prophylaxis
 Tolerisation
 Secondary Prophylaxis

^Product Name (Type)

Advate® (rFVIII)
 BeneFIX® (rFIX)
 Biostate® (pdFVIII)
 Ceprotin® (Protein C)
 Cryoprecipitate
 DDAVP (Synthetic hormone)
 Factor Eight Inhibitor Bypass Agent (FEIBA®) (Bypassing Agent)
 Factor VII Concentrate® (pdFVII)
 Factor XI bpl® (pdFXI)
 Factor XI LFB Hemoleven® (pdFXI)
 Fibrogammin P® (pdFXIII)
 Fresh Frozen Plasma (FFP)
 Haemocompletan P 1g (pdFXIII)
 Intravenous Immunoglobulin (IVIg)
 Kogenate (rFVIII)
 Kogenate FS – Blood Service (rFVIII)
 MonoFIX® - VF (pdFIX)
 NovoSeven® (rFVIIa)
 NovoSeven RT® (rFVIIa)
 Platelets
 Prothrombinex™ - VF (pdPCC)
 Recombinate® (rFVIII)
 ReFacto® (rFVIII)
 Xyntha (rFVIII)
 Xyntha Dual Chamber (rFVIII)

ABDR Patient Pamphlet

What is the ABDR? The Australian Bleeding Disorders Registry (ABDR) is a database that collects all clinical information related to the treatment of people with bleeding disorders, like an electronic medical file. This includes information about patient diagnosis, treatment details, hospital admissions and administrative information as well as details on ordering, supply and use of clotting factor products. Information is entered into the ABDR by staff at haemophilia treatment centres. The ABDR is managed by the National Blood Authority. The ABDR was first established in 1988 and has been upgraded many times with the latest significant upgrade in 2012.

Why do you need it? The ABDR provides your health care team and support staff with a record enabling them to monitor and manage your treatment over time to improve your quality of life. Depersonalised information available from the ABDR may be used by authorised organisations to understand and improve treatment for bleeding disorders. The ABDR also provides governments with information on total clotting factor product requirements to make sure there is enough available to meet the needs of all Australians with bleeding disorders.

What about privacy? Only the health care team and support staff involved in providing medical services to you have access to your personal information. Other authorised users only have access to limited, depersonalised and/or summary information where all identifying information is removed to protect your privacy.

Does information about me have to be included? A minimum amount of information about you is required to ensure the continuous supply of clotting factor product is available to meet your treatment needs.

Where can I get more information? Further information about the ABDR can be obtained from the Australian Haemophilia Centre Directors' Organisation (AHCDO) on (03) 9885 1777, email info@ahcdo.org.au or visit www.ahcdo.org.au

Endorsement from Haemophilia Foundation Australia

Haemophilia Foundation Australia supports the ABDR. It helps doctors and other treating health professionals to understand more about the care and treatment needs of people affected by bleeding disorders. The ABDR will assist and guide planning to ensure treatment product is available when it is needed. We are confident that the steps in place will mean accurate, reliable and confidential data is available and that no patient details can be identified outside haemophilia centres.

www.haemophilia.org.au

Endorsement from Australian Haemophilia Centre Directors' Organisation

The ABDR is a valuable tool that provides an overview of those affected with haemophilia and other bleeding disorders in Australia. Data from the ABDR is the best information available for clinicians to advise governments making policy decisions regarding treatment needs and product availability.

National statistics available through the ABDR will give AHCDO an overview of practise and allow opportunities for improvement. This data can be pooled to compare Australian treatment standards with international benchmarks. The ABDR will continue to provide the ability to assess quality of life and other important clinical questions arising across Australia.

AHCDO's partnership on this initiative with the National Blood Authority, Haemophilia Foundation Australia and other specialist health professional groups is vital to the pursuit of excellence in clinical treatment practices.

www.ahcdo.org.au

Copies of this pamphlet can be obtained by contacting the National Blood Authority at support@blood.gov.au or 13 000 BLOOD (13 000 25663)

Acronyms and glossary of terms

ACRONYMS

ABDR	Australian Bleeding Disorders Registry
AHCDO	Australian Haemophilia Centre Directors' Organisation
BU (BU/ml)	Bethesda unit (expressed as Bethesda units per millilitre)
DDAVP	Desmopressin (1-desamino-8-D-arginine vasopressin) a derivative of the antidiuretic hormone, used to treat patients with von Willebrand disease. It does not come under the national blood agreement funding arrangements and its use is often not recorded in the NBA's issues database.
FEIBA	Factor VIII Inhibitor Bypassing Activity
FVIIa / rFVIIa	Factor VIIa (seven 'a') / Recombinant Factor VIIa
FVIII / rFVIII	Factor VIII (eight) / Recombinant Factor VIII
HFA	Haemophilia Foundation Australia
HMA	Haemophilia A (Factor VIII deficiency)
HMB	Haemophilia B (Factor IX deficiency)
HTC	Haemophilia Treatment Centre – A specialist centre at certain hospitals where comprehensive care is undertaken for people with haemophilia. Non HTCs are other hospitals who are encouraged to work with HTCs in their region.
IDMS	The NBA's Integrated Data Management System
IU	International Units
MyABDR	a secure app for smartphones (Android and iOS) and a web site for people with bleeding disorders or parents/caregivers to record home treatments and bleeds.
NBA	National Blood Authority
OBD	Other bleeding disorders
PWH	People with Haemophilia
VWD	von Willebrand disease
WFH	World Federation of Hemophilia

GLOSSARY OF TERMS

bleeding disorders	Diseases that cause abnormal or exaggerated bleeding and poor blood clotting
blood products	Products manufactured from donated blood
fractionation	Blood plasma fractionation refers to the general processes of separating the various components of blood plasma